RECENT IMPLICATIONS OF STEM CELL IN POST-SURGICAL INTERVENTION

Sheetal Soni, Uditi Handa*, Sonali, Ujjwal Sanduja, Srishti Vats, Yash Gera, and Kumar Guarve
Guru Gobind Singh College of Pharmacy, Yamuna Nagar-135001, Haryana, India
Corresponding email ID: *uditipharmacist@gmail.com

ABSTRACT

Stem cells (SCs) are non-developed biotic cells that possess the capability of proliferation, self-renewal,conversationthedifferentiationofavarietyofcells,andregeneratingtissuemechanisms. The main objective of stem cells is that they are utilized in regenerative medicinesand tissue engineering processes. Stem cells are divided into various categories and the majorone is adult stem cells which consist of body organs and collagens stem cell categories. Stem cellalso plays a crucial role in clinical therapeutic practices mainly in surgical intervention and invariousorgandisorders.MesenchymeStemCells(MSCs) arealsousedinclinicalresearchanddevelopment processes. According to standard regulatory authorities like the United State FoodandDrugAdministration (USFDA)Stemcellscontributeanimportantroleincosmeticsmedications and are the first cell therapy that is used for the removal of wrinkles. There are various kinds of research also performed to find the SCs which improve heart functions and different cardiac cells'capabilities and also modify the muscle cells. They possessthesamepropertiesascancerandcan also undergospontaneousmalignanttransformations. It playsa crucial role in bone surgery and surgical operations with the help of engineering science and in avascular necrosis. The Different studies also showed that Stem Cell therapy is faster and increases the ligament-tendon healing process. Some researchers also indicate opposite outcomes that Mesenchyme Stem Cells can give increasing the healing of ligament-tendon injuries. It is noticed that the application of SCs therapy is for nerve cell reproduction, different nervous system injuries, and various nerve and muscle diseases. The mainly observed areas of aesthetic surgeries are gastric SCs which can be used for allogeneic fat transplantation. One of the experiments was performed on genetically altered mice and Bone Marrow Mesenchymal Stromal Cells (BM-MSCs) were carried out with the purpose of evaluating the utilization of SCs in diabetic wound therapy. One other experimental study on rats also expresses that the subcutaneous injections of allogeneic BM-MSCs reduce the apoptosis count and are necessary for the survival of the zone. Therefore,stemcellspossessbothbenefitsanddrawbacksinclinicaltherapeuticmanagement.

Keywords:*Stemcells,Mesenchyme Stem Cells,Clinicaltherapeuticpractices,Regulatoryauthority,Malignanttransformation.*

DOI Number: 10.14704/NQ.2022.20.13.NQ88132 **NeuroQuantology2022;20(13): 1042-1052**

INTRODUCTION

Theterm"stemness"definesasequencethatdifferentiatesthewide-rangingdiversificationofthe cell community. The SCs are the non-developed biotic cells that possess the capability ofproliferation,self-renewal,andconversationthedifferentiationofavarietyofcells,andregenerating tissue mechanisms. The SCs in their early stages of the 5-7 days embryo is only totipotentand are not eligiblefortherapypurposesduetomoralreasons (teratoma formation).Thepluripotentialcellscanbe developed further into cells that contain three types of germ layers that isendoblast,ectoblast,andmesoblast.Thecellsaftertheendphaseofafetusaftertheprocessoftheblastocyst are the pluripotential cells. The multipotential cells are the cells that are developed intomore than one stage. Adult SCs are the umbilical cord blood cells andare examples ofmultipotential cells as illustrated in **Figure 1** [1,2,5]. The totipotent cells have a potency very high (single cell embryo) and can be differentiated into an entire organism, resulting after the fertilization process which will be extracted from the embryo and also the placenta which will be easily isolated but have some ethical issues whereas another variety based

upon the potency of stem cells is Unipotent which only produces one cell type, but have the properties of self -renewal or regenerated power whichdistinguishthemfromnon-stemcells[6,7].



Figure 1: Illustration of sites for stem cell extraction.

Classification of SCs according to the Sources:

FetalStemCells: Theembryonicstemcellsareachieved bytheblastodermicvesiclestageof the fetus. The pluripotential has the actual capability to generate different tissues in the bodyand divided them in an extensive way. The process is happened due to the criteria of asymmetricalseparation which generate one type of stem cell and another typeofdaughtercell [2].

Postnatal Stem Cells: The adult SCs are achieved by the end stage of life furthering theprocessof the fetal stage. They are basically the type of multipotential, alike cells localized withspecial tissues having the primitive objective of preservation and renovation. The epithelial SCsthat are generated by the mesoblast is the part of adult SCs that is having the good capability toproduce the stagesof adipocerous, chondroblast, myoblastoma, and osteoblastoma.

MesenchymeStemCells:ThemesenchymeSCsareloc atedpost-natalinsidethenon-hematopoietic bone marrow stromal. The essence of body tissues is the buildup of a diversifiedcommunity of cells that contains latticed cells, fatty, bone-forming tissue cells, pronators cells,epithelium cells, and scavenger cells [3].

Hepatic Stem Cells: The individual is called to be alive by the surgical removal of seventy-five percent part of the hepatic organ by reproducing technique. Real tissue further restores for 2to 3 weeks. That is different for other organs like as the renal and pancreatic systems. The newproofsadvisethatdifferentvarietyofcellsandproc essesthatisresponsiblefororganregeneration depends upon the type ofhepaticinjury.

Pancreatic Stem Cells: There has been the contention that the pancreatic system carries realSCs. This is also declared that the secret or cells in rats' pancreas islets of Langerhans containinginsulin generating b-cells, yield gives after every forty to fifty days from the procedure ofnecrobiosis and multiplication and distinction of the new islet cells by originator epithelium cellsfound inside the pancreatic ducts.

Abdomen Stem Cells: Abdomen stem cells, a group of adult stem cells residing in thestomach, play a key role in maintaining the dynamic homeostasis of the gastric epithelium [4]. Recent studies implied that Gastric Cancer Stem Cells (GCSEs) play an important role inthe initiation and progression of Gastric Cells (GC). This subpopulation comprises cells withseveral features, such as self-renewal capability, high proliferating rate, and ability to modifytheir metabolic program, which allow them to resist current anticancer therapies [5].

Epiblast Stem Cells: The human skin is categorized into the outer epidermis and underlyingdermis.Hairsandsebaceousglandscanalso makeupthepartoftheepidermis.Themost important cell part in the epidermis is the keratinocyte which is the epithelial cell that isdivided into the basal layer of the epidermis. Once these cells leave the basal areas, they furtherundergo the terminal or final differentiation that gives an outcome in a particular cell known asQuami which basically makes the follicle layers that makes an outer skin layer inside the aliveskin cells [6].

Osteogenic and Collagens Stem Cells: The mesenchyme SCs inside the bone center canlead to differentiation into bones and collagens. The bone itself had been founded to have boththwoprocessesthatareuncommittedstemcellsan dcommittedosteoprogenitorcells[7].

Neural Stem Cells: It is also advised that a continuous neurogenic yield happens in thelimited locations of the peripheral nervous system. These two peripheral areas are eye SCs ofthe adult mammalian nervous system said to be included in this process the subventricular zoneoftheforebrainandtheelevationinthesurfaceoft hebrainofthehippocampus whichis basically called a reservoir forneural cells [8].

Ophthalmic Stem Cells: Eye diseases that cause progressive blindness include both age-related and inherited retinal diseases as well as corneal diseases. According to recent studies,therapeutic stem cell approaches for dry age-related macular degeneration, retinitis pigmentosa,and limbal stem cell deficiency [9].

Clinical Involvements of Stem cell Therapy:

Cardiothoracic Surgery: In cardiovascular diseases like heart attack is the major reason fordeath rates overall areas. Heart attacks happening to atherosclerosis can create changes and theoxygen supply damage to the heart organ. The oxygen supply damage gives causes the loss ofcardiac cells the reason for cell death. The different types of studies expressthevarietyofcellpartscontainingfetal,produc epluripotential,mesenchyme,andcardiacdiscovered SCs. There are lots of parameters selected while selecting a real cell for exampletoenhance heart functions the cardiac cells have the capability to modify into muscle cells, vascularendothelium cells, and myocytes act like paracrineprocedures, all the clinical trials based on SCs are elaborated in **Figure 2** [10].



Figure 2: Mind map of Clinical trials of stem cell therapy.

Fetal Pluripotential Stem Cells*:* The fetal SCs provide all cell types that are located in theorganisms. Although there are also many types of moral reasons for the use of fetal SCs andthelimitedsupplement oftransplantationofhumanfetuses.Theprocesscause sthe growth ofinduced pluripotential SCs. pluripotential SCs were the first program provided by the individualmice by the procedure of different kinds of steps, containing different cultures.

Mesenchyme Stem Cells: The mesenchyme SCs located in the bones center, connectivetissues, navel string blood, and reproductive structure. They have the capability to differentiate tothe ancestry like as mature bone cells, cartilage cells, fatty cells, muscle cells, and center stromal[11]. The cells are predicted to produce dissolved development factors andlymphokine which are further used in secret or cells and paracrine fashion model and provide their therapeutic action [12].

Bone Marrow: The clinical trials include the intracoronary route of autogenous bones centercomes from the mesenchyme SCs in treating individuals who take transdermal heart muscleinvention for the acute step of STEMIs at the end time of six months, the epidemiology of leftventricular ejection fraction enhanced from 6.7 percentin the therapy members individualsversus 0.7 percent in the treating member individuals [13].

Fatty Tissues: The mesenchyme SC s are also separated with the help of adipose tissue,particularlybythehelpofstromafragmentsofth efattytissues.Thistypeofcellisgraceful becauseoftheeasyprocessanditsremovalpartbytheu seoflipectomytubes[14].

Themesenchymestromacellstreatmentinindividuals sufferingfromchronicheartattack is done having the phase two study by use of fatty discovered cells that were responded byVEGF-A165, to determine the therapeutic effect of treating the group having the serious heart attackand angina pectoris [15].

Cardiac Stem Cells: These kinds of cells were located to indicate the attributes of bothtypes of mesenchyme SCs (Thy-1 and Endoglin), fetal SCs, and also beginning attribution of thecardiogenic process [16].

A lot of clinical practices estimating the security and impassiveness of coronary artery selectedthe prominin-1 in the heart recovery process after the acute heart attack and left ventricular heartwallmalfunctioningisestablishedtotheANZCTR(IDACTRN12609001045202,CHUM,Quebec,Canada)

eISSN 1303-5150

whichisrecentlygivendifferentstepsasgivenbelow(**Table1**) [17].

Table 1: Different steps of clinical practices by using different sources of stem cells

Systematic Review	Surgery/ methods	Different sourcesof stem cells	Results
Treatment of IHD and heart failure			
Ali-Hassan Sayeghet al. 2015	InborncoronaryanomalygraftingofSCsisinthecoronary artery bypass surgery	Bone marrow	It enhances the cardiac functionalparameters that basically increasetheleftventricularejectionfraction withinthenon-notabledeductiontotheleftventricularend-systolicvolume.Thetreatmentprocedurehasnolifecomplications problems.
Qin et al. 2015	Grafting of SCs includescoronaryartery bypasssurgery to the therapy of ischemic heartdiseases.	Bone marrow	It gives significant enhancementtotheLVEFandencouragessupportinleft ventricular remodeling.
treatment of lung disease and acute respiratory distress syndrome (ARDS)			
Mao et al. 2013	SCsarethemanagementofacutelunginjuryandacuterespiratory distresssyndrome (ARDS)	Toimprovetheangiogenesisofdamaged vessels	It has a management potential forvascularreconstitutionandalsohasa novelpolicyintreatingacutelungdisease andacuterespiratory distresssyndrome (ARDS)
Tissue-engineer cardiothoracic substance and organ table			
Lim et al. 2013	Tissue-engineeringofthoracictissues and organ	Hematopoieticand MSC	Livingproceduresfillcompleteimportant needs. Although, mainenhancements are still pending.

OrthopedicSurgery: TheSCscanbemainlyusedforinjurieseithersurgicallyorbyinjection with the help of a trained carrier. MSCs are taken by or through starting phases ofdifferentiation, making bones and cartilage precursors under the given laboratory condition andthenimplantingthemintoinjuries.MSCscanalsobegivenbytheI/Vroute.Thecapabilitytochange according to ranking and multiplies in the central bones after the therapy of the peripheralroute isrecovered from the management of Osteogenesis Imperfecta [18].

Spine and Peripheral Nerve Surgery: The prospective study of spine fusion indicates thatnot selecting the SCs that are used within the hypertonia arterials has the same treatment criteriain rats as auto transplantation grafting, therefore,itpreventsthedonorstem orbidity11. Thesame outcomes were achieved by Gan et al using β –TCP [19].

Osteoarthritis (OA): because of their role in stopping the process of breaking the activity ofsubstance metalloprotease, mesenchyme SCs are providing crucial effects in OA [20]. In the advanced study, researchers indicate that guinea pigs within the age regardingosteoarthritis given the treatment with mesenchyme SCs matrix injection have a greater option toreconstitute by greater type two

collagens and fewer metalloprotease contents [21].

HighestShinBoneSurgeryandSurgicalOperation:By exactusesofengineeringscience for generating normal planted surface, mesenchyme SCs have a greater capability fortransforming joint exchangesurgical process from smoothosteo integration procedure.Thethree-dimensional scaffolds prevalence used in the data with MSCs is can be used in furtherautologousosteochondralgraftsthatcanplaya nimportantroleinbiologicarthroplasty[22].

Avascular Necrosis: It is a disorder occurring mostly in younger individual patients. Thisis defined by the minimum blood supplement process in the bone marrow that the processattaches to this and further enhances the Intracoastal potential. The intensity of the beneathcartilage plates is a very crucial defining factor within the head including (central slumbering,bone grafting, thighbone osteotomy) and head compromising (hip resurfacing) methods. SCs werefurther separated and used for the study by researcher Rastogi where sixty hips in the beginningstages of AVN have been random to be treated within central slumbering and with central boneinjections [23].

Ligament-Tendon Injuries: Most of the evidence expresses that stem cell therapy is fasterandenhancesligament-tendonhealing.Although,someliteraturehasopposit eoutcomes and states that MSCs can give enhancing the healing of ligament-tendon injuries.Although, the lower doses of MCSs are very efficient in enhancing healing when the result isinterpreted withhigher doses [24].

RotatorCuLesions:These kindsoflesionsarebasicallyconnectedwithmusclepat hology, likeweaknessorimpingement.Therupturedinjuriesar esuccessfullytreatedwithopenor arthroscopictechniquesAlthough,re-ruptureratesarestillveryhightreated with infraspinatus injuries of the rats with polyglycolic acid which is combined withBM-MCSs orisolated polyglycolic acid [25].

Otolaryngology-Head and Neck Surgery: This research recognized the reviews that areespeciallynewtoOtolaryngology-HeadandNeckSurgery. Noadditional RCTsorcomparativetrialswerefurtherrecognized.Th eseclinicalpracticesare shown in **Table No. 2** [26, 27].

Table No.2: Different types of surgeries by using different stem cells

Systematicreview	Surgery	Source of stem cell	Results
Hamilton etal.2014	Regenerative medicine ofmedical nasal airway	MSC and epithelial cells.	Thefullepithelialproceduretoendoscopy(15months),ciliatedepitheliaoncytology,andnormal perfusion scan were observed thatdepend upon the mammal studies.
Wormald etal.2015	Regenerative surgery	ASC,ESC, iPSC, bone marrow-derived MSC,amniotic fluid-derived SCs	Outcomesbasicallydependonanimalsandoutsidethebodystudy.

Neurosurgery: There are lots of branded cells searched by the adult tissue containingNeuralstem-ad for treatment of brain disease, SB623, Multistem (Multipotential originator Cell,ALD-401, and PDA001 for the patients of stroke. Of the variety of SCs used in clinical research,not any is having actual use in the treatment till now [28].

InNewZealandandAustralia,itisobservedthattheuse

eISSN 1303-5150

ofSCstherapyforthecerebraldisorder,nervecellreproduction,peripheralnervoussysteminjury,andnerveandmusclediseases[29,30,31].Arelatedstudyanalyzedtheimpassiveness of SCs for traumatic mental injury is givenin **Table 3** [32].

Table No.3:Evaluation of the impassiveness of SCs for the traumatic mental injury

Study	Surgery method	Source of stem cell	Study result
Cerebral disorder			
Novaket al. 2016	Not reported	Hearingnerves,brainnerves,neuraloriginators,andallogeneic,umbilicalcord blood stem cells	The SCs are taking in the important therapy foreffectivetreatmentforthecerebraldisorderandnotany evidence is right now.
Face nerve reconstitution			
Euler et al. 2014	Not reported	BonemarrowMSC,ASC, dental pulp cells,and neural SCs	TheSCsdiscoveredbythevarioussourcesofpresentoutcomesconnectedtofacenervereconstitutions and generates impactive functionalresults (related to animal research)
Neuromuscular diseases			
Finsterer andZarrouk Mahjoub2016	Not reported	Hematopoietic stemcells (autologous)	thisindicatestheincompletesolutionofg.i.t, muscle and brain malfunctioning to theneuronsand muscles disorder.
Mental injury			
Wang et al. 2013	Not reported	Placental vessels,mesenchyme SCs	Research including the multicenter and big samplefor the random clinical research, which is neededandexplainingtheroleofplacentalvesselsmesenchymeSCsgraftingofvarioustraumaticmental injuries.

AestheticSurgicalprocess: Themostlyobservedlocationsofaestheticsurgeriesareabdomen SCswhichcanbeappliedinallogeneicfattransplantation.allogeneicfat transplantation right now becomes famous for the delicate tissue combination process in theentirebody [33].

Cell Therapy in Cosmetic Surgical Process: There are the uses of stem cells in cosmeticmedications.InJune2011,aftertheperiodof 10yearsperiodofliterature,theUSFoodandDrug Administration (USFDA) approved laViv, the first cell therapy for the removal of wrinklessurrounding the nasal sides and mouth. The laViv therapy involves harvesting a patient's ownfibrinogenfor up to90 days and then further again injecting the injection into the skin duringaset period of therapy [34].

Hereditary Risks of Cell Processing: Potential spoiling and diminishing of cells make anissue when cell-related products go to damage continuously greater than normal malfunctioning,involving cell expansion in culture and distinctions [35]. The cells are eliminated by apatientandchangesinthesimilarsurgeryprocedures teps are notathigherriskofdiseasetransmission in comparison to the surgical process. Although cell culture basically includes theuse of nonhuman serum, which is achieved by the fetal calves, so defining a potential risk forprion infection [36].

eISSN 1303-5150

Regenerative Surgical Process: Allogeneic adipose grafting whose resorption rate exceedsaround 70%, is used for the correction of facial symmetry and as filler after cancer treatment, andaftertheresultofradiotherapy.Otherproductstha tcanbeappliedforthesepurposeslikexenogeneic collagen, biopolymers, and cellular dermal matrix that can develop immunologicalreactions at the recipient site. These kinds of features of the ideal filler for soft tissues are that itmust be easy to use, effective, and low-cost and that it must provide long-lasting therapeuticeffects without causing any immunological reaction as shown in Figure 3.

Treatment of Chronic Wounds and Burn: An experiment was performed on geneticallymodified mice and BM-MSCs were carried out with the objective of estimating the usefulness ofstem cells in diabetic wound therapy [37]. An experimental study on rats also indicatesthatsubcutaneousinjectionsofallogeneicB M-MSCsreducetheapoptosiscountandarenecessary for the survival of the zone. The drawback of this procedure is the of their importanceof administering the stem cells very soon after the injury [38].



Figure 3: Regenerative Surgical Process by stem cell therapy.

CONCLUSION

This study showed the basic properties of stem cells and their sources for the use of differentbodyfunctionslikehepatic,pancreatic,abdo men,andmanyothers.Thedifferenttypesofclinicalinv olvementsandlatesttherapieswillhelpthepatient'sth erapeuticmanagement like removal of wrinkles in cosmetics, enhancing heart functions and modifying muscle cells activity, bone surgery, malignant tumors, tendon ligament injuries, anesthetics surgery, Diabetes Mellitus, andwell as to do the research to generate the latest techniques and therapies of stem cells.

ACKNOWLEDGMENT

The authors of the Department of Pharmacy, Guru Gobind College of Pharmacy, Yamunanagar (Haryana), India are gratefully acknowledged for their support and space.

CONFLICTS OF INTEREST

The authors' declaration of conflict of interest is none.

FUNDING RESOURCES

This short communication manuscript received no funding.

REFERENCES

1. Shostak S, (Re) defining stem cells, 2006, 28, 301-8. doi: 10.46582/jsrm.1401003
2. Shahriyari L, Komarova NL. Symmetric vs. asymmetric stem cell divisions: an adaptation against cancer? 2013. doi: 10.1371/journal.pone.0076195
3. Bianco C, Riminucci M. The bone marrow stroma in vivo: Ontogeny structure, cellular composition and changes in disease; In Marrow Stromal Cell Culture. In J. Bresdord and M. Owen(Eds). 2010. doi:10.1017/CBO9780511623219.004
4. Xiao S, Zhou L. Gastric Stem Cells: Physiological and Pathological Perspectives. Front. Cell Dev Biol, 2020. doi: 10.3389/fcell.2020.571536
5. Addeo M, Paola GD, Verma HK, Laurino S, Russi S. Gastric Cancer Stem Cells: A Glimpse on Metabolic Reprogramming. Front. Oncol. 2021. doi.org/10.3389/fonc.2021.698394
6. Blanpain C, WE Lowry, Geohegan A. Self-renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche, 2004, 530–32. https://doi.org/10.1016/j.cell.2004.08.012
7. Gronthos S, Franklin DM, Leddy HA, Surface protein characterization of human adipose tissue-derived stromal cells; Journal of cellular physiology, 2011, 54–63. doi: 10.1002/jcp.1138
8. Lois C, Alvarez A. Proliferating sub ventricular zone cells in the adult mammalian fore brain can differentiate into neurons and glia, 1993, 2074-77 doi: 10.1073/pnas.90.5.2074
9. Caras IW, Collins LR, Creasey AA. A stem cell journey in ophthalmology: From the bench to the clinic. Stem cells journals, 2021,10(12):1581-87. doi.org/10.1002/sctm.21-0239
10. Heyndrickx GB, Baig H, Nellens P, Leusen I, Depression of regional blood flow and wall thickening after brief coronary occlusions; Am Physio l Soc, 1978, 234(6), 653–59. doi: 10.1152/ajpheart
11. Pittenger MF, Martin MJ. Mesenchymal stem cells and their potential as cardiac therapeutics. Circ Res, 2004,95(1). doi: 10.1161/01.RES.0000135902.99383.6f.
12. Monsel A, Zhu YG, Gennal S, Preclinical evidence and ongoing clinical trials using mesenchymal stem cells; Anesthesiology, 2015, 121(5):1099–1121. DOI: 10.1097/ALN.0000000000000446
13. Wollert KM, Meyer GP, Lotz J, SR Lichtenberg, Intracoronary autologous bone-marrow cell transfer after myocardial infarction, the BOOST randomised controlled clinical trial, 2004, 364(9429). doi: 10.1016/S0140-6736(04)16626-9
14. Casteilla L, Benard VP, Laharrague P. Adipose derived stromal cells: their identity and uses in clinical trials; an update. World J Stem Cells, 2011, 3(4), 25–33. https://doi.org/10.1016/j.cell.2004.08.012
15. Qayyum AHS, Sørensen MH. Adipose-derived mesenchymal stromal cells for chronic myocardial ischemia (My Stromal Cell Trial): study design; Regen Med, 2012, 7(3):421–28. doi: 10.2217/rme.12.17.
16. Chong JJ, Reinecke H, Torok-Storb B. Progenitor cells identified by PDGFR-alpha expression in the developing and diseased human heart; Stem Cells Dev, 2013, 22(13),1932–43. DOI:10.1089/scd.2012.0542
17. Yasoba, Atukorale, Lambert R, A Cameron of ASERNIP-S; Royal Australasian College of Surgeons
18. Horwitz EM, Prockop DJ, Gordon PL, Clinical responses to bone marrow transplantation

in children with severe osteogenesis imperfect; Blood, 2011, 1227-31 DOI: 10.1182/blood. v97.5.1227

19. Gan Y, Dai K, Zhang P. The clinical use of enriched bone marrow stem cells combined with porous beta tri calcium phosphate in posterior spinal fusion; Biomaterials, 2008, 3973-82, doi: 10.1016/j.biomaterials.2008.06.026
20. Lozito TP, Tuan RS. Mesenchymal stem cells inhibit both endogenous and exogenous MMPs via secreted TIMPs; J Cell Physiol. 2011, 226, 385-96. DOI: 10.1002/jcp.22344
21. Sato M, Uchida K, Nakajima H, Direct transplantation of mesenchymal stem cells into the knee joints of Hartley strain guinea pigs with spontaneous osteoarthritis; Arthritis Res Ther. 2012. doi: 10.1186/ar3735.
22. Maclaine SE, McNamara LE, Bennett AJ. Developments in stem cells: implications for future joint replacements; Pro cIn Mech Eng H, 2013, DOI: 10.1177/0954411912471492
23. Rastogi S, Sankineani SR, Nag HL, Intralesional autologous mesenchymal stem cells in management of osteonecrosis of femur: a preliminary study; Musculoskelet Surg, 2013, 223-8 doi: 10.1007/s12306-013-0273-0..
24. Saether EE, Chamberlain CS, Leiferman EM, Primed Mesenchymal Stem Cells Alter and Improve Rat Medial Collateral Ligament Healing. Stem Cell Reviews and Reports; Stem Cell Rev, 2015, 10(1),1–12. doi: 10.1007/s12015-013-9479-7. [PubMed: 24174129].
25. Yokoya S, Mochizuki Y,Natsu K, Regeneration using a bio absorbable material with bone marrow derived mesenchymal stem cells in a rabbit model; Am J Sports Med, 2012, 40(6):1259–68. doi: 10.1177/0363546512442343
26. Hamilton N, Bullock AJ, Macneil S, Tissue engineering airway mucosa. A systematic review; Laryngoscope, 2014, 124(4), 961-8, doi: 10.1002/lary.24469
27. Wormald JC, Fishman JM, Juniat S. Regenerative medicine in otorhinolaryngology, J laryn otol, 2015, 129(8), 732-9. doi: https://doi.org/10.1136/bmj.m718
28. Mack GS. ReNeuron and stemcells get green light for neural stem cell trials; Nature Biotechnology, 2005, 29, 95–7 DOI: 10.1038/nbt0211-95
29. Finsterer J, Mahjoub SZ. Treatment of muscle weakness in neuromuscular disorders; Expert review of neurotherapeutics, 2016, doi: 10.1080/14737175.2016.1206471
30. Novak I, Walker K, Hunt RW. Concise Review: Stem Cell Interventions for People with Cerebral Palsy: Systematic Review with Meta-Analysis; Stem Cells Translational Medicine, 2016. DOI: 10.5966/sctm.2015-0372
31. Lucena EDS, Guzen EFP, Lopes D. Experimental considerations concerning the use of stem cells and tissue engineering for facial nerve regeneration: A systematic review; J Oral Maxillofac Surg. 72(5),1001-12, DOI: 10.1016/j.joms.2013.11.006
32. Wang S, Cheng H, Dai G, Umbilical cord mesenchymal stem cell transplantation significantly improves neurological function in patients with sequelae of traumatic brain injury; Brain Research, 2013, 1532, 76-84. DOI: 10.1016/j.brainres.2013.08.001
33. Coleman SR, Saboeiro AP. Fat grafting to the breast revisited: safety and efficacy; Plast Reconstr Surg, 2014, 775–85, doi: 10.1097/01.prs.0000252001. 59162.c9.
34. Schmidt C, FDA approves first cell therapy for wrinkle-free visage; Nature Biotechnology, 2011, 674–75. doi: 10.1038/nbt0811-674

35. Halme DG, Kessler DA, FDA regulation of stem-cell-based therapies; The New England journal of medicine, 2006,1730–35, doi: 10.1056/NEJMhpr063086
36. Fekete N, Rojewski MT, Furst D. GMP-compliant isolation and large-scale expansion of bone marrow-derived MSC, 2012, doi: 10.1371/journal.pone.0043255.
37. Wu Y, Chen L, Scott PG. Mesenchymal stem cells enhance wound healing through differentiation and angiogenesis; Stem Cells, 2007, 2648–59. doi: 10.1634/stemcells.2007-0226.
38. Öksüz S, Ülkür E, Öncül O. The effect of subcutaneous mesenchymal stem cell injection on statis zone and apoptosis in an experimental burn model; Plast Reconstr Surg, 2013, 463–71. doi: 10.1097/PRS.0b013e31827c6d6