Employing Clinical Dynamics to Reduce Head CT Scan Ordering for Trifling Cranium Trauma Patients

Abdullah Hasan Jabbar[1*], Murtadha M-Hussein A-Kadhim[2], Karrar Mahdi Badi[3], Salim Oudah Mezan[4], Muhammad Hasnain Jameel[5*], Abdulkareem Thjeel Jabbar[6]

Abstract

The current study is to figure out how routine care procedures could help prevent patients who have suffered mild head injuries from needing a CT scan. Patients who are awake (Al-Hussein teaching hospital in Al-Muthanna Province- Iraq) were brought to the emergency room unconscious due to serious head injury and memory loss. The CT scan was then done to assess any head predictors. Extreme headache sufferers, fatigue, and vomiting Physical examination revealed a depressed skull fracture. were checked for potential risk factors. Patients who have there are no known risk factors and who present with an irregular CT rate, along with those who have a risk factor or more, and a high CT rate, are compared with relation to their rate of surgical operation for a head injury. Of the 50 patients who were part of the study, 72% had no head injury risk factor, while 28% had at least one risk factor. If the total number of CT scans down, 16 (9%) showed an odd outcome. In this category, 3.7% had no risk factors, and 11% had one or more risk factors. It was not considered clinically significant that the CT scan anomalies were found in the low-risk community. Nine patients, which had at least one of the risk variables, needed surgery. According to the "minimal four-criteria" technique, over 72% of minor head trauma patients may be detected without a CT scan. They are identified with the aid of neurosurgery among individuals who had an abnormal CT scan. National savings inpatient charges may be a result of this solution. More research may support these findings.

Key Words: Clinical Dynamics, Head CT Scan, Cranium Trauma.

DOI Number: 10.14704/nq.2022.20.6.NQ22110 **NeuroQuantology 2022; 20(6):1161-1167**

Introduction

Based on the current medical literature, there is also uncertainty surrounding patients who have had a concussion and should be treated CT scanning in the emergency room (ED) [1]. It is fuelled by the demand to quickly locate any wounded patient, against enforcing rising standards to minimize rising treatment costs. Before this work was completed, retrospective research and previous clinical experience of patients who had sustained confusion following a head injury showed the importance of a CT scan of the head to rule out a neurosurgical lesion. Recent findings have shown that 20-35s The CT scans of individuals with a GCS of 13 or 14 are irregular. outcomes and it offers an inference that treatments are needed for many of these patients. Following head injuries, the workup of patients with a typical GCS and no neurological results is much less established. Head CT scans should be done on all patients who lose consciousness (LOC) or have amnesia following head trauma [2].

Corresponding author: Abdullah Hasan Jabbar & Muhammad Hasnain Jameel

Address: [1*]Optical Department, College of Health and Medical Technology, Sawa University, Ministry of Higher Education and Scientific Research, Al-Muthanna, Samawah, Iraq & Al-Hussein Teaching Hospital, Directorate of Al-Muthanna Health, Ministry of Health, Iraq; [2]Department of Biotechnology, College of Biotechnology, AL-Qasim Green University, Ministry of Higher Education and Scientific Research, Iraq; [3]Optical Department, College of Health and Medical Technology, Sawa University, Ministry of Higher Education and Scientific Research, Al-Muthanna, Samawah, Iraq; [4,6]General Directorate of Education, Al-Muthanna Governorate, Ministry of Education, Republic of Iraq; [5*]Shaanxi Key Laboratory for Theoretical Physics Frontiers, Institute of Modern Physics, Northwest University, Xi'an, China.

E-mail: [1*]abdullah.hasan.j@sawa-un.edu.iq; [5*]mhasnainjamil@gmail.com

Most individuals in this group need little or no care, and the anomalies found on CT scans are often mild. There is now hope that a less restrictive solution to addressing problems will be adopted.
It is found that among Patients who have had a mild to severe head injury and have a GCS of greater than minor than IS constituted the overwhelming majority who arrived in the ED, and these patients could dramatically cut hospital costs (2,6.X). Previous research has shown that around 80% of the patients who had CT scans after mild head injuries were classified by the four clinical risk factors [3]. To figure out whether or not we could confidently choose which GCS patients to treat. Of 15 traumatic brain injury (TBI) necessitates a CT scan in the emergency department., we investigated these risk factors in the current research. In addition to that, we sought to quantify the benefits that could be accrued in treatment expenses by our therapeutic requirements.

Materials and Methods

A large urban Level 1 trauma centre was used for the research, which was performed from October 2020 to February 2021.

Patient Selection

Inclusive Criteria

1. Adults aged 18 and up are the first group.
2. This study included patients with a history of a car accident, a fall, or an assault.

Exclusion Criteria

1. Cases of children.
2. Injuries to the skin.

Equipment Used

A Single Slice Spiral is used to scan patients: The investigation was conducted using GE CT/e equipment. It's a fifth-generation scanner that's been tweaked.

Slice Thickness – 1mm, 2 mm, 5 mm and 10 mm Available gantry tilt ± 20 degree

Matrix size of 512

KV – 80 to 130 MAS – 60 to 200

Figure 1. CT Scan [4]

CT Technique

All of the patients had their heads immobilized and positioned correctly. IV Diazepam was given to uncooperative patients (5-20 mg). To align the scan plane with the orbitomeatal line, the gantry tilt was set between 0 and 20 degrees. Brain and bone window settings were used to examine the pictures produced. Six to eight hours passed on average between the scan and the head injury. Patients were assessed using the proforma provided [5].
Our trauma centre mandated all those patients who experienced a lack of consciousness (LOC) or after a head injury, even if you have amnesia, you should have a head CT scan. Though they were conscious in the emergency room. In this situation, the word LOC meant the witness found the patient unconscious and told the ambulance crew or doctor in charge of the patient's treatment.
According to the concept, a patient would be called amnesic if they could not recall or explain the event contributing to their head injuries. In all patients with a healthy mental state, state, who presented less than 24 hours following a blunt head wound, amnesia or a history of LOC, and who had no symptoms of a current or deteriorating brain injury, admission was allowed. Patients who have used medications or alcohol, have undergone multiple system abuse, have a bleeding condition, or have an apparent coagulopathy are not disqualified. As long as the psychiatric state is stable and the GCS is 15, they will be admitted as patients.
Patients were registered on an ongoing basis, one at a time, and consecutively. Patients who came to the ED within two hours of suffering an injury must be included in the report. Both patients were CT-scanned in 8 hours after they arrived in the ED. The attending physician or senior emergency resident in charge of the patient's care. Treatment

completed a report sheet on each patient before the CT scan. Once patients appeared in the emergency room, their mental health is normal state was described as a GCS. Extreme headache was characterised as being more severe than other kinds of headaches when asking the patient questions. Both patients were asked whether they felt nauseated and responded using a "yes" or "no" response. "Vomiting" was recorded whether it happened before the emergency room entrance or in the ED. On evaluation by a resident or attending physician, a palpable deformity in the outer table of the skull was identified as a sign of skull depression.

The final reading taken by the visiting neuroradiologist was used as the official result. When multiple radiologists analysed There was no attempt to check inter-rater reliability using the CT images. Acute inflammatory intracranial lesions (subarachnoid haemorrhage, parenchymal hematoma, epidural hematoma, subdural hematoma) or a skull fracture is two features of an irregular CT scan. Patients who do not have all of the risk factors assessed for concussion are contrasted to patients who have any of these risk factors and who may need neurosurgical intervention. Neurosurgical intervention is repairing an acute traumatic injury or inserting a tracking bolt, craniotomy. Patients with irregular CT scans were monitored before hospital discharge to assess if an improvement in the mental state occurred.

Statistical Analysis

All of the data was gathered and translated to percentages as needed. To compare CT findings of different variables and parameters, the Chi-square test and Spearman's correlation coefficient were employed, and the p-value was computed using the MINITAB (USA 13.1) program.

Results

Over two months, 50 people have enrolled in the study. 6.4% of the CT scans were irregular, but only 0.2% needed neurosurgical intervention. The four main risk factors are measured in patients with an irregular and normal CT scan. The four risk factors for developing CML are included in the diagnostic criteria for CML. Using these diagnostic criteria on the patients in the sample showed that 36 (72% of the total group) had no risk factors, while 14 (28% of the total group) had at least one risk factor.

Table 1. Age distribution in 50 patients of head injury

Age Group (Years)	Number of Patients	Percentage	Neuro deficit & Death	
			Number	Percentage
Adult (18-50)	37	74%	26	68.42%
Elderly (>50)	13	26%	12	31.58%
Total	**50**	**100%**	**38**	**100%**
18-30	19	38%	16	42.11%
31 -50	16	32%	10	26.32%
>50	15	30%	12	31.58%
Total	**100**	**100%**	**38**	**100%**



Figure 2. Number of Patients



Figure 3. Number of Neuro deficit & Death

Table 2. Sex distribution in 50 patients of head injury

Age Group (Years)	Male	Percentage	Female	Percentage
Adult (18-50)	27	67.5%	7	70%
Elderly (>50)	13	32.5%	3	30%
Total	**40**	**100%**	**10**	**100%**
18-30	21	52.5%	5	50%
31-50	14	35%	3	30%
>50	5	12.5%	2	20%
Total	**40**	**100%**	**10**	**100%**



Figure 4. Sex distribution in 50 patients of head injury

Table 3. Association of fractures with various lesions

Fractures	Number of Patients
Isolated fracture	3
Fracture associated with other lesions	20
Pneumocephalus	12
EDH	6
Contusions	4
SDH	2
Intracerebral haemorrhage	3

Three patients had isolated fractures, whereas the other 78 had lesions associated with them. Contusion was the most frequent lesion, with 40 instances reported. There were 26 instances of pneumocephalus.

Table 4. Fracture detection with conventional skull radiograph and CT

Imaging Modality	Number of Patients	Percentage
Fractures detected only on CT	41	82%
Fractures detected only on plain radiograph	0	0%
Fractures detected on both modalities	2	2%

Only two instances had plain radiographs as the primary modality of inquiry, which CT validated. In 80% of the patients, CT revealed fractures.

Table 5. Extradural Haemorrhage: Relationship with Overlying Fractures

Item	Number of Patients	Percentage
EDH with fracture	15	93.75%
EDH without fracture	1	6.25%
Total	**16**	**100%**

EDH was associated with an overlying fracture in 93.75% of cases. This is a well-established association.

Table 6. Density patterns of SDH

Density Pattern	Number of Patients
Hyperdense	26
Isodense	1
Hypodense	-
Mixed	-
Layering	1

Acute SDH was predominantly hyperdense.

Table 7. Distribution of SDH

Sites of SDH	Number of Patients
Frontal	2
Temporal	2
Parietal	3
Occipital	1
Fronto-parietal	2
Frontotemporal	1
Temporo-parietal	7
Temporo-occipital	1
Fronto-temporo-parietal	6
Temporo-parieto-occipital	1

SDH was commonly found in the temporoparietal region (7 nos), followed closely by those in the front temporoparietal region (6 nos).

Table 8. Traumatic Intracranial Haemorrhage-Location and Associated Lesions

Location	Number
Frontal	5
Temporal	2
Parietal	1
Occipital	0
Associated Findings	
Oedema	4
Subarachnoid haemorrhage	1
Intraventricular haemorrhage	1
Extradural hematoma	1
Fracture	5

ICH was seen in 8% of cases and most commonly found in the frontal region. It is equally associated with subarachnoid and intraventricular haemorrhage.

Table 9. Contusions

Contusions	Number	Percentage
Non-haemorrhagic	10	20%
Haemorrhagic	40	80%
Total	**50**	**100%**

Hemorrhagic contusion was seen in 83.3% of cases and non-hemorrhagic in16.6% of cases.

Table 10. Types of Lesions and Mortality in Minor Injury

Lesion	GCS			
No.	**13**	**14**	**15**	**Total**
Fracture	5	4	13	22
Hematoma				
• SDH	1	2	2	5
• EDG	0	0	2	2
• ICH	0	0	1	1
Oedema	1	2	7	10
Contusion	0	0	15	15
SAH	1	0	0	1
Pneumocephalus	1	0	3	4
Death	1	0	0	1
Number of patients	5	4	17	26

According to the instances, there were 26 patients with a GCS score of 13 to 15. Fractures were found in 84.6 percent of patients, along with edema in 38 percent and contusions in 57 percent. Hematomas were responsible for 30% of the overall number of cases. One patient with a GCS score of 13 died, but none of the patients with higher scores perished, demonstrating that CT may predict patient outcomes.



Figure 5. Types of lesions and mortality in minor injury

The two categories concerning the percentage of CT scan displaying irregular findings and the percentage of interventions conducted by neurosurgery. Patients who did not have any health risk factors showed a slightly lower incidence of irregular CT scans were performed on all patients in this category, and none of them required surgical intervention. (confidence interval 95% = 0-0.03%). When we looked at the patients who needed neurosurgical intervention, 3 of them had depressed skull fractures on evaluation and moderate headache, and both nausea and severe headache. Three patients who had a depressed skull fracture underwent elevation of the fracture. An epidural hematoma was found in this situation, which involved the removal, where a serious headache resulted. During protracted hospitalization, the final patient was discovered to have Initial CT scans revealed numerous contusions, necessitating a partial lobectomy. For identifying patients with an abnormal CT scan, the patient had a sensitivity of 65 percent and a precision of 63 percent. if the four guidelines for the head injury had been added. Patients that needed surgical intervention would have been 100% sensitive, though.

Discussion

Patients with mild head injuries are equally common in the ER departments, but there are significantly different approaches to caring for these patients [6]. Since brain injuries, several trauma hospitals obtain head CT scans for all patients, even though the patient's state of consciousness in the ED remains uncertain [7]. Our institution faced complications as a result of this process because of the time consumption. Other patients' CT scans had to wait an extra three to four hours as a result. Even if The CT scan is a costly process, and clinical therapy for these individuals is also expensive. seldom shifts regardless of the CT scan outcome, the proposed solution is Because a

CT scan is pricey, it is highly expensive. A continuing quest for clinical variables that will help doctors to make better decisions. recognize those that will benefit from a rapid CT scan is now needed because of this issue. There are a few findings that have been found. Four therapeutic factors were suggested [8] CT scans of the head might be reduced by half. 60%. 90% sensitivity and 66% accuracy for recognizing patients who have a skewed CT scan and those who require treatment to resolve the abnormality.

The analysis results were minimal, as there were only 373 patients, and additional calculations could not be done. There are ten essential risk factors to conduct CT scans on patients. This technique was quite successful: sensitivity was 96%, and specificity was 21% [9]. Despite this, only 16% of CT scans were determined to be unnecessary, and it can be thought of as a minor allowance for using ten parameters. Cranial damage, focal developmental deficit, and the appearance of signs of basilar skull fracture on a The use of a CT scan assisted in reducing the number of CT scan performed. [10]. These specific and sensitive parameters show a sensitivity of 92% and a precision of 46% for irregular CT scans. CT scans were decreased by just 21%, but "cranial damage" was not found to be present. In the current analysis, we analyzed four objective and transparent clinical factors. Though some physicians may consider this degree of sensitivity to be too low, 65% of patients with an irregular CT scan using these four variables are measurable.

On the other hand, this degree of sensitivity is wrong. 95% CI = O-0.03% accordingly, there is a 1 in 3333 possibilities If we adopt our criterion, we'll lose someone with a surgically correctable lesion. We discovered that 90 percent of patients with an extra-axial hematoma and 75 percent of patients with a skull fracture required surgical intervention based on these variables. [11]. For clinicians who are worried about detecting and treating all patients with abnormal CT scans, regardless of their age or whether or not the intervention is needed, it is typical for them to feel dissatisfied. It is found that 41 patients out of 1302 were (3 percent) An intracranial lesion can be identified on a CT scan even if there are no risk factors. [12]. Our logic is as follows: Unexpectedly, when we examined one of these 41 patients, we found no improvement of mental status and needed no medication (2 d average).

The CT scan is not an incredibly successful device to diagnose minor brain damage, according to studies conducted by Levin et al. (lo), in which they looked at 50 patients with a moderate head injury. Twenty-five percent of the patients had irregular CT scans compared to 80 percent of the patients who had anomalies Magnetic resonance imaging (MRI) (MRI). Third, a rare type CT scan does not usually lead to a favorable prognosis for neurological healing. CT and MRI abnormalities were not sufficiently specific to Which patients are likely to have serious neurologic problems dysfunction. They determined that patients with concussions and regular CT scans have the lowest frequency of post-concussion symptoms. In contrast, patients who have had CT scans that reveal a concussion and focal contusions. had the highest rate of delayed neurocognitive dysfunction [13]. Only similar percentages of people had constant pain, dizziness, sleep disorders, and so on in both categories.

If we had expanded the research duration to two years, we might have spared almost 1300 CT scans that were needed. A cut of this size would have resulted in $1.7 million in savings for our head tracenterntre alone. About 400,000-450,000 individuals visit the doctor's clinic per year with head injuries [14]. GCS 15 or lower is the standard for those with a head injury. One study showed that 80% have a GCS of 15 or lower (14). Based on optimistic figures that say that 65% of all people with a head injury present with a GCS of 15, around 275,000 individuals who have had head injuries have a GCS of 15. If our requirements were wutilizedised worldwide, this approach could halve the number of head CT scans require red 61% (166,750). That would contribute to significant financial savings [15].

Conclusions

In conclusion, when determining whether patients with minor head injuries and a GCS of 15 require head CT scans, we found that four key clinical characteristics (nausea, vomiting, severe headache, and skull fracture) were predictive. depression) might be employed. The study showed that integrating clinical data to distinguish patients who may need neurosurgical intervention and other suspected brain disorders was very successful. These traits are intrinsic fact hats that can be found in patients in the emergency room As a result, these clinical predictors will assist reduce the number of CT scans required up to a certain point. 61%,

possibly resulting in significant benefits to the patients and the use of the CT scanner in the case of an emergency. Therefore, preliminary tests that use these standards can help to validate these results.

References

Jennings, R.M., Burtner, J.J., Pellicer, J.F., Nair, D.K., Bradford, M.C., Shaffer, M, & Tieder, J.S. (2017). Reducing head CT use for children with head injuries in a community emergency department. Pediatrics, 139(4).

Chilamkurthy, S., Ghosh, R., Tanamala, S., Biviji, M., Campeau, N.G., Venugopal, V.K., & Warier, P. (2018). Deep learning algorithms for detection of critical findings in head CT scans: a retrospective study. The Lancet, 392(10162), 2388-2396.

Nigrovic, L.E., Stack, A.M., Mannix, R.C., Lyons, T.W., Samnaliev, M., Bachur, R.G., & Proctor, M.R. (2015). Quality improvement effort to reduce cranial CTs for children with minor blunt head trauma. Pediatrics, 136(1), e227-e233.

Burstein, B., Upton, J.E., Terra, H.F., & Neuman, M.I. (2018). Use of CT for head trauma: 2007–2015. Pediatrics, 142(4).

Dayan, P.S., Ballard, D.W., Tham, E., Hoffman, J.M., Swietlik, M., Deakyne, S.J., & Traumatic Brain Injury-Knowledge Translation Study Group. (2017). Use of traumatic brain injury prediction rules with clinical decision support. Pediatrics, 139(4).

Puffenbarger, M.S., Ahmad, F.A., Argent, M., Gu, H., Samson, C., Quayle, K.S., & Saito, J.M. (2019). Reduction of computed tomography use for pediatric closed head injury evaluation at a nonpediatric community emergency department. Academic Emergency Medicine, 26(7), 784-795.

Marin, J.R., Weaver, M.D., Barnato, A.E., Yabes, J.G., Yealy, D.M., & Roberts, M.S. (2014). Variation in emergency department head computed tomography use for pediatric head trauma. Academic Emergency Medicine, 21(9), 987-995.

Akares, S.A., & Bahomil, A.A. Radiological imaging diagnosis of adult renal cystic diseases and management. Iraqi JMS. 2020; 18 (1): 12-20. doi: 10.22578. IJMS, 18(3).

Abdullah Hasan Jabbar, Salim Oudah Mezan, Salih Meri Al Absi et al., Assessment of anticholinesterase effect of polyvinylpyrrolidone/silver nanocomposite biosynthesized by Pandanus atrocarpus extract, Materials Today: Proceedings. https://doi.org/10.1016/j.matpr.2020.12.582.

S. Meri Al Absi, A. Hasan Jabbar, S. Oudah Mezan et al., An experimental test of the performance enhancement of a Savonius turbine by modifying the inner surface of a blade, Materials Today: Proceedings. https://doi.org/10.1016/j.matpr.2020.12.309

Dagash, M.T., Dagash, H.T., Jassim Obaid Al-Essawi, A., Saadallah, L., & Abdulkareem, M.A.K. (2020). Incidence of Malignant Neck Masses (MNMs) in Fallujah District, Iraq; Descriptive Cross-Sectional Study 2017-2018. Executive Editor, 11(7), 1402.

Al-Hajj, S., Hammoud, Z., Colnaric, J., Ataya, M., Macaron, M.M., Kadi, K., & TBI Working Group. (2021). Characterisation of traumatic brain injury research in the Middle East and North Africa Region: a systematic review. Neuroepidemiology, 1-12.

Mahboub, B.H., Vats, M.G., Al Zaabi, A., Iqbal, M.N., Safwat, T., Al-Hurish, F., & Al-Hajjaj, M.S. (2017). Joint statement for the diagnosis, management, and prevention of chronic obstructive pulmonary disease for Gulf Cooperation Council countries and Middle East–North Africa region, 2017. International journal of chronic obstructive pulmonary disease, 12, 2869.

Salim Oudah Mezan, Salih Meri Al Absi, Abdullah Hasan Jabbar et al., Synthesis and characterization of enhanced silica nanoparticle (SiO2) prepared from rice husk ash immobilized of 3-(chloropropyl) triethoxysilanea, Materials Today: Proceedings. https://doi.org/10.1016/j.matpr.2020.12.564

Kanne, J.P. (2020). Chest CT findings in 2019 novel coronavirus (2019-nCoV) infections from Wuhan, China: key points for the radiologist.