



EVALUATION OF KNOWLEDGE AND ATTITUDE ABOUT PERIODONTAL DISEASE AND ITS TREATMENT AMONG RURAL PUBLIC OF EASTERN UTTAR PRADESH

Dr. Janardhana Amaranath B.J., Dr. Prateek Singh, Dr. Shruti Gupta, Dr. Lynn Johnson, Dr. Neelam Das, Dr. Saurabh Tripathi, Dr. Deepika Trivedi

Rama Dental College Hospital & Research Centre, Rama University, Mandhana, Kanpur, UP;

drjanardhana.rdc@ramauniversity.ac.in

Abstract:

Introduction: Majority of the Indians lives in rural India. Periodontal diseases are highly prevalent in India and periodontitis is one of the main causes of tooth loss. Periodontal diseases are related to various systemic diseases. Accessibility to dental care is one of the major challenges of the rural population and hence, dental health education is an essential pre requisite for improving the periodontal health in rural population.

Objective: To evaluate the knowledge related to periodontal disease, its impact on systemic health and to assess the attitude towards periodontal treatment among rural population of eastern Uttar Pradesh.

Material and Methods: Questionnaire survey on knowledge and attitude towards periodontal disease and its treatment was conducted among 1000 participants age ranging between 35-65 years, The responses recorded were subjected to statistical analysis to obtain results.

Results: As per the findings, most of the respondents had poor knowledge about periodontal disease and its treatment. In terms of knowledge related to periodontal disease, its impact on systemic health and their attitude towards periodontal treatment was not statistically significant.

Conclusion: It was concluded that rural public had poor knowledge about periodontal disease, its impact on systemic health was also very poor and they also had a very poor attitude towards periodontal treatment. Hence, special attention is needed to educate and motivate the rural population regarding importance of maintaining periodontal health.

Keywords: Attitude; Knowledge; Periodontal disease; Periodontal treatment; Rural public; Systemic health.

DOI Number: 10.48047/nq.2020.18.10.NQ20245

NeuroQuantology 2020; 18(10):179-185

INTRODUCTION

The main reason for tooth loss in India is periodontal disease. 90% of people have periodontal and gingival disorders. According to epidemiological data from the Global Oral Health Data Bank, only a small fraction (3%) of

individuals between the ages of 35 and 44 and seniors over the age of 60 in underdeveloped nations like India have healthy periodontium. The mouth is thought of as the body's entrance and serves as a mirror to reflect the general state of health.¹ Nowadays, it is



acknowledged that oral health and general health go hand in hand.² Some chronic systemic disorders, including diabetes, congestive heart failure, chronic kidney disease, obesity, stroke, and even some coagulative diseases, may have a close connection to poor periodontal health.³ Periodontal disease prevalence varies widely across the world depending on factors including age, sex, race, ethnicity, education, geographic location, environmental conditions, oral hygiene practises, way of life, social traits, and dental awareness. India's rural population is 64.61 percent. Poverty and disease, both of which can have a serious influence on wellness, are frequently linked to rural locations.⁴ The average level of education, the kind of jobs people have, their wages, and other related characteristics of wellbeing varies between urban and rural areas. There are still significant gaps in health and access to healthcare for rural populations.

AIM

To evaluate the knowledge related to periodontal disease, its impact on systemic health and to assess the attitude towards periodontal treatment among rural population of eastern Uttar Pradesh.

MATERIAL AND METHODS

This questionnaire survey was conducted among rural public of Eastern Uttar Pradesh. Out of 1055 participants, 55 subjects could not complete the questionnaire proforma

PERCENTAGE

- 0-20
- 21-40
- 41-60
- 61-80
- 81-100

STATISTICAL ANALYSIS

Statistical analysis was carried out using Statistical Package of Social Sciences (SPSS Inc. Chicago IL, version 11.0 for windows.) For quantitative variations percentage were calculated. Means were compared among the three groups. Mean, standard deviation and p value were calculated using independent t test.

were excluded, all the other responses were considered. Ethical clearance was obtained by institutional ethical committee. Written informed consent was obtained from each study participants. Both male and female subjects within the age group of 35-65 years, resident of rural population were included in the study.

The purpose of the study was explained to every participant and they were asked to fill the questionnaire. The questionnaire proforma consisting of total 15 questions in native language (Hindi) were given to them to record their responses with respect to each question by ticking the answer "yes" or "no". And the questionnaire consists of three sections, first section consisted of 5 questions to assess the knowledge about periodontal diseases. Second section consists of 5 questions to assess the knowledge on impact of periodontal diseases on general health. The third section consists of 5 questions to assess the attitude towards periodontal treatment. Participation was voluntary, and all the participants were assured about the confidentiality of the data received. The percentage analysis of knowledge of periodontal disease its impact on systemic health and to evaluate the attitude towards periodontal treatment, among rural public of Eastern Uttar Pradesh are graded based on the following grading system:

GRADING

- Very poor
- Poor
- Average
- Fair
- Good

RESULTS

The data from the questionnaire comprised of 73.8% male and 26.2% female. Most of the participants of the study were in the age group of 45-50 years with mean and standard deviation of 44.85 ± 8.60 years. [table 1]



Table 1: Demographic representation of age and gender of rural public of Eastern Uttar Pradesh

Variable	Category	Mean ± SD / n(%)
Age (in years)		44.85 ± 8.60
Gender	Male	738 (73.8)
	Female	262 (26.2)


Table 2 and Graph 1 shows that majority of the subjects (78.3%) had knowledge about periodontal disease but they were unaware about dental plaques as the main cause for periodontal disease (80.6%). 84% of the subjects did not agree that it is necessary to have good periodontal care for healthy and firm dentition. Only 4.2% and 9.5% of the subjects knew that periodontal disease if untreated can ultimately results in loss of teeth, and periodontal disease can manifest with many symptoms such as bleeding gums, tooth sensitivity, tooth mobility, swollen gums, receding gums. etc.

Table 2: Evaluation of knowledge of periodontal disease among rural public of Eastern Uttar Pradeshn(%)

Question	Yes	No
1. Do you know what is periodontal / gum disease (pyorrhoea)?	783 (78.3)	217 (21.7)
2. Are you aware that dental plaque/poor oral hygiene is the main cause for periodontal/gum disease?	194 (19.4)	806 (80.6)
3. Do you agree that it is necessary to have good periodontal/gum care for healthy and firm dentition?	160 (16.0)	840 (84.0)
4. Do you know that periodontal disease if untreated can ultimately results in loss of teeth?	42 (4.2)	958 (95.8)
5. Do you know that gum disease can manifest with many symptoms such as bleeding gums, tooth sensitivity, tooth mobility, swollen gums, receding gums etc.?	95 (9.5)	905 (90.5)

181

Graph 1: Evaluation of knowledge of periodontal disease among rural public of Eastern Uttar Pradesh.



None of the subjects were aware about the fact that oral health is important to maintain good systemic health and diabetes mellitus can lead to worsening of periodontal health and periodontal diseases is one of the complications of diabetes mellitus. Only 3.1% of the subjects were aware




about the fact that good oral hygiene maintenance and periodontal care can lead to better blood sugar control. Very few subjects knew that periodontal disease can be a risk factor for coronary heart disease (13.8%) and periodontal disease can be a risk factor for adverse pregnancy outcome (preterm low birth weight) (12.4%) as shown in table 3, graph 2

Table 3: Evaluation of Knowledge on impact of periodontal disease on general health among rural public of Eastern Uttar Pradesh

Question	Yes	No
6. Do you know that the oral health is important to maintain good general/systemic health?	0	1000 (100)
7. Do you aware that diabetes mellitus can lead to worsening of periodontal/gum health and periodontal diseases is one of the complications of diabetes mellitus?	0	1000 (100)
8. Are you aware that good oral hygiene maintenance and periodontal health/gum care can lead to better blood sugar control?	31 (3.1)	969 (96.9)
9. Do you know that periodontal/gum disease can be a risk factor for coronary heart disease?	138 (13.8)	862 (86.2)
10. Do you know that periodontal/gum disease can be a risk factor for adverse pregnancy outcome (preterm low birth weight)?	124 (12.4)	876 (87.6)

182

Graph 2: Evaluation of Knowledge on impact of periodontal disease on general health among rural public of Eastern Uttar Pradesh




Only 17.8% of the subjects agreed that daily brushing of the teeth is necessary for good periodontal health. Only 26.2% of the subjects agreed that cleaning of teeth by dentist is must for everyone. Only 2.4% of the subjects agreed that consulting periodontist is necessary for all. Only 13.8% of the subjects agreed that regular gum care is necessary to have healthy and firm dentition. Only 12.9% of the subjects agreed that continuous monitoring of periodontal health is mandatory for better systemic health, as shown in table 4, graph 3.

Table 4: Evaluation of attitude towards periodontal treatment among rural public of Eastern Uttar Pradesh

Question	Yes	No
11. Do you think daily brushing of teeth is necessary for good	178 (17.8)	822 (82.2)

periodontal/gum health?		
12. Do you think cleaning of teeth by dentist is must for everyone?	262 (26.2)	738 (73.8)
13. Do you think consulting periodontist/gum care specialist is necessary for all?	24 (2.4)	976 (97.6)
14. Do you think regular gum care is necessary to have healthy and firm dentition?	138 (13.8)	862 (86.2)
15. Do you think continuous monitoring of gum health is mandatory for better general health?	129 (12.9)	871 (87.1)

Graph 3: Evaluation of attitude towards periodontal treatment among rural public of Eastern Uttar Pradesh



There was no difference in overall knowledge of periodontal disease, knowledge on impact of periodontal disease on systemic health and attitude towards periodontal health among male and female subjects as shown in table 5.

Table 5: Comparison of knowledge, impact, and attitude towards periodontal treatment (in %) among male and female subjects

Domain	VAR00001	Mean	Std. Deviation	P value
Knowledge	Male	25.420	14.804	0.828
	Female	25.649	14.282	
Impact	Male	5.935	13.504	0.766
	Female	5.649	13.049	
Attitude	Male	15.068	14.938	0.107
	Female	13.359	14.121	

Independent t test; * indicates significant difference at $p \leq 0.05$

DISCUSSION

Periodontal diseases are one of the common and major dental diseases amongst the human population around the globe. There have been very few studies in India regarding periodontal disease in rural area. In Uttar

Pradesh, 77.7% of the population belongs to the rural category.

This study was the first of its kind, a questionnaire study was conducted with 5 questions in each group to explore the knowledge related to periodontal disease, its impact on systemic health and to evaluate the



attitude towards periodontal treatment respectively among rural public of Eastern Uttar Pradesh. The questionnaire was in native language (hindi) for the ease of understanding of the subjects.

In this study, 78.3% rural population had knowledge about periodontal disease but they were unaware about dental plaque and poor oral hygiene as the main cause for periodontal disease which is in accordance with the study by Bala K et al.⁵ in 2019 they have found that there was adequate knowledge but poor attitude regarding oral health. In our study 84% of the subjects did not agree that it is necessary to have good gum care for healthy and firm dentition. Only 4.2% and 9.5% of the subjects knew that periodontal disease if untreated can ultimately results in loss of teeth and gum disease. This is in accordance with studies by Sekhon *et al.*⁶, Ramoji Rao *et al.*⁷ in 2015 and 2017 respectively where they found that there was a higher prevalence of periodontal disease in rural subjects due to lack of awareness.

In the present study only 3.1% of the subjects were aware about the fact that good oral hygiene maintenance and gum care can lead to better blood sugar control. Very few subjects i.e., 13.8% knew that gum disease can be a risk factor for coronary heart disease and 12.4% knew periodontal disease can be a risk factor for adverse pregnancy outcome (preterm low birth weight), which is in accordance with the studies conducted by Gandhimadhi D. et al.⁸, Dhadse P. et al.⁹ both in the year 2010 and Jepsen S. *et al.*¹⁰ 2020.

Only 17.8% of rural population agreed that daily brushing of teeth is necessary for good periodontal health, along with 26.2% of the subjects agreed that cleaning of teeth by dentist is must for everyone and 13.8% of the subjects agreed that regular gum care is necessary to have healthy gum which is in accordance with the study conducted by Jain N. et al.¹¹ in 2012 stated that there was an acute lack of oral hygiene awareness and limited knowledge of oral hygiene practices among patients of Jodhpur.

Overall, it was found that rural population had poor knowledge about periodontal disease, its impact on systemic health was also very poor

and they also had a very poor attitude towards periodontal treatment.

About 65% of population lives in rural areas of India, hence it is of utmost requirement to educate, motivate rural population about oral and periodontal health and to provide accessibility for dental and periodontal treatment to achieve comprehensive health among both among urban and rural population.

CONCLUSION

Rural public had a poor knowledge about periodontal disease, the impact of periodontal disease on systemic health was also very poor and there was a very poor attitude towards periodontal treatment. Hence, special attention is needed to educate and motivate the rural population regarding importance of maintaining periodontal health for firm and healthy dentition, through comprehensive dental health educational programs by conducting various dental camps and public oral and periodontal health awareness programs.

Funding: No funding sources

Source of Support: Nil

Conflict of Interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee. RDCHRC/ETHICSCOMMITTEE/0156

REFERENCES

1. Kassak KM, Dagher R, Doughan B. Oral hygiene and lifestyle correlates among new undergraduate university students in Lebanon. *J Am Coll Health* 2001; 50:15-20.
2. Igari K, Kudo T, Toyofuku T, Inoue Y, Iwai T. Association between periodontitis and the development of systemic diseases. *Oral Biol Dent*. 2014;2(1):4.
3. Gil-Montoya JA, de Mello AL, Barrios R, Gonzalez-Moles MA, Bravo M. Oral health in the elderly patient and its impact on general well-being: A nonsystematic review. *Clin Interv Aging* 2015; 10:461-7.
4. Jain N, Mitra D, Ashok KP, Dundappa J, Soni S, Ahmed S. Oral hygiene-awareness and practice among patients attending OPD at

Vyas Dental College and Hospital, Jodhpur. *J Indian Soc Periodontol.* 2012;16(4):524-528.

5. Bala K, Gupta R, Ara A, Sahni B. A KAP study of oral health status among adults in a rural area of Jammu District. *Int J Community Med Public Health* 2019;6:135-41..

6. Sekhon TS, Grewal S, Gambhir RS. Periodontal health status and treatment needs of the rural population of India: A cross-sectional study. *J Nat Sc Biol Med* 2015;6:111-5.

7. Ramoji Rao MV, Katari PK, Vegi L, Bypureddy TT, Prabhakara Rao KS, Tejaswi KS. Prevalence of periodontal diseases among rural population of Mustabad, Krishna District. *J Int Soc Prevent Communit Dent* 2016;6:S59-63.

8. Gandhimadhi D, Mythili R. Periodontal infection as a risk factor for preterm low birth weight. *J Indian Soc Periodontol.* 2010;14(2):114-120.

9. Dhadse P, Gattani D, Mishra R. The link between periodontal disease and cardiovascular disease: How far we have come in last two decades ?. *J Indian Soc Periodontol.* 2010;14(3):148-154.

10. Jepsen S., Suvan J, Deschner J, The association of periodontal disease with

metabolic syndrome and obesity. *Periodontology* 2000. 2020;83:125-153.

11. Jain N, Mitra D, Ashok KP, Dundappa J, Soni S, Ahmed S. Oral hygiene-awareness and practice among patients attending OPD at Vyas Dental College and Hospital, Jodhpur. *J Indian Soc Periodontol* 2012;16:524-8.

Legends

Table 1: Demographic representation of age and gender of rural public of Eastern Uttar Pradesh

Table 2: Evaluation of knowledge of periodontal disease among rural public of Eastern Uttar Pradesh n (%).

Table 3: Evaluation of Knowledge on impact of periodontal disease on general health among rural public of Eastern Uttar Pradesh.

Table 4: Evaluation of attitude towards periodontal treatment among rural public of Eastern Uttar Pradesh.

Table 5: Comparison of knowledge, impact, and attitude towards periodontal treatment (in %) among male and female subjects.