



# Formulation Development and Characterization of Ethosomes containing Ethanolic Flowers extract of *Sarcosteema acidum* (Roxb.) Voigt

Priyanka Vijay Patil\*, Rajendra Singh Bapna, Rakesh Patel and Karunakar Shukla  
Faculty of Pharmacy, Dr. A.P.J. Abdul Kalam University, Indore (M.P.) – India  
\*Corresponding Author

## Abstract

Formulation of herbal drugs are becoming the promising tool for the cure of many diseases with the less toxic effects, better therapeutic effect, have good patient compliance and to minimize a problems of the stability, toxicity, poor bioavailability various novel drug delivery system are presently preferred. Development of NDDS such as phytosomes, ethosomes, herbal transdermal patches, nanoparticles is of great interest using herbal extract. The present work aims to developed ethosomes of ethanolic extract of *Sarcosteema acidum* (Roxb.) Voigt Flowers. The developed ethosomes were evaluated. Results indicates that EESAF formulation code SAF-5 have maximum EE i.e.,  $93.34 \pm 0.44$ .

**Key-words:** Ethosomes, Leaves Extract, *Sarcosteema acidum* (Roxb.) Voigt

**DOI Number:** 10.48047/nq.2022.20.22.NQ10488

**NeuroQuantology2022;20(22): 4814-4817**

4814

## Introduction

Ethosomes are developed by mixture of phospholipids and ethanol in which the ethanol concentration is maintained at higher level. This carrier will penetrate into the skin deeply to improve the delivery of drugs in to deeper layers of the skin. Ethosomes is non invasive vesicles and are commonly formulated as a carrier for the delivery of the herbal drugs into deeper layers of the skin. [1] *Sarcosteema acidum* (Roxb.) Voigt, is annual herb having height of 3-5 feet commonly known as somlata. The species is of flowering plant in the family Apocynaceae, typically found in the arid parts of peninsular India where it is used in religious sacrifices. The plant is religiously linked

to Hinduism and is believed to be a major ingredient of the Soma in Ancient India. This plant is used in treatment of respiratory ailments like mild bronchospasms, allergic rhinitis, sinusitis, coryza or common cold. This plant is an ancient discovery of siddhars and they used it in extraction of somarasam which is considered to be an amrutam of life, and used in somayagam. [2] The present study was designed to formulate and evaluated ethosomes of ethanolic flowers extract of *Sarcosteema acidum* (Roxb.) Voigt.

## Material and Methods

### Extract



Ethanolic leaves extract of *Sarcosteema acidum* (Roxb.) Voigt (EESAF) was taken after extraction. The extract was prepared using soxhlet apparatus. [3]

**Development of Ethosomes containing extract**  
 Ethosomal formulations were prepared by using the cold method. The ethanolic vascular system was composed of phospholipid (2.0% to 4%W/V), ethanol (20% to 40% V/V), propylene glycol (20 % V/V), extract (EESAF, 0.5% W/V) and distilled water to 100% (V/V). Phospholipid was dissolved along with the drug in ethanol. This mixture was heated to 400 C ± 10 C and a fine stream of distilled water was added slowly,

with constant mixing at 700 rpm with a mechanical stirrer in a closed container. Mixing was continued for an additional 5 minutes, while maintaining the system at 400 C ± 10 C. The preparation was left to cool at room temperature for 30 min and then it was sonicated at 40 C for five cycles of 3 minutes each with a minute rest between cycles using a probe sonicator. Nine formulations were prepared using different concentration of phospholipid and ethanol among them optimized formulation was selected for characterization and evaluation studies. [4-5]

**Table 1: Compositions of ethosomes of *Sarcosteema acidum* (Roxb.) Voigt**

Formulation Code	Conc. of Phospholipid (w/v)	Conc. of Ethanol (v/v)	Conc. of Propylene Glycol (v/v)	Conc. of Extract (w/v)	Conc. of distill water (v/v)
SAF-1	2%	20%	20%	0.5%	Up to 100%
SAF-2	3%	30%	20%	0.5%	Up to 100%
SAF-3	4%	40%	20%	0.5%	Up to 100%
SAF-4	2%	30%	20%	0.5%	Up to 100%
SAF-5	3%	40%	20%	0.5%	Up to 100%
SAF-6	4%	20%	20%	0.5%	Up to 100%
SAF-7	2%	40%	20%	0.5%	Up to 100%
SAF-8	3%	20%	20%	0.5%	Up to 100%
SAF-9	4%	30%	20%	0.5%	Up to 100%

4815

**Evaluation of Ethosomes [4-5]**

**Image analysis of ethosomes by optical microscope**

Visualization done by image analysis optical microscope (Labomed Microscope, Leica ATC2000, India). The optical microscope is attached with the software Digipro V 4.0, through which image analysis was done, photographs were captured

**Particle size analysis and Zeta potential**

ZP, vesicle size and PDI were measured by zetasizer (Malvern Instruments, Malvern)

**Entrapment efficiency**

The total volume of the ethosomal suspension was measured. 5ml of this formulation was diluted with distilled water up to 8 ml and centrifuged at 15,000 rpm for 45 min at 40C using a cooling centrifuge. After centrifugation, the supernatant and sediment were recovered, their volume was measured. Then sediment was lysed using n- propanol and filtered through a 0.45 µm nylon disk filter. The concentration of extract in the supernatant and sediment was analyzed by UV- spectroscopic method at 260 nm. The percent drug entrapment was calculated using the following equation:

$$\% \text{ Entrapment efficiency} = \frac{\text{Amount of entrapped drug recovered}}{\text{Total amount of drug}} \times 100$$



### Stability testing

The ability of vesicles to retain the drug (i.e., drug-retentive behavior) was assessed by keeping the ethosomal suspensions at different temperatures, i.e.,  $4 \pm 2$ ,  $25 \pm 2$  (room temperature, RT),  $37 \pm 2$  and  $45 \pm 2$  °C for different periods of time (1, 20, 40, 60, 80 days). The vesicular suspensions were kept in sealed vials (10 ml capacity) and their image analysis done by optical microscopy.

### Results and Discussion

Nine different batches of ethosomes were prepared by using the cold method. In the investigation extract (EEPFL, 0.5% W/V) is used and along with phospholipid (2.0% to 4%W/V), ethanol (20% to 40% V/V), propylene glycol (20 % V/V), and distilled water to 100% (V/V). Nine formulation of i.e., SAF-1 to SAF-9 were prepared and further evaluated for various parameters. All the images depict smooth surface. It has been reported that if the vesicular size is <300 nm, they can deliver their contents into deeper layers of skin to some extent. For all developed formulation of *Sarcosteema acidum* (Roxb.) Voigt the vesicle

size was found be in range. Size analysis of the formulation depending upon the concentration of SPC and ethanol was found to range from 208.14 to 259.11 nm. ZP is an important parameter that affects stability. All the ethosomal formulation was found to have negative ZP due to the net charge of the lipid composition in the formulation. The negative ZP is responsible for enhanced percutaneous permeation of drug. The ranges of EE of ethosomes were between 34.21% and 94.23%. The ethanol concentration in the ethosome system should not be too high, and generally, should be kept below 40%. As increasing concentration of ethanol results in leaking of the drug from the lipid bilayer due to which EE decreases therefore ethanol concentration only up to 40% was considered. The ability of vesicles to retain the drug (i.e., drug-retentive behavior) was assessed by keeping the ethosomal suspensions at different temperatures, i.e.,  $4 \pm 2$  °C,  $25 \pm 2$  °C (room temperature, RT),  $37 \pm 2$  and  $45 \pm 2$  °C for different periods of time (1, 20, 40, 60, 80 days). The results of stability study were presented in Table 3.

4816

**Table 2: Evaluation of Ethosomes of *Sarcosteema acidum* (Roxb.) Voigt**

Formulation Code	Vesicle Size (nm)	Zeta Potential (mV)	Entrapment Efficiency (%)
SAF-1	208.14 ±0.01	-39.1	48.29±0.04
SAF-2	212.17±1.21	-35.3	59.22±0.11
SAF-3	231.23±0.08	-34.4	69.20±0.18
SAF-4	223.20±0.21	-35.6	87.29±0.21
SAF-5	259.11±0.01	-30.1	94.23±0.40
SAF-6	232.39 ±0.02	-30.0	88.21±0.22
SAF-7	214.35±0.02	-34.3	83.14±0.02
SAF-8	241.01±1.39	-40.5	82.11±0.12
SAF-9	229.11±0.22	-41.2	78.23±0.10

**Note:** All values are Mean±SEM; n=3



**Table 3: Stability studies of ethosomes containing *Sarcosteema acidum* (Roxb.) Voigt flower extract**

S/No.	Ethosomes	Stability condition	20 days	40 days	60 days	80days
1.	SAF-5	4±2 °C	Stable	Stable	Stable	Stable
2.		25±2 °C	Stable	Stable	Stable	Stable
3.		37±2 °C	Fusion of vesicles	Fusion of vesicles	Complete deformation of vesicles	Complete deformation of vesicles
4.		45±2 °C	Fusion of vesicles	Fusion of vesicles	Complete deformation of vesicles	Complete deformation of vesicles

### Conclusion

From the results obtained it was concluded that the formulation code SAF-5 have maximum EE i.e., 94.23±0.40 having vesicle size 259.11±0.01 nm with zeta potential of -30.1 mv. Also results of stability studies indicate that temperature of 4±2 °C and 25±2 °C have stable ethosomes while in other two temperature there is deformalities. The ethanol concentration in the ethosome system should not be too high, and generally, should be kept 40%. As increasing concentration of ethanol results in leaking of the drug from the lipid bilayer due to which EE decreases therefore ethanol concentration only up to 40% was considered as best for formulation of ethosomes.

### References

1. Touitou E, Dayan N, Bergelson L, Godin B, Eliaz M. Ethosomes - Novel vesicular carriers for enhanced delivery: Characterization and skin

penetration properties. *J Control Release* 2000;65:403-18.

2. Singh, N. P. (1988). *Flora of Eastern Karnataka, Volume 1*. Mittal Publications. p. 416. ISBN 9788170990673
3. Harborne JB. *Phytochemical methods*, Chapman and Hall, 1984, 1<sup>st</sup> Edition, London.
4. Li G, Fan Y, Fan C, Li X, Wang X, Li M, et al. Tacrolimus-loaded ethosomes: Physicochemical characterization and *in vivo* evaluation. *Eur J Pharm* 2012;82:49-57.
5. Seema A. Recent development of herbal formulation a novel drug delivery system. *International Ayurvedic Medical Journal*. 2014; 2 (6):953-958.

