



Thermodynamic probability of thermal decomposition of ammonia

¹Olimjon Panjiev, ²Arziqul Panjiev, ³Erkin Yoqubov

¹Karshi Engineering and Economic Institute, Ph.D. tech. Sciences, Associate Professor

²Karshi Engineering and Economic Institute, Ph.D. tech. Sciences, Associate Professor

³Karshi State University, Ph.D. chem. Sciences, Associate Professor

E-mail olimjon573@mail.ru

2076

Annotation

Technical solutions have been developed to create a technology for obtaining calcium cyanamide from lime, carbonate anhydride and industrial ammonia. A technological scheme for obtaining nitrogen fertilizer and an effective defoliant, calcium cyanamide, has been developed and optimal technological parameters of the process have been determined. An experimental batch of calcium cyanamide has been produced at an experimental installation.

Keywords. Carbon dioxide, ammonia, expansion gas, calcium cyanamide, waste gases.

DOI Number: 10.14704/NQ.2022.20.12.NQ77181


NeuroQuantology2022;20(12): 2076-2083

Аннотация

Разработаны технические решения по созданию технологии получения цианмида кальция из извести, ангидрида карбоната и промышленного аммиака. Разработана технологическая схема получения азотного удобрения и эффективного дефолианта – цианмида кальция и определены оптимальные технологические параметры процесса на экспериментальной установке произведена опытная партия цианмида кальция.

Ключевые слова. Диоксид углерода, аммиак, экспансерный газ, цианамид кальция, отходящие газы.

Introduction. In the synthesis of calcium cyanamide, it is possible to propose the following chemical reaction, which can carry out additional consumption of expensive ammonia:



Literature review. In order to carry out thermodynamic calculations of the probability of the occurrence of this reaction, the following equations were compiled:

The thermodynamic calculations of the thermal decomposition of ammonia were based on the following, derived by us mathematically by multiplying by the reduced coefficient 4.184, the results of thermodynamic calculations were converted from the metric system of measures to the international SI system.

1. In accordance with the Hess law, the change in the enthalpy of the reaction is determined under standard conditions:

$$\Delta H_{298}^{\circ} = \sum \Delta H_{\text{пр}}^{\circ} - \sum \Delta H_{\text{и.в.}}^{\circ} = [(0,5\Delta H_{\text{N}_2}^{\circ} + 1,5\Delta H_{\text{H}_2}^{\circ}) - (\Delta H_{\text{(NH}_3)}^{\circ})] \cdot 10^3 \text{ cal/mol}$$

2. From the following expression, the thermal effect of the reaction is determined at 298 K:

$$Q_{298} = -\Delta H_{298}^{\circ} \text{ cal/mol}$$

3. For standard conditions, we calculate the absolute value of the entropy using the equation:

$$\Delta S_{298}^{\circ} = \sum \Delta S_{\text{пр}}^{\circ} - \sum \Delta S_{\text{и.в.}}^{\circ} = (0,5\Delta S_{\text{N}_2}^{\circ} + 1,5\Delta S_{\text{H}_2}^{\circ}) - \Delta S_{\text{NH}_3}^{\circ}, \text{ cal/mol.deg}$$

4. At a temperature of 298 K, the change in the Gibbs energy is:

$$\Delta G_{298}^{\circ} = \Delta H_{298}^{\circ} - T^* \Delta S_{298}^{\circ}, \text{ cal/mol}$$

5. According to the Kirchoff equation, the dependence of the true molar isobaric heat capacity of the reagents on temperature is determined:

$$\Delta C_p^{\circ} = \Delta a + \Delta b \cdot T + c^1/T^2,$$



where: Δa , Δb , Δc^1 - the constant coefficients of the equation are defined as the algebraic difference between the corresponding parameters of the reaction products and starting materials, taking into account their stoichiometric coefficients;

where: T- temperature in degrees Kelvin.

$$5.1 \Delta a = \sum \Delta a_{np} - \sum \Delta a_{i.b.} = (0,5\Delta a_{N_2} + 1,5\Delta a_{H_2}) - \Delta a_{NH_3}$$

$$5.2 \Delta b = \sum \Delta b_{np} - \sum \Delta b_{i.b.} = (0,5\Delta b_{N_2} + 1,5\Delta b_{H_2}) - \Delta b_{NH_3}$$

$$5.3 \Delta c^1 = \sum \Delta c^1_{np} - \sum \Delta c^1_{i.b.} = (0,5\Delta c^1_{N_2} + 1,5\Delta c^1_{H_2}) - \Delta c^1_{NH_3}$$

6. The integration constant of the equation for the dependence of the change in the enthalpy of the reaction on temperature is calculated by the formula:

$$\Delta H^{\circ}_0 = \Delta H^{\circ}_{298} - \Delta a * 298 + \Delta b * 298^2 / 2 + \Delta c^1 / 298$$

7. For different temperatures, the change in the enthalpy of the components is calculated by the following formula

$$\Delta H^{\circ}_T = \Delta H^{\circ}_0 + \int_{298}^T \Delta C_p dT = \Delta H^{\circ}_0 + \int_{298}^T (\Delta a + \Delta b \cdot T + \Delta c^1 / T^2) dT$$

After integrating this expression, we have the following formula:

$$\Delta H^{\circ}_T = \Delta H^{\circ}_0 + \Delta a * T + \Delta b * T^2 / 2 - \Delta c^1 / T$$

8. The thermal effect of a chemical reaction at different temperatures is determined as follows:

$$Q_{pT} = -\Delta H^{\circ}_T, \text{ cal/mol}$$

9. According to the van't Hoff isobar equation, the dependence of the equilibrium constant of a chemical reaction on temperature is determined:

$$d \ln K_{pT} / dT = \Delta H^{\circ}_T / R * T^2$$

where:

$$d \ln K_{pT} = (\Delta H^{\circ}_T / R * T^2) dT$$

or:

$$d \ln K_{pT} = \Delta H^{\circ}_T / (R * T)$$

10. The decimal logarithm of the equilibrium constant of a chemical reaction is calculated by the formula at a temperature of 298K:

$$\lg K_{p298} = -\Delta G^{\circ}_{298} / R * 298$$

11. The integration constant of the decimal logarithm dependency equation is found from the equation:

$$K_0 = \lg K_{p298} + \Delta H^{\circ}_0 / (4,576 * \lg 298) - (\Delta a * \lg 298) / 1,987 - (\Delta b * \lg 298) / 9,150 - \Delta c^1 / (9,150 * \lg 298^2) - (\Delta c^1 * \lg 298^2) / 27,45$$

12. The value of the decimal logarithm of the equilibrium constant of a chemical reaction is determined by the formula:

$$\lg K_{pT} = -\Delta H^{\circ}_0 / (4,576 * T) + (\Delta a * \lg T) / 1,987 + (\Delta b * \lg T) / 9,150 * \Delta c^1 / (9,150 * T^2) + (\Delta c^1 * 298^2) / 27,45 + K_0$$

13. The value of the change in the Gibbs energy of a chemical reaction depending on the temperature of the solution:

$$\Delta G^{\circ}_T = -R * T * \lg K_{pT} = -4,576 * T * \lg K_{pT}, \text{ кал/моль}$$

The C++Builder 6 computer program for calculating the thermodynamic decomposition probability was as follows:

```
//-----
#include <vcl.h>
#include <math.h>
#pragma hdrstop
#include "Unit1.h"
```



```
//-----  
#pragma package(smart_init)  
#pragma resource "*.dfm"  
TForm1 *Form1;  
//-----  
__fastcall TForm1::TForm1(TComponent* Owner)  
: TForm(Owner)  
{  
}  
//-----  
void __fastcall TForm1::Button1Click(TObject *Sender)  
{  
double n1=1, n2=0, n3=0, n4=0, H1=-10980, H2=0,  
H3=0, H4=0, n5=0.5,n6=1.5, n7=0, n8=0, H5=0, H6=0,  
H7=0, H8=0, S1=46.047, S2=0,S3=0, S4=0, S5=45.77,  
S6=31.195, S7=0, S8=0, A1=7.122, A2=0,A3=0, A4=0,  
A5=6.663, A6=0, A7=0, A8=0, B1=0.00609, B2=0, B3=0,  
B4=0, B5=0.001021,B6=0.000779, B7=0, B8=0,  
C11=-39900, C12=0, C13=0,C14=0, C15=0, C16=11900,  
C17=0, C18=0, T=298, T1=373, K1=4.576,K2=1.987,  
K3=9.150, K4=27.45,DHO298, DS298, GO298, DA, DB,  
DC1, DCPT1,HO, DHTO1, LOGKP298, KO, LOGKPT1, DGOT1;  
  
DHO298 = (n5 * H5 + n6 * H6 + n7 * H7 + n8 * H8) - (n1 * H1 + n2 * H2 + n3 * H3 + n4 * H4);  
DS298 = (n5 * S5 + n6 * S6 + n7 * S7 + n8 * S8) - (n1 * S1 + n2 * S2 + n3 * S3 + n4 * S4);  
GO298 = DHO298 * (T * DS298);  
DA = (n5 * A5 + n6 * A6 + n7 * A7 + n8 * A8) - (n1 * A1 + n2 * A2 + n3 * A3 + n4 * A4);  
DB = (n5 * B5 + n6 * B6 + n7 * B7 + n8 * B8) - (n1 * B1 + n2 * B2 + n3 * B3 + n4 * B4);  
DC1 = (n5 * C15 + n6 * C16 + n7 * C17 + n8 * C18) - (n1 * C11 + n2 * C12 + n3 * C13 + n4 * C14);  
DCPT1 =float (DA + DB * T1 +DC1) / float(T1 * T1);  
HO = float (DHO298 - DA * T -(DB * (T * T))) / float (2- DC1 / T);  
DHTO1 = HO + DA * T1 +(DB * (T * T)) - float (2- DC1 / T1);  
LOGKP298 = log10(fabs(- GO298 / float(K1 * T)));  
KO = LOGKP298 + float (HO) / float((K1 * T) - (DA * (log10(T))) / float( K2 - (DB * T1)) / float (K3 + DC1) /  
float(K3 * (T1 * T1)));  
LOGKPT1 = log10(fabs(- HO / (K1 * T1 ) + (DA * log10(T1)) / K2 + (DB * T1) / (K3 * (T * T)) + KO));  
DGOT1 = - K1 * T1 * LOGKPT1;  
StringGrid1->Cells[0][0]=FloatToStrF(DHO298,ffFixed,15,2);  
StringGrid1->Cells[1][0]=FloatToStrF(DS298,ffFixed,15,2);  
StringGrid1->Cells[2][0]=FloatToStrF(GO298,ffFixed,15,2);  
StringGrid1->Cells[3][0]=FloatToStrF(DA,ffFixed,15,2);  
StringGrid1->Cells[4][0]=FloatToStrF(DB,ffFixed,15,2);  
StringGrid1->Cells[5][0]=FloatToStrF(DC1,ffFixed,15,2);  
StringGrid1->Cells[6][0]=FloatToStrF(DCPT1,ffFixed,15,2);  
StringGrid1->Cells[7][0]=FloatToStrF(HO,ffFixed,15,2);  
StringGrid1->Cells[8][0]=FloatToStrF(DHTO1,ffFixed,15,2);
```



```
StringGrid1->Cells[9][0]=FloatToStrF(LOGKP298,ffFixed,15,2);
StringGrid1->Cells[10][0]=FloatToStrF(KO,ffFixed,15,2);
StringGrid1->Cells[11][0]=FloatToStrF(LOGKPT1,ffFixed,15,2);
StringGrid1->Cells[12][0]=FloatToStrF(DGOT1,ffFixed,15,2);
}
```

//-----

According to the above program, a number of calculations were made, changing 122 temperature values from 373 to 1473K in increments of 100 degrees.

The results obtained are presented in Tables 1-4.:

Table 1

Values of thermal effect, entropy and change of Gibbs energy for the reaction of ammonia decomposition under standard conditions

Qp 298		DS 298		GO 298	
cal/mol	J/mol	cal/mol	J/mol	cal/mol	J/mol
-10980	-4594,32	23,635	98,889	+3936,77	+16471,445

Table 2

Values of thermal effect of ammonia decomposition reaction at different temperatures.

QPT1	T1, K	373	473	573	673	773
	кал/моль	-116674,204	-12046,9	-12393,931	12703,199	12972
	Дж/моль	-48815,581	-50404,229	-51856,207	53150,184	542,75

Table 3 continued

Values of thermal effect of ammonia decomposition reaction at different temperatures.

QPT1	T1, K	873	973	1073	1173	1273	1373	1473
	cal/mol	13199,737	13384,890	-13527,271	-13626,375	-13682,162	13694,412	13663
	J/mol	55227,699	56002,379	-56598,101	-57012,753	-57246,165	57297,419	57166

Table 4

Values of the logarithm of the equilibrium constant of the ammonia decomposition reaction at different temperatures.

T1, K	373	473	573	673	773	873
LOGKT1	-1,2411	+0,1918	+1,1587	+1,8587	+2,3913	+2,8104
T1, K	973	1073	1173	1273	1373	1473
LOGKT1	+3,1493	+3,4285	+3,7440	+3,8611	+4,0314	+4,1783

Table 5

The values of the change in the Gibbs energy of the reaction of decomposition of ammonia at different temperatures.



ΔGOT1	T1, K	373	473	573	673	773
	cal/mol	+2118,309	-415,141	-3038,167	-5724,142	-8458,621
	J/mol	+8862,984	-1736,95	-2711,69	-3949,81	-35390,87 2080

ΔGOT1	T1, K	873	973	1073	1173	1273	1373	1473
	кал/моль	1122	-14022,094	-16834,099	-20096,474	-22491,865	-25328,673	-28163,612
	Дж/моль	4697	-58668,441	-70433,87	-84083,647	-94105,963	-105975,16	-117836,55

- **Test methodology (Research Methodology).** According to the compiled program, we calculated the values of thermodynamic quantities: DHO298, DS298, GO298, QPT1, DHOT1, LOGKPT1, DGOT1 and etc. for temperatures from 873 to 1473K in steps of 100K. To do this, each time in operand 138, the value of T1 was changed to the corresponding temperature in degrees Kelvin.

Results and discussion (Analysis and results). The results of the above thermodynamic calculations using a computer, presented in table. 3 show that under standard conditions the reaction under study is endothermic, does not proceed spontaneously, and is thermodynamically impracticable.

The thermal effect of the reaction (Table 1) in the temperature range 873–1473 K is endothermic; the reaction proceeds with the absorption of heat from the outside.

Conclusion and proposals (Conclusion/Recommendations).




Fig.1. Dependence of the heat effect of the ammonia decomposition reaction on temperature based on new constants






Fig.2. Dependence of the logarithm of the equilibrium constant on the decomposition temperature of ammonia




Fig. 3. Dependence of the magnitude of the change in the Gibbs energy on the temperature of decomposition of ammonia

At 373K the value of the decimal logarithm had a negative sign (Table 3.4), and at 458K it was equal to zero. With a further increase in

temperature, a slight increase was observed and at 1073K the value of the decimal logarithm was - + 3.4285. At a temperature of 1073K, which is



optimal in the process of obtaining calcium cyanamide from calcium oxide, carbon dioxide and ammonia [119, p. 5], the value of the Gibbs energy change is -70433.87J/mol , which follows from Fig.3.3.

On the basis of the method used, thermodynamic calculations were carried out based on the old constants. According to the old physicochemical constants, the temperature of the onset of ammonia decomposition turned out to be 456K, and according to the new ones, it was 459K [119, p. 6].

Thus, the ammonia decomposition temperature of 459K can be considered more reliable, since it was determined according to new literature data.

2. Thermodynamically calculated values of the change in the Gibbs energy showed that their absolute values decrease with increasing temperature.

3. The results of computer thermodynamic calculations show that the thermal effect of the reaction in the temperature range of 873–1473 K is endothermic, the reaction proceeds with external heat absorption.

List of used literature

1. Toirov Z.K., Panjiev O.Kh., Bozorov O.N., Bobokulov A.N. "Chemical technology of inorganic substances." Textbook. - T. - "Philosophers". - 2018. - pp. 27-30.
2. Panzhiev A.X., Panzhiev O.X. Thermodynamic Studies of the Possibility of Free Carbon Formation during the Synthesis of Calcium Cyanamide by the Carbide-Free Method. International Journal of Progressive Sciences and Technologies (IJPSAT), International Journals of Sciences and High Technologies. 2 September. 2020. 111-116-pp
3. Panzhiev A.X., Panzhiev O.X. Kinetics of calcium Cyanamid obtaining process from lime, carbon dioxide and ammonia. International Journal on Integrated Education. Vol.3 № 9 (2020): IJIE. 260-263-pp
4. Panzhiev A.X., Panzhiev O.X. The expander gas and ammonia ratio influence on the calcium cyanamide yield. International Journal of Trend in Scientific Research and Development (IJPSAT), Special Issue on International Research Development and Scientific Excellence in Academic Life Available Online; www.ijtsrd.com e-ISSN; 2456-6470. 2021. January
5. Panzhiev A.X., Panzhiev O.X., Toirov Z.K. The impact of temperature on the syntheses of calcium cyanamide from ammonia, carbon dioxide and lime obtained from Jamansai limestone. International conference on integrated innovative development of Zarafshan region; Achievements, challenges and prospects 27-28 november, Uzbekistan 2019, 369-372-pp
6. Panzhiev A.Kh., Panzhiev O.Kh., Toirov Z.K. Influence of temperature on the synthesis of calcium cyanamide from ammonia, carbon dioxide of lime obtained from Jamakay limestone. UNIVERSUM scientific journal issue N2. (68), February 2020, Moscow, 68-71s
7. Panzhiev A.X., Panzhiev O.X., Norkulov M.B. The effect of temperature on the synthesis of calcium cyanamide and the composition of the exhaust gases. International engineering journal for research and development. Published; may- 28, 2020 353-362-pp
8. Panzhiev A.X., Panzhiev O.X. Thermodynamic Studies of the Possibility of Free Carbon Formation during the Synthesis of Calcium Cyanamide by the Carbide-Free Method. International Journal of Progressive Sciences and Technologies (IJPSAT), International Journals of Sciences and High Technologies. 2 September. 2020. 111-116-pp



9. Panzhiev A.X., Panzhiev O.X. Kinetics of calcium Cyanamid obtaining process frocess from lime, carbon dioxide and

ammonia. International Journal on Integrated Education. Vol. 3 № 9 (2020): IJIE. 260-263-pp.

