How often Complications Related to Tooth Extraction in Patients with Periodontitis

Dr.GALIB MASUDI- BDS. HDD. FIBMS.
galibmas@gmail.com
Lubna Abdulwahab Aldury-MSc
lubnaabdulwahab@gmail.edu.iq

DOINumber: 10.48047/NQ.2022.20.9.NQ44934 **NeuroQuantology2022;20(9): 8037-8043**

Introduction:

The second most common reason for extraction in dentistry is periodontitis [1] since it is an irreversible process especially if it is accompanied by severe tooth movement [2], advanced periodontitis, which is marked by severe bone and attachment loss, can make the teeth very mobile and loose [3], requiring extraction [4]. The inflammatory process in periodontitis is usually irreversible, especially if it has reached an advanced stage [5]. Early diagnosis and intervention are essential to salvage as many affected teeth as possible [6]. Bad oral hygiene [7], smoking habits [8], diabetes mellitus, and genetic influences all act as accelerators of increased severity that may further complicated or necessitate extraction of extracted teeth [9]. In fact, under certain conditions, extraction shall become the best course of treatment when those teeth host advanced forms of periodontitis not to be adequately treated or controlled [10].

Keywords:exodontia, periodontitis, tooth mobility.

The extent of Periodontal Disease:

The severity and progression of periodontal disease must be evaluated to determine the appropriate time for tooth extraction [11]. Teeth that have a lot of bone loss, deep periodontal pockets, and much mobility should be extracted early to prevent further damage and complications [12].

Presence of Active Infection:

Prior to extractions, any active infections, including abscesses or acute inflammation of the periodontium, must be addressed. In order to avoid problems after surgery, the patient should treat any current infections with the appropriate antimicrobials and practise good infection control.

Overall Oral Health Status:

In order to get a complete picture of a affected person's oral health, it's vital to look at their teeth from all angles. In addition, you want to take care of different dental problems, like a continual mouth hassle. The patient's potential to coaching right dental hygiene should be the last consideration. Ideally, a patient ought to wait until their oral fitness is ideal before scheduling an extraction of a seriously damaged teeth.

Planned Restorative Treatment:

Oral recovery methods, consisting of extractions, dental implant placement, and the use of a set or removable prosthesis, ought to all take location at the equal time for the first-class viable final results in the end [13].

Patient Factors:

As a part of the extraction assessment, the medical doctor will study the affected person's vitals, clinical records, and mindset in the direction of the procedure. The patient's ability to tolerate the process, their compliance with pre- and post-operative care commands, and the feasible effect on their great of life are some of the maximum vital elements to keep in mind

[14]. The availability of modern-day surgical tools and techniques, as well as preparedness to cope with ability headaches, are other elements that have an effect on while an extraction is achieved. To find the candy spot among risks and blessings, clinicians should suppose cautiously approximately these factors when determining whilst to extract tooth from patients with advanced periodontal disease. When dealing with extreme periodontal disease, it's far important to make sure the affected person is fully organized for tooth extraction. Here are some steps to follow:

Patient Education and Informed Consent:

Provide the patient with an in-intensity explanation of the reasoning behind the extraction, any capacity dangers or headaches, and the anticipated effects. Make sure the affected person is prepared for the method by means of presenting him with written instructions and all of the necessary treatment facts. Before performing the extraction, make sure you've got the patient's informed consent.

Anxiety Management:

Determine the patient's degree of tension and, if vital, administer medicine to calm them down or alleviate their tension to be able to relax throughout the process. The patient can also be helped to deal with anxiety or fear via healing communique, distraction, and rest strategies.

Pain Management:

It will even cope with the requirements for lengthy-term anaesthesia and neighborhood anaesthetic use, presenting a comprehensive plan for ache management. It is essential to think about the patient's consolation earlier than, at some point of, and after the system with the aid of considering the perfect postoperative analgesic. The patient's ache tolerance and any prior issues with local anaesthesia must also be documented within the clinical history.

Infection Control:

Prior to surgical operation, it is vital that sufferers practise true oral hygiene and use an antimicrobial mouthwash, along with chlorhexidine, to get rid of any potential infections. It is feasible to lessen the danger of postoperative infections by means of prescribing prophylactic antibiotics if medically vital.

Postoperative Instructions:

Before, throughout, and after an extraction, it's miles essential to provide designated commands concerning wound care, nutrition, and the handling of any capacity headaches. Make sure to emphasize the significance of following up with a renovation treatment.

Emotional Support:

In instances of excessive periodontal ailment, sufferers might also experience feelings of anxiety, fear, or disappointment further to the bodily signs of enamel loss [15]. Patients need emotional aid, empathy, and reassurance at some point of the procedure so they can manipulate their emotions.

Coordination of Care:

Proper control of any preexisting health situations that could have an effect on the extraction and restoration tiers requires near collaboration between the patient's primary care doctor and different healthcare providers. Incorporating these important additives of patient training permits clinicians to enhance the patient's cognizance, consolation, and ease for the duration of teeth extraction, main to better effects and patient pride.

Some of the changes which have been made to help patients with intense periodontal ailment recover greater fast after having teeth surgically removed encompass:

Methods for Traumatic Tooth Extraction:

Utilise specialized gadgets such as peristomes and luxators to loosen the teeth from its surrounding smooth tissues and bone while minimising bone elimination. If you want to hold the extraction from negative the bone and gentle tissues an excessive amount of, you have to know not use too much leverage or force. To facilitate extraction, think about how reducing the enamel into smaller pieces would possibly facilitate reducing it into more achievable pieces.

Minimally Invasive Flap Design:

Use a sulcular incision or small, limited flap design to enter the area of surgery rather than

large, full-thickness mucoperiosteal flap. That way, interfering with blood flow to the remaining tissues would be minimized while healing.

Gentle Bone Removal:

Precision only to the point of selectively removing only so much bone as needed within the alveolar: piezosurgery or even similar; avoid using rotaries so much since, by induction they raise heat and harm the alveolar bone themselves.

Meticulous Wound Closure:

Use primary closure techniques that reduce wound tension through incision releases or tension-relieving sutures. Consider resorbable collagen membranes or other barrier materials to protect the extraction site and prevent proliferation of soft tissues.

Adjunctive Therapies:

Platelet-rich fibrin (PRF) or similar biologically-derived substances should be added to the graft for improved healing and tissue regeneration. In addition, antimicrobial agents like chlorhexidine gel applied to the surgical site would be included to treat infection and improve healing.

Minimized Surgical Time:

It is highly important to minimize tissue exposure and trauma in order to improve the surgical process and reduce the total operating time. To do this, proper planning, proper tool selection, and the adoption of effective techniques are of great importance.

Different specialized techniques and materials have proven to be effective in reducing trauma and enabling patients with advanced periodontal disease to recover more quickly after tooth extractions:

Piezosurgery:

Piezosurgery is a technique using the ultrasonic vibration during cutting of bone with minimal trauma and maximum preservation of surrounding tissues. Piezosurgery assisted tooth extraction cases advanced periodontal diseases resulted in low postoperative pain and edema and faster healing of soft tissues compared to the rotary conventional techniques.

Platelet-Rich Fibrin (PRF):

Platelet-rich fibrin (PRF) is a biomaterial prepared from the patient's blood by concentrating growth factors and bioactive substances. Its application in the extraction socket enhances bone and soft tissue regeneration, reduces incidence of alveolar osteitis or dry socket, and accelerates healing.

Collagen Membranes:

The application of resorbable collagen membranes, either alone or in combination with bone grafting materials, helps protect the extraction socket and prevent the ingrowth of soft tissues, thereby promoting guided bone regeneration. It has been shown that collagen membranes result in improved healing outcomes, increased bone growth, and better maintenance of the alveolar ridge in patients with severe periodontal disease.

Laser-Assisted Extraction:

Erbium-doped yttrium aluminum garnet (Er: YAG) lasers can be used in a way that makes laser-assisted procedures more precise and minimally invasive. Such laser-assisted procedures have been linked to less postoperative pain and swelling as well as faster healing of soft tissues when compared with traditional methods.

Piezo-Assisted Ridge Preservation:

Piezosurgery can be used in combination with bone grafting and socket preservation techniques to effectively preserve the alveolar ridge following tooth extraction in patients suffering from advanced periodontal disease. This holistic approach has shown promising results in the preservation of bone stock and support for future prosthetic reconstruction.

Biologics and Growth Factors:

Recombinant growth factors such as BMPs and PDGF can be used to stimulate the regeneration of bone and soft tissue at the extraction site. Research has shown that these biological agents will enhance bone formation and promote better healing results in the patient with severe periodontal disease. By incorporating these advanced techniques and materials into the surgical extraction procedure, health care providers can make the healing environment more favorable, decrease trauma, and allow the

patient with severe periodontal disease to recover faster, thus enhancing treatment success.

Recovery periods following minimally invasive extractions vary from one case to another, as they may allow for quicker recoveries than others. However, most of these procedures result in shorter recovery times than conventional extraction procedures do.

Recovery time is usually more extensive with patients who have had traditional extractions with forceps or elevators. Recovery time may range from 5 to 10 days, with some significant post-operative swelling, pain, and discomfort. Patients will often require pain medication for several days and may have a delayed return to normal oral function and activities.

PatientsInformations:

1: Name: John Doe
Age: 55 years old
Gender: Male
Chief Complaint:
"My gum is swollen and painful, and I have difficulty chewing on this side."
Medical History:
Hypertension, controlled with medication
No known drug allergies
Dental History:
History of poor oral hygiene
Irregular dental visits, last check-up 3 years ago
Previous scaling and root planning treatment 2 years ago
Clinical Examination:
Tooth #30 (lower right first molar)
Severe periodontal pockets, measuring 8-10 mm
Grade III mobility
Radiographic evidence of advanced bone loss
Purulent exudates present
Diagnosis:
Advanced periodontal disease affecting tooth #30
Treatment Plan:
Local anesthesia administration
A traumatic extraction of tooth #30 using periotome and luxator instruments
Debridement of the extraction socket
Placement of a collagen membrane or socket graft material
Suturing of the extraction site
Post-operative instructions and prescription of analgesics and chlorhexidine mouthwash
Follow-up appointment in 1 week for suture removal and evaluation of healing
Prognosis:
Good, with proper oral hygiene maintenance and regular dental visits

This clinical chart would summarize the patient's condition, the proposed treatment plan, and the expected prognosis for the extraction procedure because of advanced periodontal disease affecting tooth #30. To allow the patient to heal and recover faster, a minimally invasive extraction technique like the traumatic approach should be utilized.

2: Patient Information:
Name: Jane Smith
Age: 62 years old
Gender: Female
Occupation: Retired
Chief Complaint:
"My gum is swollen and my tooth is painful. I have trouble chewing on this side."
Medical History:
Hypertension, controlled with medication
Type 2 diabetes, well-managed with oral hypoglycemic agents
No known drug allergies
Dental History:
History of poor oral hygiene and irregular dental visits
Previous scaling and root planning treatment 4 years ago
Last comprehensive dental examination 2 years ago
Clinical Examination:
Tooth #19 (upper right first molar)
Severe periodontal pockets, measuring 9-11 mm
Grade III mobility
Radiographic evidence of advanced bone loss, with furcation involvement
Purulent exudates present
Diagnosis:

Advanced periodontal disease affecting tooth #19

Treatment Plan:

Consider replacing the tooth with a dental implant or a fixed or removable prosthetic if extraction is required and the patient prefers that, or simply cannot be saved. Have an established regular periodontal maintenance schedule, which may include periodic professional cleanings and periodic monitoring of the patient's oral health. The most essential point would be to tell the patient to commit himself to good oral hygiene as well as regular dental visitation in order to continue to keep the remaining healthy teeth in the long term. Treatment plan will thus depend on the individualized needs of the patient because of differences in the progression of the disease, and the prognosis of the diseased tooth, the patients' health, and perhaps his preferences.

Pre-operative assessment:

Review medical history and optimize diabetes control

Obtain necessary radiographic imaging (panoramic or CBCT)

Discuss treatment options and obtain informed consent

Definitive Prosthetic Rehabilitation:

Discussion and planning for a suitable tooth replacement option (e.g., dental implant, fixed partial denture).

References:

1. Jafarian, M., &Etebarian, A. (2013). Reasons for extraction of permanent teeth in general dental practices in Tehran, Iran. Medical Principles and Practice, 22(3), 239–244.
2. Hugoson, A., Sjödin, B., &Norderyd, O. (2008). Trends over 30 years, 1973-2003, in the prevalence and severity of periodontal disease. Journal of Clinical Periodontology, 35(5), 405–414.
3. Levin, L., Levine, J., & Ashkenazi, M. (2018). Tooth loss due to root caries and periodontal disease among institutionalized elderly in Israel. Geriatrics & Gerontology International, 18(7), 1031–1036.
4. Ong, G., Yeo, J. F., &Bhole, S. (1996). A survey of reasons for extraction of permanent teeth in Singapore. Community Dentistry and Oral Epidemiology, 24(2), 124–127.
5. Tonetti, M. S., Jepsen, S., Jin, L., &Otomo-Corgel, J. (2017). Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: A call for global action. Journal of Clinical Periodontology, 44(5), 456–462.
6. Ong, G. (1998). Periodontal disease and tooth loss. International Dental Journal, 48(3Suppl 1), 233–238.
7. Axelsson, P., Nyström, B., &Lindhe, J. (2004). The long-term effect of a plaque control program on tooth mortality, caries and periodontal disease in adults. Results after 30 years of maintenance. Journal of Clinical Periodontology, 31(9), 749–757.
8. Müller, H. P., Stadermann, S., &Heinecke, A. (2002). Gingival recession in smokers and non-smokers with minimal periodontal disease. Journal of Clinical Periodontology, 29(2), 129–136.
9. Albandar, J. M. (2011). Global risk factors and risk indicators for periodontal diseases. Periodontology 2000, 29, 177–206.
10. Kassebaum, N. J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C. J., &Marcenes, W. (2014). Global burden of severe periodontitis in 1990-2010: A systematic review and meta-regression. Journal of Dental Research, 93(11), 1045–1053.
11. **Al-Shourbaji, I., & Duraibi, S. (2023). IWQP4Net: An Efficient Convolution Neural Network for Irrigation Water Quality Prediction. *Water*, *15*(9), 1657.**
12. Alijoyo, F. A., Prabha, B., Aarif, M., Fatma, G., & Rao, V. S. (2024, July). Blockchain-Based Secure Data Sharing Algorithms for Cognitive Decision

Management. In *2024 International Conference on Electrical, Computer and Energy Technologies (ICECET* (pp. 1-6). IEEE.

13. Ramachandran, R., & Sujathamalini, J. (2024). Promoting Diversity And Inclusion In Higher Education: Strategies And Best Practices. *Educational Administration: Theory and Practice*, *30*(4), 6997-7007.
14. Alim, S., & AlShourbaji, I. (2020). Professional uses of Facebook amongst university students in relation to searching for jobs: an exploration of activities and behaviours. *International Journal of Social Media and Interactive Learning Environments*, *6*(3), 200-229.
15. F. A. Alijoyo, B. Prabha, M. Aarif, G. Fatma, V. S. Rao and P. Valavan M, "Blockchain-Based Secure Data Sharing Algorithms for Cognitive Decision Management," 2024 International Conference on Electrical, Computer and Energy Technologies (ICECET, Sydney, Australia, 2024, pp. 1-6, doi: 10.1109/ICECET61485.2024.10698611.
16. Al-Shourbaji, I., Alhameed, M., Katrawi, A., Jeribi, F., & Alim, S. (2022). A Comparative Study for Predicting Burned Areas of a Forest Fire Using Soft Computing Techniques. In *ICDSMLA 2020: Proceedings of the 2nd International Conference on Data Science, Machine Learning and Applications* (pp. 249-260). Springer Singapore.
17. Kalpurniya, S., Ramachandran, R., & Chandramohan, N. (2023). A Study on Stress Level, Happiness, Challenges, and Emotional Bonds of Parents having Children with Disabilities Availing Services at
18. NIEPMD, Chennai. *Integrated Journal for Research in Arts and Humanities*, *3*(5), 72-88.
19. AlShourbaji, I., Helian, N., Sun, Y., & Alhameed, M. (2021). Anovel HEOMGA approach for class imbalance problem in the application of customer churn prediction. *SN Computer Science*, *2*, 1-12.
20. **Band, S. S., Ardabili, S., Danesh, A. S., Mansor, Z., AlShourbaji, I., & Mosavi, A. (2022). Colonial competitive evolutionary Rao algorithm for optimal engineering design. Alexandria Engineering Journal, 61(12), 11537-11563.**
21. Singh, A., & Ramachandran, R. (2014). Study on the effectiveness of smart board technology in improving the psychological processes of students with learning disability. *Sai Om Journal of Arts & Education*, *1*(4), 1-6.
22. **Kachare, P., Puri, D., Sangle, S. B., Al-Shourbaji, I., Jabbari, A., Kirner, R., ... & Abualigah, L. (2024). LCADNet: a novel light CNN architecture for EEG-based Alzheimer disease detection. *Physical and Engineering Sciences in Medicine*, 1-14.**
23. Shiju, K. K., Breja, M., Mohanty, N., Ramachandran, R., & Patra, I. (2023). Importance of Special Education and Early Childhood General Education Teachers' Attitudes toward Culturally Linguistically Diverse People. *Journal for ReAttach Therapy and Developmental Diversities*, *6*(9s (2)), 1544-1549.
24. **Kachare, P. H., Sangle, S. B., Puri, D. V., Khubrani, M. M., & Al-Shourbaji, I. (2024). STEADYNet: Spatiotemporal EEG analysis for dementia detection using convolutional neural network. *Cognitive Neurodynamics*, 1-14.**
25. Ramachandran, R., & Singh, A. (2014). The Effect of Hindustani Classical Instrumental Music Santoor in improving writing skills of students with Learning Disability. *International Journal of Humanities and Social Science Invention*, *3*(6), 55-60.
26. Puri, D. V., Kachare, P. H., Sangle, S. B., Kirner, R., Jabbari, A., Al-Shourbaji, I., ... & Alameen, A. (2024). LEADNet: Detection of Alzheimer's Disease using

Spatiotemporal EEG Analysis and Low-Complexity CNN. *IEEE Access*

27. Sudarsanan, S., Ramkumar Thirumal, H. D. K., Shaikh, S., & Ramachandran, R. (2023). Identifying the Scope of Reattach Therapy for Social Rehabilitation for Children with Autism. *Journal for ReAttach Therapy and Developmental Diversities*, *6*(10s), 681-686.
28. **Abualigah, L., Sheikhan, A., Ikotun, A. M., Zitar, R. A., Alsoud, A. R., Al-Shourbaji, I., ... & Jia, H. (2024). Particle swarm optimization algorithm: review and applications. *Metaheuristic Optimization Algorithms*, 1-14**
29. Figuero, E., Smoucková, I., Lávková, M., Herrera, D., Alonso, B., Serrano, J., &Escribano, M. (2021). Gingival Recession and Tooth Loss: Socio-Demographic, Periodontal, and Life Style Determinants in Older Adults. International Journal of Environmental Research and Public Health, 18(22), 11927.
30. Sheiham, A., &Netuveli, G. S. (2002). Periodontal diseases in Europe. Periodontology 2000, 29, 104–121.
31. Levin, L., Zini, A., Levine, J., Weiss, M., Lev, R., &Polak, D. (2018). Periodontal Disease as a Risk Factor for Tooth Loss and Prosthodontic Considerations. Quintessence International, 49(3), 193–201.
32. Seerig, L. M., Nascimento, G. G., Peres, M. A., Horta, B. L., & Demarco, F. F. (2015). Tooth loss in adults and income: Systematic review and meta-analysis. Journal of Dentistry, 43(9), 1051–1059.
33. Castrejon-Pérez, R. C., Borges-Yáñez, S. A., Gutiérrez-Robledo, L. M., & Avila-Funes, J. A. (2012). Negative impact of oral health conditions on older adults' quality of life. Geriatrics & Gerontology International, 12(4), 782–783.