



A PRELIMINARY STUDY ON VARIOUS PHYSIOLOGICAL PARAMETERS AND ESTIMATION OF TOTAL PHENOLIC, FAVONOIDAL CONTENTS OF HYDRO ALCOHOLIC EXTRACTS OF *Kalanchoe pinnata* AND *Ficusracemosa*

Nayakanti Bhasker Babu^{a,b*}, Donakonda Madhuri^b.

(a) Research Scholar JNTUA. Ananthapur, Andhrapradesh.

(b) Creative Educational Society's College of Pharmacy, JNTUA, Kurnool, AndhraPradesh, 518218.

*Presenting Author : bhasker.nayakanti@gmail.com

ABSTRACT

Kalanchoe pinnata (KP) is commonly known as Cathedral bells belongs to the family *Crassulaceae*. *Ficusracemosa* (FR), a well known standard-sized tree in the family *Moraceae*. Both the plants are collected from Kurnool, Andhrapradesh, and Authenticated by Dr. K. Venkata Ratnam, Assistant Professor, Department of Botany, Rayalaseema University, Kurnool. Voucher specimens were preserved in the herbarium with voucher numbers RU871, RU872. In the present study Leaves of both the plants are extracted with aqueous alcohol in 70:30 ratio, The Percentage yield was 3.9, 5.5 in KP and FR respectively. preliminary phytochemical screening is done and found various phytoconstituents like Alkaloids, Tannins, Flavonoids, Glycosides, Phenols and Terpenoids in both extracts, various physiological parameters like Loss on Drying, Total ash, Acid insoluble ash, Water soluble ash, Swelling Index, Foaming index were performed. The total phenolic contents were estimated as 54.85±0.149% w/w and 48.89±0.261 %w/w equivalent of gallic acid respectively. Flavonoid contents were found as 65.24±0.672mg/g and 46.32±2.273mg/g of quercitin equivalent respectively. My further studies are aiming to structural elucidation and the anticancer activities of specific compounds in both the extracts.

Keywords: *Kalanchoepinnata*, *Ficusracemosa*, *Crasulaceae*, *Moraceae*.

DOI Number: 10.48047/nq.2022.20.22.NQ10224

NeuroQuantology2022;20(22):2356-2364

2356

INTRODUCTION

Medicinal plants have been used in virtually all cultures as a source of medicine. The widespread use of herbal remedies and healthcare preparations, as those described in ancient texts and obtained from commonly used traditional herbs and medicinal plants has

been traced to the occurrence of natural products with medicinal properties. The use of traditional medicine and medicinal plants in most developing countries, as a normative basis for the maintainance of good health, has also been widely observed .



Kalanchoe is a succulent and perennial plant that grows 3-5 feet tall which is commonly known as 'air plant', it has tall hollow stems, fleshy dark green leaves that are distinctively scalloped and trimmed in red, and bell-like pendulous flowers belongs to the family Crassulaceae. *Kalanchoe* is botanically classified with two main Latin names which refer to the same plant: *Brophyllum pinnatum* and *Kalanchoe pinnatum*. *Kalanchoe* is a medicinal plant largely used in folk medicine for the treatment of kidney stones, gastric ulcers, pulmonary infection, rheumatoid arthritis etc. KP has become naturalized in temperate regions of Asia, Australia, New Zealand etc. In Hawaii it is regarded as an invasive species. It is also widely distributed in Philippines.

Ficus racemosa is an evergreen, moderate to large-sized spreading, lacriferous, deciduous tree 15-18m high, without prominent aerial roots belongs to the family Moraceae. It is popular medicinal plant present in India, it has long been used in Ayurveda, the ancient system of Indian medicine, for various diseases including diabetes. *Ficus* is an exceptionally large pan-tropical genus with over 700 species distributed widely throughout the warmer parts of Asia, Africa, America and Australia. It is commonly known as 'gular', and all parts of this plant are regarded medicinally important in Ayurveda and it has been used extensively in the treatment of biliary disorders, jaundice, dysentery, diabetes.

Breast cancer is the common cause of cancer death in women worldwide. Rates vary about five-fold around the world, but they are increasing in regions that until recently had low rates of the disease. Many of the established risk factors are linked to oestrogens. Risk is increased by early menarche, late menopause, and obesity in postmenopausal women, and prospective studies have shown that high concentrations of endogenous oestradiol are associated with an increase in risk. It is categorized into 3 major subtypes based on the presence or absence of molecular markers for estrogen or progesterone receptors and human epidermal growth factor 2. More than 90% of

breast cancers are not metastatic at the time of diagnosis.

METHODOLOGY

All the plants collected were shade dried and powdered. The powdered material was initially extracted by maceration technique.

Plant Extraction:

The freshly collected leaves of *Kalanchoe Pinnata* and *Ficus racemosa* were shade dried and coarsely powdered. The powder was passed through sieve no.40. The sieved powder was stored in airtight container for further use. Initially, 100g of each dried plant material powder was macerated with hydro alcohol 70% ethanol for 7 days. It was then filtered and the solvent was evaporated and percentage yield was calculated and stored in desiccators until further use.

Phytochemical screening of *Kalanchoe Pinnata* and *Ficus racemosa* leaves

Preliminary phytochemical screening was carried out for detection of phytoconstituents.

Test for Alkaloids:

A small portion of the solvent free extract was stirred separately with few drops of dilute hydrochloric acid and filtered. The filtrate was tested with various reagents for the presence of alkaloids.

- A. **Dragendorff's reagent** (Potassium bismuth iodide solution): Reddish brown precipitate.
- B. **Mayer's reagent** (Potassium mercuric iodide solution): Cream colour precipitate.
- C. **Wagner's reagent** (Iodine potassium iodide solution): Reddish brown precipitate.
- D. **Hager's reagent** (Saturated solution of picric acid): Yellow precipitate.

Test for Carbohydrates:

- **Molisch's test:** To the test solution add few drops of alcoholic α -naphthol, then add few drops of concentrated sulphuric acid through the sides of test tube, purple to violet colour ring appears at the junction. It indicates the presence of carbohydrates.

- **Fehling's test:** To 1ml of Fehling's solution A (aqueous solution of Copper sulphate- CuSO_4) add 1ml of Fehling's solution B (solution of potassium tartrate) and 2ml of test solution. Then mix well and boil. Formation of red precipitate of cuprous oxide indicates the presence of carbohydrates.
- **Benedict's test:** To 1ml of sample add 2ml of Benedict's reagent (a solution of sodium citrate and sodium carbonate mixed with a solution of copper sulphate) followed by heating in boiling water for 3 minutes and not more than 7 minutes. Formation of reddish precipitate within 3 minutes indicates the presence of complex sugars and brown / yellow precipitate indicates presence of simple sugars.
- **Barfoed's Test:** The extracts were treated with Barfoed's reagent and heated. Appearance of reddish orange colour precipitate indicates the presence of non-reducing sugars.
Test for Proteins:
- **Biuret's Test:** When the extracts were treated with copper sulphate solution, followed by addition of sodium hydroxide solution, appearance of violet colour indicates the presence of proteins.
- **Million's Test:** When the extracts were treated with Million's reagent, appearance of pink colour indicates the presence of proteins.
Test for Steroids:
- **Liebermann Burchard Test:** When the extracts were treated with concentrated sulphuric acid, few drops of glacial acetic acid, followed by the addition of acetic anhydride, there is a formation of violet ring in between the two layers, and the appearance of green colour in the aqueous upper layer indicates the presence of steroids.
- **Salkowski test:** Treat the extract with few drops of concentrated sulphuric acid, red colour at lower layer indicates presence of steroids and formation of yellow colored

lower layer indicates presence of triterpenoids.

- **Sulfur powder test:** Add small amount of sulfur powder to the test solution, it sinks at the bottom. It indicates the presence of steroids.

Test for Sterols: When the extracts were treated with 5% potassium hydroxide solution, appearance of pink colour indicates the presence of sterols.

Test for Phenols:

When the extracts were treated with neutral ferric chloride solution, the appearance of violet colour indicates the presence of phenols.

When the extracts were treated with 10% sodium chloride solution, the appearance of cream colour indicates the presence of phenols.

Test for Gums and Mucilage:

The extracts were treated with 25ml of absolute alcohol, and then solution was filtered. The filtrate was examined for its swelling properties.

Test for Glycosides:

General Test:

Test-A: Extract 200mg of drug with 5ml of dilute sulphuric acid by warming on a water bath. Filter it. Then neutralize the acid extract with 5% solution of sodium hydroxide. Add 0.1 ml of Fehling's solution A and B until it becomes alkaline (test with pH paper) and heat on a water bath for 2 minutes. Note the quantity of red precipitate formed and compare it with that of formed in test B.

Test-B: Extract 200mg of drug using 5ml of water instead of sulphuric acid. After boiling add equal amount of water as used for sodium hydroxide in the above test. Add 0.1ml Fehling's solution A and B until alkaline (test with pH paper) and heat on water bath for 2 minutes. Note the quantity of red precipitate formed. Compare the quantity of precipitate formed in the Test-B with that of formed in the Test-A. If the precipitate in Test-A is greater than in Test-B then glycosides are present.



Since, Test-B represents the amount of free reducing sugar already present in the crude drug, where as test A represents free reducing sugar plus those related on acid hydrolysis of any glycoside in the crude drug.

When a pinch of the extracts were dissolved in the glacial acetic acid and few drops of ferric chloride solution was added, followed by the addition of concentrated sulphuric acid, formation of red ring at the junction of two liquids indicates the presence of glycosides.

Test for Saponins:

- **Foam Test:** 1ml of extracts are diluted to 20 ml of distilled water and shaken well in a test tube. The formation of foam in the upper part of the test tube indicates the presence of saponins.

Test for Terpenes: When the extracts were treated with tin and thionyl chloride, appearance of pink colour indicates the presence of terpenes.

Test for Tannins:

- **Ferric chloride test:** Treat the extract with ferric chloride solution, blue colour appears if hydrolysable tannins are present and green colour appears if condensed tannins are present.
- **Gelatin test:** To the test solution add 1% gelatin solution containing 10% sodium chloride. Precipitate is formed. It indicates the presence of tannins.

Test for Flavonoids:

- 5ml of extract solution was hydrolyzed with 10%v/v sulphuric acid and cooled. Then, it was extracted with diethyl ether and divided into three portions in three separate test tubes. 1ml of diluted sodium carbonate, 1ml of 0.1N sodium hydroxide, and 1ml of strong ammonia solution were added to the first, second and third test tubes respectively. In each test tube, development of yellow colour demonstrated the presence of flavonoids.
- **Shinoda test:** To the test solution add few magnesium turnings and concentrated hydrochloric acid drop wise, pink scarlet,

crimson red or occasionally green to blue colour appears after few minutes.

- **Zinc hydrochloride test:** To the test solution add a mixture of zinc dust and concentrated hydrochloric acid. It gives red colour after few minutes which indicate the presence of flavonoids.
- **Alkaline reagent test:** to the test solution add few drops of sodium hydroxide solution, intense yellow colour is formed which turns to colourless on addition of few drops of dilute acid indicates the presence of flavonoids.

Physicochemical Parameters

The ash values and extractives values were performed according to the official methods described in the Indian pharmacopeia and WHO guidelines on quality control methods -for medicinal plant materials (Kokate CK. et al., 2007; India Pharmacopoeia. 1996).

Determination of Total Ash

About 2 to 3 grams (accurately weighed) of plant powder was taken in a silica crucible previously ignited and weighed. It was incinerated by gradually increasing the heat not exceeding dull red heat (450°C) until free from carbon, cooled and weighed. The percentage of ash was calculated with reference to the air dried powder

Determination of Acid Insoluble Ash

The total ash was boiled for 5 minutes with 10% w/v dilute hydrochloric acid and filtered through an ash less filter paper (Whatmann no. 41). The filter paper was ignited in the silica crucible, cooled and acid insoluble ash was weighed.



Determination of Alcohol Soluble Extractive

Plant powders were macerated with 100 ml of alcohol of the specified strength in a closed flask for 24 hrs, shaking frequently during 6 hrs and allowing it to stand for 18 hrs. It was filtered rapidly taking precautions against loss of alcohol and 25 ml of the filtrate was evaporated to dryness in a tarred bottomed shallow dish at 105°C and weighed. The percentage of alcohol soluble extractive was calculated with reference to the air dried powder.

Determination of Water Soluble Extractive

About 5 g of the each powder was added to 50 ml of water at 80°C and to it 2 gm of Keiselghur was added and filtered. 5 ml of the filtrate was transferred to a tarred evaporating dish, the solvent was evaporated on a water bath, drying was continued for half an hour, finally it was dried in a hot air oven for 2 hours and weighed. The percentage of water soluble extractive was calculated with reference to air dried drug.

Loss on drying

The Loss on Drying Test is designed to

measure the amount of water and volatile matters in a sample when the sample is dried under specified conditions.

The weighing bottle was dried for about 30 min and cooled in the desiccator and weighed it again accurately. 2gms of sample was taken into weighing bottle and the sample was spread as a thin layer less than 5mm and weighed it accurately and placed in oven dried at a temperature of about 10 degrees for 1/2hr and cooled and weighed until constant weight was obtained.

Swelling index

The swelling index is the volume in ml taken up by the swelling of 1gm of herbal material under specified conditions. Its determination is based on the addition of water or a swelling agent as specified. About 2gms of powdered drug was taken in a glass stoppered measuring cylinder and added water of about 25ml and shaken repeatedly for about 1hour and allowed to stand it for 24hours and volume in ml is read is measured. The procedure is repeated for two times.

2360

The swelling index was calculated by using the formula

$$\text{Final weight-initial weight Swelling index} = \text{-----}$$

Final weight

Foaming index

The foaming ability of an aqueous decoction of herbal materials and their extracts is measured in terms of a foaming index. About 1gram of powdered drug was taken in to a 500ml conical flask and 100ml of boiling water was added to it and maintained at moderate boiling for 30 min collected and filtered into 100ml conical flask. The decoction was taken into 10 stoppered test tubes add it in successive portions of 1ml, 2ml,.....10ml and the volume is adjusted to 10ml and the test tubes was stoppered and shaken for 15 seconds lengthwise 2 shakes per second and allowed to stand for 15 min and the length of foam was measured.

Foaming index was calculated using the

following formula: Foaming index = 1000/a
Where, a = the volume in ml of the decoction used for preparing the dilution in the tube where foaming to a height of 1 cm is observed.

Quantitative estimation of Phytochemicals

Determination of Total Phenolics

The amount of total phenolic content of the extracts was determined by Folin-Ciocalteu reagent as oxidizing agent, gallic acid as standard. Exactly 0.1 ml of the extract was transferred to a 100 ml Erlenmeyer flask and the final volume was adjusted to 46 ml by addition of distilled water. 1 ml of Folin-Ciocalteu reagent diluted 10 times was added and incubated at room temperature for 3 min. 3



ml of 2% sodium carbonate solution was added and the mixture was shaken on a shaker for 2 h at room temperature. The absorbance was measured at 760 nm. Gallic acid was used as the standard (250-15.625 µg/ml) for a calibration curve. The phenolic compound content was expressed in terms of mg/ml (Nagulendhran et al., 2007).

Estimation of Total Flavonoids

Aluminum chloride colorimetric method was used for flavonoids determination. One milliliter of sample was mixed with 3 ml of

methanol, 200 µl of 10% aluminum chloride, 200 µl ml of 1 M potassium acetate and 5.6 ml of distilled water and remains at room temperature for 30 min. The absorbance of the reaction mixture was measured at 420 nm with ultraviolet (UV) visible spectrophotometer. The content was determined from extrapolation of calibration curve which was prepared by using quercetin solution (6.25-100 µg/ml) in methanol. The concentration of flavonoid was expressed in terms of mg/ml (Akinpeluet al., 2010).

Results and Discussion

Phytochemical screening of *Kalanchoe Pinnata* and *Ficusracemosa* leaves

Test	Hydroalcoholic extract of <i>Kalanchoe Pinnata</i>	Hydroalcoholic extract of <i>Ficusracemosa</i>
Alkaloids	+	+
Carbohydrates	+	+
Tannins	+	+
Flavonoids	+	+
Steroids & Terpenoids	+	+
Glycosides	+	++
Saponins	+	+
Phenols	+	+
Proteins	-	-
Fixed oils & Fats	+	-

2361

+indicates Presence -- indicates Absence

Percentage yield of plants

The percentage yield of the extracts were found to be 3.9 and 5.5 for KP, FR respectively all

extracts were semisolid in nature so, further freeze or spray drying is necessary nature for KP and FR are green in color.

Table: Percentage yield of plants

Plant Material	Solvent used	Percentage yield	Nature
<i>Kalanchoe Pinnata</i>	Hydro-alcohol (70:30)	3.9%	semisolid
<i>Ficusracemosa</i>	Hydro-alcohol (70:30)	5.5%	semisolid

Physiological Parameters

The physiological characters for the both of the plants were shown in given table. All are found to within limits as per ayurvedic pharmacopeia

Table : Different physiological parameters of *Kalanchoe Pinnata* and *Ficusracemosa* leaves



Parameters	Plants powders	
	<i>Kalanchoe Pinnata</i>	<i>Ficusracemosa</i>
Loss on drying	6.74	9
Total ash	17.66	16.7
Acid insoluble Ash	8.16	9.4
Water soluble Ash	9.61	5.2
%CFC	13.5	36.1
Swelling Index	4	8
Foaming index	7	9


Total phenolic content: Total phenolic content in the *KP leaves* extract was showed to 54.85 ± 0.149 %w/w equivalent of gallic acid and *FR* was 48.89 ± 0.261 %w/w equivalent of gallic acid respectively.

Total flavonoid content: Estimation of total flavonoid content of *KP* has showed high total flavonol content 65.24 ± 0.672 mg/g of quercetin equivalent and *FR* showed total flavonol content is 46.32 ± 2.273 mg/g

Calibration data for total phenolic contents

Gallic acid concentration ($\mu\text{g/mL}$)	Absorbance (Mean \pm S.D)
20	0.543 ± 0.008
40	0.782 ± 0.001
60	0.967 ± 0.007
80	1.398 ± 0.003
100	1.487 ± 0.002

2362



Estimation of total phenolic content of Hydroalcoholic extract of *Kalanchoe Pinnata* and *Ficusracemosa* leaves




Treatment	Concentration (µg/mL)	Total phenolic content (mg/g of gallic acid equivalent) (Mean ±S.D)
<i>Kalanchoe Pinnata</i>	1000	54.85±0.149
<i>Ficusracemosa</i>	1000	48.89±0.261

Results are mean of three values ± SD

Calibration data for total flavonoid contents

Quercetin concentration (µg/mL)	Absorbance (Mean ±S.D)
20	0.652±0.003
40	0.857±0.006
60	1.234±0.001
80	1.476±0.008
100	1.675±0.005



2363

Estimation of total flavonoid content of Hydroalcoholic extract of *Kalanchoe Pinnata* and *Ficusracemosa* leaves

Treatment	Concentration (µg/mL)	Total flavonoid content (mg/g of quercetin equivalent) (Mean ±S.D)
KP	1000	65.24±0.672
FR	1000	46.32±2.273

Results are mean of three values ± SD

Conclusion:

Kalanchoe pinnata and *Ficus racemosa* leaves are taken and extracted with Aqueous alcohol (70:30) by the process called Maceration. The preliminary phytochemical screening was done and found to

be many secondary metabolites like Alkaloids, Glycosides, Flavonoids, Carbohydrates, Tannins, Steroids, Saponins and Tannins etc. The physicochemical parameters like LOD, Ash values like Total



Ash, Acid insoluble ash, Water soluble ash, Extractive values, Swelling index and Foaming index

etc., Total Phenolic and Flavonoidal studies are done. These plants are well known and have source

of curing ailments for time immemorial. In future studies the isolated principles needs to be evaluated

in scientific manner using specific animal models. Clinical studies are needed to confirm the actions.

Science Communication and Information Resources, Council of Scientific and Industrial Research, New Delhi; 1997:31.

7. Plants.usda.gov

8. www.raintree.com

9. S Huissain and DK Hore, Collection and conversation of major medicinal plants of Darjeeling and Sikkim, Indian Journal of Traditional Knowledge, 2007;6(2):352-357.

10. DonatusEbereOkwuet.al, Exotic Multifaceted Medicinal Plants of drug and pharmaceutical industries, African Journal of Biotechnology, 2009;8(25). 7271-7282.

ACKNOWLEDGEMENTS:

I Thank my Guide, Principal of Creative Educational society's college of pharmacy for providing infrastructure and facilities to carry out the research work. I also thank college management for their funding of my research work.

REFERENCES:

1. QuaziMajaz.A et al, The Miracle Plant (*Kalanchoe pinnata*): A Phytochemical and Pharmacological Review, IJRAP, 2011, 2(5), 1470-1482.
2. H. Agarwal, Anti inflammatory activity screening of *Kalanchoe pinnata* methanol extract and its validation using a computational simulation approach, Informaties in medicine unlocked, 14, 2019, 6 – 11.
3. Faiyazahmed, Traditional uses, medicinal properties and phytopharmacology of *Ficus racemosa*. A review, 2009, 48(6), 672-681.
4. Adrienne G. Waks et al, Breast Cancer Treatment A Review, Jama Journal of The American Medical Association, 2019, 321 (3), 288-300.
5. Kritikar K.R. & Basu B.D., Indian Medicinal Plants with illustrations, Volume-5, second edition, Oriental enterprises. 2003: 1394-1396
6. The wealth of India, A Dictionary of Indian Raw Materials and Industrial products, Vol 5, First supplement series, National Institute of

