Formulation and Evaluation of Herbal Gel from Zingiber Officinale and Garlic (Allium Sativum) for Anti-Inflammatory Action and Relieving Muscular Stress

Amulya Gupta, Dr. Pushpendra Jain
Department of Pharmaceutics, IIMT college of pharmacy knowledge park III, Plot No. 20-A Greater Noida 201310, Affiliated with A.K.T.U. Lucknow, 226031

Corresponding Author: Department of Pharmaceutics, IIMT College of Pharmacy, Dr. A.P.J. Abdul Kalam Technical University,Sec-11, Jankipuram, Vistar Yojna, 226031, Lucknow , India. E-mail address: amulyagupta283@gmail.com

ABSTRACT:

Aim of the study **:** The goal of the research is to create a topical medication delivery system for herbal formulations that has a deeper skin penetration potential and minimal systemic distribution for the treatment of inflammation and muscular tension.

Methodology : The powdered plant material of *Zingiber Officinale and Garlic (Allium Sativum)*were evaluated by standard procedure for the determination of following physicochemical parameters**.**Extract was prepared using Soxhlet apparatus.The base was preapared on the bases trial and error method by evaluating its phase separation, spreadability, easily removability and Non-irritant properties. The extracts were filtered and concentrated under reduced pressure. The final composition of water miscible base are given in formulation table. The dried methanol extract of both plants was mixed.To accomplish appropriate consistency of gel definition grouping of gelling specialist were improved and ready with various gel formers, Carbomer 934, HPMC , Carboxy methylcellulose sodium and different centralization of thickness enhancer were attempted lastly gel that showed great spread capacity and consistency was chosen.

Result: Physicochemical parameters like colour, odour, texture and the Consistency was found best of formulation (FM5). Optimized formulation (FM5) was selected on the basis of smooth texture, compatible viscosity & consistency, good spreadability&extrudibility, absence of foreign particles, non-irritant and stable in accelerated stability study over the period of a month. It was found to be the best among the all five formulations**.**

Conclusion: Further we can conclude that, this gel formulation can also be used to standardize the gel in anti-inflammatory. These elements may be used as lead compounds in the creation of potential topical herbal gel anti-inflammatory and anti-arthritic medications, according to the results of the overall investigations.

KEYWORDS: anti-inflammatory, Zingiber officinale,Allium Sativum, polyherbal gel**,**inflammation.

Number: 10.14704/nq.2022.20.9.NQ44316 **Neuro Quantology 2022; 20(9):2697-2706**

INTRODUCTION

Herbal medicine

The practice of herbal medicine is the oldest form of healthcare which has been used for decades in developing and developed countries. Primitive people have depended on nature for food, shelter, clothing and medicine to cure ailments. These humans distinguished useful herbs with beneficial effects from those that were inactive or toxic [1]. According to literature approximately 50,000 plant species are stated to have medicinal properties [2]. Thus, the basis of modern medicinal drugs such as aspirin, morphine, digitoxin and quinine were synthesized through scientific validation of herbal medicine [3]. Plant based drugs awareness advanced gradually and has been passed on, therefore setting a foundation for many traditional medicine systems around the globe [4].

Inflammation

Inflammation is part of the complex biological response of body tissues to harmful stimuli, such as pathogens, damaged cells, or irritants and is a protective response involving immune cells, blood vessels, and molecular mediators. The function of inflammation is to eliminate the initial cause of cell injury, clear out necrotic cells and tissues damaged from the original insult and the inflammatory process, and initiate tissue repair.[5]

Inflammation can be classified as either acute or chronic. Acute inflammation is the initial response of the body to harmful stimuli, and is achieved by the increased movement of plasma and leukocytes (in particular granulocytes) from the blood into the injured tissues. A series of biochemical events propagates and matures the inflammatory response, involving the local vascular system, the immune system, and various cells within the injured tissue. Prolonged inflammation, known as chronic inflammation, leads to a progressive shift in the type of cells present at the site of inflammation, such as mononuclear cells, and is characterized by simultaneous destruction and healing of the tissue from the inflammatory process.[6]

The accompanying side effects and markers may be among them: general feeling of sickness, exhaustion, and fever. These are side effects of an exceptionally dynamic safe framework that requires a great deal of energy, which may not be accessible for different undertakings. More safe framework cells and antibodies can be produced in the event that a fever expands the speed of digestion. modifications in the blood, like an expansion in immunological cells. Septicemia is an extremely unprecedented yet serious contamination related result (blood harming). This issue might appear as chills, feeling very unwell, or an extremely high temperature. [7]

Site of inflammation

An area of irritation might be confined to a solitary tissue or it might reach out over the whole body. Itis is a postfix used to assign different confined irritations, including vasculitis, hepatitis, myocarditis, an infected appendix, colitis, and dermatitis, to specify a couple. However, hardly any nearby aggravations, such asthma or pneumonia, don't stick to this norm. Because of physical closeness, relentless limited aggravation can once in a while prompt ongoing foundational irritation. For example, foundational provocative sickness is welcomed on by the development of bacterial biofilm close periodontal vasculature [8]. Like this, expanded degrees of provocative middle people in the blood animate vein endothelial cells, which supports the beginning of atherosclerosis.[9]

NSAIDs (Nonsteroidal anti-inflammatory drugs)

FDA-endorsed for use as antipyretic, mitigating, and pain relieving meds are nonsteroidal calming meds (NSAIDs). [10] These activities make NSAIDs helpful for dealing with messes like gout, headaches, pyrexia, dysmenorrhea, joint inflammation, and strong agony. They can likewise be utilized as narcotic saving medications in certain circumstances of extreme injury. [11-13]

As per their substance creation and selectivity, NSAIDs are regularly arranged into the accompanying gatherings: acetylated salicylates (anti-inflamatory medicine), non-acetylated salicylates (diflunisal, salsalate), propionic acids (naproxen, ibuprofen), acidic acids (diclofenac, indomethacin), enolic acids (meloxicam, piroxicam), anthran (celecoxib, etoricoxib).

Topical Drug Delivery System

Because of prescription entrance into the more deeply layers of the skin or mucous movies, skin drugs have exceptionally restricted impacts at the application site. The critical advantage of utilizing an effective getting sorted out approach is that it evades primary pass processing. Successful definitions additionally stay away from the dangers and costs of intravenous treatment as well as the numerous maintenance circumstances, for example, pH swings, protein presence, and stomach discharging spans.[14] All powerful conveyance, however it likewise utilizes froths, sprinkles, relieved powders, game plans, and startlingly calmed stick systems, is overwhelmed by serious areas of strength for semi in its designs. The cutaneous medicine conveyance framework is utilized when other solution the executives systems are ineffective. It is essentially utilized for help with discomfort, contraception, and urinary incontinence.[15-16]

Formulation approaches

An essential part of further developing penetration utilizing novel definition frameworks is the utilization of colloidal carriers. Dynamic molecules that have been caught are moved into the skin by submicron-sized particles. These carriers incorporate strong lipid nanoparticles, liposomes, and nanoemulsions. As per most of information, the carriers gather in the layer corneum or other upper skin layers. As a rule, it is far-fetched that these colloidal carriers will enter living skin. By the by, it is as yet dubious in the event that these carriers are powerful. [17,18]

MATERIALS AND METHODS:

Collection, Identification and Authentication of Plant Specimens

Due to their robust pharmacological effects, minimal toxicity, and commercial feasibility, plants have been studied for their therapeutic qualities globally in light of current scientific discoveries. The present study evaluates the activity of Zingiber Officinale and Garlic (Allium Sativum) extract in accordance with the data gathered from local vaidhyas and other traditional anti-inflammatory medicine practitioners.

The Zingiber Officinale and Allium Sativum plant materials were purchased from a neighbourhood market.

Garlic (Allium Sativum) and Zingiber Officinale components have been cleaned, rinsed, and dried over the course of six days. After drying, the plant material was combined into a coarse powder and stored at room temperature for further research.

Physicochemical Evaluation

The following physicochemical properties were determined by normal technique using the powdered plant material of Zingiber Officinale and Garlic (Allium Sativum).Extract was prepared using Soxhlet apparatus.

Loss on drying

Loss on drying is the lack of mass expressed as a percentage of mass. A Petri dish containing precisely 5–6g of medicine powder is placed in a hot-air broiler set at 105°C for 4-5 hours. After cooling in the dessicator, each situation's weight misfortune was recorded. This method was repeated until the desired weight was obtained. [19,20]

Loss on drying (%) = loss in weight X 100/ W

W= weight of the drugs in grams

Determination of Ash Value [21]

Ash value is a method for evaluating the purity and quality of unprocessed powdered pharmaceutical materials. Since the natural matter was burnt, which tells us more about the existence of the active ingredient in the unprocessed medicine, the debris is actually the rough material without the natural matter.[22]

Total ash value [23,24]

Take 2 to 3g of properly measured powdered separate, then place it in a platinum or silica plate that has been tarred, recently lit, and gauged. Disperse the drug in powder form on the bottom of the plate. Burn gradually increasing the heat while not exceeding dull red heat until released from carbon, then cool and measure. If a carbon-free debris cannot be obtained in this way, then burn the singed mass with warm water, collect the buildup on ashless channel paper, add the filtrate, spread the buildup, and light at low temperature. The vacant silica cauldron was immediately taken and put into the muffle heater with the help of a tong for a start at 6000C for 30 minutes. 2g of powdered drug was put to the silica cauldron after it had been removed and weighed. After that, it was placed in a silent heater set at 500–6000°C for 2–3 hours, or until it became white. finally weighed it. The total amount of debris was measured using an air-dried example.

Total ash value = (z-x/y) × 100

Where,

X = weight of the silica crucible

Y = weight of the drug powder (g)

Z = weight of the silica cruicible with powder ash

Acid-insoluble ash[25]

With about 30 ml of diluted hydrochloric corrosive, the debris was bubbled for 10 to 15 minutes, collecting the insoluble material in a cauldron. It was lit, gauged, and washed with warm water. For the air-dried medication, the amount of corrosive insoluble debris was measured. 25 cc of dil HCl were combined with the detritus obtained by the aforementioned procedure. For five minutes, bubble it. The accumulation was then placed on debris-free channel paper at that time. This was followed by

lighting it in a suppress heater at 5600C for 60 minutes. Regarding an air-dried sample, the amount of corrosive insoluble debris was assessed.

Acid insoluble ash value % = (a /y) × 100

where,

A = weight of the remaining residue

Y = weight of crude powder taken (g)

Water-soluble ash

The ash from the total ash value was boiled in 25 cc of water for 5 minutes. The insoluble material was added to filter paper with no ash. Then, a low temperature ignition was used to maintain a steady weight. By deducting the weight of water insoluble ash from the overall ash value, the weight of water soluble ash was calculated. With reference to air died value, the proportion of water soluble ash was computed.[26]

Extractive Value

Methodology In a closed cup, 5 gm of coarsely powdered, air-dried medication was macerated with 100 ml of soluble (chloroform, ethanol, and water) for 24 hours before being allowed to stand for 18 hours. After then, it was promptly divided in order to prevent liquor loss. A 25 ml sample of the filtrate was evaporated to dryness in a shallow dish that was lined with tartar, then dried at 1050°C and gauged. Regarding the air-dried medication, the amount of alcohol solvent extractive was determined. The quantity of dissolvable components used for extraction in a certain dissolvable is referred to as its extractive worth.[27]

Extractive value was determined using the formula

Extractive value(%) = (weight of residue)/(weight of dry powder) x 100

Formulation and development of Herbal Gel [28]

The base was preapared on the bases trial and error method by evaluating its phase separation, spreadability, easily removability and Non-irritant properties. The extracts were filtered and concentrated under reduced pressure. The final composition of water miscible base are given in formulation table. The dried methanol extract of both plants was mixed.

To accomplish appropriate consistency of gel definition grouping of gelling specialist were improved and ready with various gel formers, Carbomer 934, HPMC , Carboxy methylcellulose sodium and different centralization of thickness enhancer were attempted lastly gel that showed great spread capacity and consistency was chosen. Carbomer 934, HPMC and Carboxy Methyl cellulose sodium were utilized as thickness enhancer and upgraded for herbal gel plan. Gel was arranged containing methanol separate as per the equation given in Table underneath with various polymers in changing fixations.

Gelling specialist is the primary element of the definition. The centralization of thickness enhancer or gel previous is a significant boundary as a less focus will prompt straightforward arrangement or moisturizer with extremely low consistency, while high fixation might prompt development of gels with high consistency prompting non-uniform dissemination of medication and issue with treatment of gel.

S.no	Ingredients	FM1	FM2	FM3	FM4	FM5
1.	Methanol drug Extract	4	4	4	4	4
2.	HPMC	-	1	-	1.5	2
3.	Carbopol 934(g)	1	1.5	2.5	1	1.5
4.	Sodium hydroxide	1	1	1	1	1
5.	Propyleneglycol 400	4	4.5	4	4.5	4
6.	Carboxymethyl cellulose	1	1.5	2	3	2.5
7.	Purified water q.s	100	100	100	100	100

Table 1 Formulation and development of Herbal Gel

Physicochemical parameters evaluation of herbal gel

Extrudability

A closed, collapsible tube carrying a formulation was forcibly squeezed at the crimped end to ascertain its extrudability. Formula extruded once the cap was taken off until the pressure subsided. It was calculated the weight in grammes needed to extrude a 0.5 cm ribbon of the formulation in 10 seconds. It was recorded the average extrusion pressure in g.

pH

In 100 ml of pure water, 1.0 g of gel was properly measured and dispersed. Using a computerised pH metre that was calibrated at 4.0, 7.0, and 9.0 before usage with normal support arrangements, the pH of the scattering was determined. Threefold pH estimates were made, and typical characteristics were identified.[29]

To ensure that the detailing may be used without the risk of irritating the skin, the pH of the product is not fixed. Since the pH of the gel was estimated to be 6.6 0.5, which is extremely close to the neutral pH, the term may be used without risking skin irritation. This further proved that the chosen plan components did not alter the definition's pH.

Viscosity

Using a Brookfield viscometer, the viscosity of gels was measured (Spindle type S-24 at 30 rpm) The spindle was submerged in 200 g of gel for roughly 5 minutes before the reading was obtained.[30]

Spreadability

It should have excellent spreadability, which is one of the models for an effective detailing to satisfy the required features. It is a phrase used to describe the extent to which a plan quickly spreads after being applied to skin or another affected area. The usefulness of a term also depends on how well-known it is. On a glass slide that was 7.5 cm long, 0.5 g of gel was placed in a circle that was 1 cm in diameter and spaced apart to determine the spreadability of details. A second glass slide was then placed on top of the first. A 20 g weight was dragged across the upper 7.5 cm and then isolated from the bottom side, noting the length of time it took for the separation. Spreadability was assessed using.[31]

S = M.L / T

Where,

S = Spreadability

M = Weight tide to the upper slide

L = Length of a glass slide

T = Time taken to separate the slide completely from each other.

Homogeneity

After the generated gels were put in the container, they were all visually inspected for uniformity.[32]

Stability studies of topical herbal gel formulation

The major goal of the stability testing is to offer proof of how the medication product's quality changes over time under the impact of temperature and humidity. According to ICH requirements, a stability chamber was used for a three-month period to conduct the stability research for the topical herbal gel formulation

RESULT:

Organoleptic Properties Of Zingiber Officinaleand Garlic (Allium Sativum)

Organoleptic properties are the critical parameter for the rapid identification and consumer acceptance. Sensory evaluation-visual macroscopy, colour, odour, taste, fracture are the common features helped in identification of the crude drug. The organoleptic properties of **Zingiber officinale and Garlic (Allium Sativum)**have been mentioned in Table Below:

Table 5.1: The organoleptic properties of Zingiber officinale

S.NO	PROPERTIES	OUTCOME
1.	Colour	Light Brown
2.	Shape	Irregular
3.	Odour	Characteristic Odour
4.	Texture	Rough
5.	Taste	Pungent

Table 5.2: The organoleptic properties of Garlic (Allium Sativum)

S.NO	PROPERTIES	OUTCOME
1.	Colour	Whitish Yellow
2.	Shape	Irregular
3.	Odour	Strong Characteristic
4.	Texture	Coarse Powder
5.	Taste	Strong Pungent

Physicochemical Standardization of Proposed Plant Drug

Standardization of drugs means confirmation of its identity and determination of its quality and purity. The quality control standards of various medicinal plants, used in indigenous system of medicine, are significant nowadays in view of commercialization of formulations based on medicinal plants. The quality of herbal drugs is the sum of all factors, which contribute directly or indirectly to the safety, effectiveness, and acceptability of the product. Lack of quality control can affect the efficacy and safety of drugs that may lead to health problems in the consumers. Standardization of drugs is needed to overcome the problems of adulteration and is most developing field of research now. Therefore, there is an urgent need of standardized drugs having consistent quality.

Table 2 Standardization parameters of Zingiber Officinale and Garlic (Allium Sativum)

S.No	Parameters % w/w	Methanol ext	Methanol ext
1.	Ash value	7.45	6.74
2.	Water soluble matter	10.25	8.45
3.	Water soluble ash	3.65	3.21
4.	Acid insoluble ash	1.83	1.97

Table 3 Evaluation parameters for Methanol gel extract

Parameters	Formulation-1	Formulation-2	Formulation-3	Formulation-4	Formulation-5
Extrudability	++	+	++	+	+++
Spreadability	28.76	30.85	25.09	24.56	26.73
Viscosity	29430	31244	41150	42340	43213
Homogeneity	+	+++	+++	++	+++
Ph	6.3	6.2	6.2	6.3	6.2

+++ Very good, ++ good, + satisfactory



Figure 1: Spreadability of Methanol gel extract



Figure 2: Viscosity of Methanol gel extract



Figure 3: pH of Methanol gel extract

FUTURE PERSPECTIVE:

Despite the ample research in the area of for anti inflammatory use, very limited of them have reached to the clinical examinations, because of several boundaries in their developing phase, for instance, their physicochemical analysis, toxicity analysis, quality control, in-vitro testing, and optimization of several formulation development parameters. The research focus is key to aim the diseases in the posterior part of the eye via sustained /controlled release system. Furthermore, upcoming work should be focused to develop the formulation with low toxicity profile and better drug absorption. So, additional attention is needed for engineering therapeutics in continuing contemporary investigations. Researches in the field of novel polymers would help this resolution well. The polymer scientist canproduce a biocompatible and biodegradable carrier deprived of side-effects with patient suitability and affordability.

CONCLUSION:

Since the introduction of human development, restorative plants have been utilized to treat diseases.Medicinal plants are part and parcel of human life to combat diseases form the dawn of human civilization. Plants used in traditional medicine contain a wide range of phytocompounds that can be used to treat chronic infectious diseases. Various phytocompounds found in plants utilized in conventional medication can be used to treat ongoing irresistible diseases. Thus, Zingiber officinale and Garlic (Allium Sativum) both have a few dietary and phytochemical impacts that will without a doubt be useful in the treatment and counteraction of irritation and muscle uneasiness. Especially bioactive substances like flavanoids might be responsible for these plants' restorative impacts, which bring about clear physiological capabilities in the body. People ought to explore how conventional green sorts might be integrated into their ordinary eating routine and assist them with living sound, normal lives.

ACKNOWLEDGEMENTS:
I appreciate the administration of this excellent professional institution for giving the top lab facilities I needed to do my study. The conception and writing of this page were greatly assisted by all of the above authors.

CONFLICT OF INTEREST:

The Authors declare no conflict of interest.

REFERENCES:

1. Kunle OF, Egharevba HO, Ahmadu PO. Standardization of herbal medicines –A review. Journal of Biodiversity and Conservation. 2012;4(3):101-112. Herbal Medicine.
2. Barboza GE, Cantero JJ, Núñez C, Pacciaroni A, Espinar LA. Medicinal plants: A general review and a phytochemical and ethnopharmacological screening of the native Argentine Flora. Kurtziana. 2009;34:7-365. ISSN:1852-5962
3. Wachtel-Galor S, Benzie IFF. Herbal medicine. In: Benzie I, Wachtel-Galor S, editors. Herbal Medicine: Biomolecular and Clinical Aspects. 2nd ed. Boca Raton: CRC Press/ Taylor & Francis; 2011. ISBN: 13: 978-1-4398-0713-2
4. Pal SK, Shukla Y. Herbal medicine: Current status and the future. Asian Pacific Journal of Cancer Prevention. 2003;4:281-288.
5. Ferrero-Miliani L, Nielsen OH, Andersen PS, Girardin SE (February 2007). "Chronic inflammation: importance of NOD2 and NALP3 in interleukin-1beta generation". Clinical and Experimental Immunology. 147 (2): 227–35.
6. Chen, Linlin; Deng, Huidan; Cui, Hengmin; Fang, Jing; Zuo, Zhicai; Deng, Junliang; Li, Yinglun; Wang, Xun; Zhao, Ling (14 December 2017). "Inflammatory responses and inflammation-associated diseases in organs". Oncotarget. Impact Journals, LLC. 9 (6): 7204–7218.
7. Granger DN, Senchenkova E (2010). "Leukocyte–Endothelial Cell Adhesion". Inflammation and the Microcirculation. Integrated Systems Physiology—From Cell to Function. Morgan & Claypool Life Sciences
8. Nanci, A. and Bosshardt, D. D. Structure of periodontal tissues in health and disease. Periodontology 2000, 40: 11-28, 2006. DOI 10.1111/j.1600-0757.2005.00141.x
9. Seymour, G. J., Ford, P. J., Cullinan, M. P., Leishman, S., and Yamazaki, K. Relationship between periodontal infections and systemic disease. Clin Microbiol Infect, 13 Suppl 4: 3-10, 2007.
10. Phillips WJ, Currier BL. Analgesic pharmacology: II. Specific analgesics. J Am AcadOrthop Surg. 2004 Jul-Aug;12(4):221-33.
11. Dawood MY. Primary dysmenorrhea: advances in pathogenesis and management. Obstet Gynecol. 2006 Aug;108(2):428-41.
12. Shekelle PG, Newberry SJ, FitzGerald JD, Motala A, O'Hanlon CE, Tariq A, Okunogbe A, Han D, Shanman R. Management of Gout: A Systematic Review in Support of an American College of Physicians Clinical Practice Guideline. Ann Intern Med. 2017 Jan 03;166(1):37-51.
13. Oyler DR, Parli SE, Bernard AC, Chang PK, Procter LD, Harned ME. Nonopioid management of acute pain associated with trauma: Focus on pharmacologic options. J Trauma Acute Care Surg. 2015 Sep;79(3):475-83.
14. Surver, C. and Davis, F.A., Bioaviability and Bioequivalence, In Walter, K.A..(Ed.) , Dermatological and Transdermal Formulation, Marcal Dekker, INC. NewYork , 119,2002,pp. 403,323,326,327,403.
15. Bhowmik, Debjit. (2013). TOPICAL DRUG DELIVERY SYSTEM.
16. Singh Malik, D., Mital, N., & Kaur, G. (2016). Topical drug delivery systems: a patent review. Expert opinion on therapeutic patents, 26(2), 213–228.
17. Touitou, E., Dayan. N., Bergelson. L., Gidin, B., Eliaz, M., Ethosmes – novel vesicular carrier for enhanced delivery: characterization and skin penetration properties. J. Control Rel. 2000; 65: 405-418.
18. Sun, Y. M., Huang, J. J., Lin, F. C., Lal, J. Y., Biomaterials. 1997;18: 527 – 533.
19. Müller, Joachim &Heindl, Albert. (2006). Drying of Medicinal Plants. 10.1007/1-4020-5449-1_17.
20. Masud, Md & Hossain, Muhammad &Aktar, Morgina& Islam, Md & Hasan, Zakiul. (2014). Drying and Dehydration Kinetics of Ginger. 9. 2028-9324.
21. Kokate CK. Practical Pharmacognosy, 4th edition, Vallabh Prakashan. Delhi; 1994.
22. Ismail, B.P. (2017). Ash Content Determination. In: Food Analysis Laboratory Manual. Food Science Text Series. Springer, Cham.
23. Rao, Bayya. (2020). A Study on Ash Values and Pharmacopoeial Assay Methods in Herbal Pharmaceuticals.
24. Kim, D., Kim, B., Yun, E., Kim, J., Chae, Y., & Park, S. (2013). Statistical quality control of

total ash, acid-insoluble ash, loss on drying, and hazardous heavy metals contained in the component medicinal herbs of "Ssanghwatang", a widely used oriental formula in Korea. Journal of natural medicines, 67(1), 27–35.

25. Rao, Y., & Xiang, B. (2009). Determination of total ash and acid-insoluble ash of Chinese herbal medicine Prunellae Spica by near infrared spectroscopy. Yakugakuzasshi : Journal of the Pharmaceutical Society of Japan, 129(7), 881–886.
26. Kadam, Vasantrao. (2013). Determination of water soluble Ash values of some Medicinal Plants of Genus Sesbania. Journal of Pharmaceutical and Biological Research. 1. 1-4.
27. Pawar, Sushama& M.G., Jadhav. (2015). Determination of Extractive value and Phytochemical analysis.
28. Misal, G. & Dixit, G. &Gulkari, Vijay. (2012). Formulation and evaluation of herbal gel. Indian Journal of Natural Products and Resources. 3. 501-505.
29. Zhu, Y. J., & Chen, F. (2015). pH-responsive drug-delivery systems. Chemistry, an Asian journal, 10(2), 284–305.
30. Hancock, B. C., Dupuis, Y., &Thibert, R. (1999). Determination of the viscosity of an amorphous drug using thermomechanical analysis (TMA). Pharmaceutical research, 16(5), 672–675.
31. DeMartine, M. L., & Cussler, E. L. (1975). Predicting subjective spreadability, viscosity, and stickiness. Journal of pharmaceutical sciences, 64(6), 976–982.
32. Skibsted, E. T., Boelens, H. F., Westerhuis, J. A., Witte, D. T., &Smilde, A. K. (2006). Simple assessment of homogeneity in pharmaceutical mixing processes using a near-infrared reflectance probe and control charts. Journal of pharmaceutical and biomedical analysis, 41(1), 26–35.