



Awareness of Body Posture Among Students in A Private Dental Teaching Institution in Kanpur City, India – A Cross Sectional Study

Dr. Prateek Singh, Nidhi Shukla, Vaibhav Bansal, Surbhit Singh, Rohma Yusuf

Rama Dental College Hospital and Research Centre, Rama University, Mandhana, Kanpur, Uttar Pradesh, India

drprateek@ramauniversity.ac.in

Abstract

Background: This study was conducted to assess the postural awareness of dental students in A Private Dental Teaching Institution in Kanpur City, India.

Materials and Methods: A previously developed questionnaire was used for data collection in the study. The questionnaire was prepared by observing the positions of students working in the clinics and the mistakes they make with regard to their postures. The questionnaires were distributed among the dental students who were present and reported to work in the clinics. Levels of postural awareness and the relationship between postural awareness and the degree of musculoskeletal disorder (MSD) among the students was evaluated. This study was carried out in Rama Dental college, Hospital & Research Centre, Kanpur, India.

Statistical Analysis: Categorical variables were compared using Chi-square test. *P* values of less than 0.05 were considered statistically significant.

Results: A total of 184 dental students from the age group of 20–25 years were included in the survey, of which 144 dentists responded (83%). When their postural awareness was evaluated, results showed that 89% of the students had poor-to-medium levels of postural awareness. The relation between postural awareness and prevalence of MSDs indicated that 75% of the students with poor awareness, 49% of the students with average awareness, and 40% of the students with good awareness have MSDs. The results were statistically significant (0.002127, which is <0.005) stating that better awareness about proper postures while working helps to minimize the risk of MSDs.

Conclusion: Evaluation of levels of postural awareness showed that 21% of the students had poor postural awareness, 67% had average awareness, and 11% had good postural awareness. The analysis of results showed that those students with low-to-average postural awareness had significantly greater prevalence of MSDs.

DOI Number: 10.48047/nq.2019.17.10.2576

NeuroQuantology 2019; 17(10):09-12

BACKGROUND

Dental professionals commonly experience musculoskeletal pain during the course of their careers that may even cause a career-ending disability.¹ Dentists often assume static positions that are uncomfortable and asymmetric.² Historically, eISSN1303-5150

researchers have established that back pain is one of the most common problems among dental personnel.³ Prolonged shoulder flexion and upper arm abduction as well as high static muscle activity levels during common dental tasks were considered major factors



leading to neck and shoulder discomfort.⁴ Neck discopathy results in cervical pains or cervico-acromial pains, which are particularly common among dental practitioners.⁵⁻⁶ Movements such as craning, excessive bending and twisting of neck, bending forward from the waist, elevation of the shoulders, and general bending and twisting of neck and back are the common postural faults in dentists.

Some symptoms of musculoskeletal disorders (MSDs) are as follows: Excessive fatigue in the shoulders and neck; tingling, burning, or other pain in arms; weak grip; cramping of hands; numbness in fingers and hands; clumsiness and dropping of objects; and hypersensitivity in hands and fingers. Median nerve and cubital nerve defects are seen in a number of dentists due to defective ergonomics.⁷⁻⁹

Some signs of MSDs are decreased range of motion, loss of normal sensation, decreased grip strength, loss of normal movement, loss of coordination, etc.

MATERIALS AND METHODS

The survey with close-ended, self-administered questionnaires that focused on various positions while working was used as the data collection method. The questionnaire was prepared keeping in mind the various faulty positions of students working in the dental clinics and was aimed at drawing their attention to their strained positions that can potentially cause MSDs.

The questions were designed such that they evaluated the level of postural awareness by eliciting the details of the body posture of the

dentist, the manner of working (without rest or with short breaks), and the presence or absence of MSDs. Using the convenience sampling, the questionnaires were distributed to all the 184 male and female dental students present and working in the student clinics of Rama Dental College, Hospital & Research Centre, Kanpur. All the students were aged between 20 years and 25 years. Among the 184 students, 144 (83%) responded. **Statistical analysis:** The resultant data were subjected to statistical analysis. The level of knowledge of postural awareness was assessed and correlated with the presence or absence of MSDs. The categorical variables were compared using Chi-square test. *P* values of less than 0.05 were considered statistically significant.

RESULTS

A total of 184 dental students were given the questionnaire but only 144 responded and participated in the study giving a response rate of 83%. Of the 144 dental students who responded, 21.6% (29) had poor postural awareness (with a score less than 5), 67.16% (90) had average postural awareness (score of 5–7), and only 11.19% (15) had good postural awareness (score 8–10). About 45.5% participants were males and 54.5% were females. When the postural awareness was compared between male and female dental students, the *P* value was statistically insignificant (*P* = 0.18, i.e. >0.05) stating that postural awareness did not have any influence on the gender of the dentists [Table 1].

Level of postural awareness	Poor (%)	Average (%)	Good (%)
Number and percentage of students	29 (21.6)	90 (67.16)	15 (11.19)
Presence of musculoskeletal disorders	22 (75.86)	44 (48.88)	6 (40)

Table 1: Relationship between postural awareness levels and musculoskeletal disorder

The relationship between postural awareness and the degree of MSD among
 eISSN1303-5150

the students was evaluated, and the results are significant (<0.05) implying that better



awareness about the proper posture while working helps to eliminate the risk of MSDs. Eighteen percent of the surveyed dentists said they exercise regularly to prevent MSD and nearly 58% of the dentists with MSDs neglect and do not exercise regularly.

As many as 32% of the respondents claim that they are not suffering from any MSDs and thus do not require any kind of exercise. Some of dental students (67.16%) answered that they follow sitting dentistry most of the time and while working in the sitting posture, they bend forward slightly and work. They keep only the front portion of the foot touching the floor. They also responded saying that their backbone is never supported by the back of the sitting chair. All respondents said that they have no custom-made chair, but use the dental stool provided by the manufacturer. All of these responses strongly support that the awkward and static posture is the biggest risk factor for the MSDs.

DISCUSSION

Ergonomics is an applied science concerned with designing products and procedures for maximum efficiency and safety. According to Bethany Vallachi and others, dentists frequently assume static postures, which require more than 50% of the body's muscles to contract to hold the body motionless during longer breaks.^{1,9}

Postural awareness techniques are follows: Tilt the seat angle slightly forward up to 15° to increase the low back curve. Use a saddle-style operator stool that promotes the natural low back curve. Adjust the chair so that your hips are slightly higher than your knees and distribute your weight evenly by placing your feet firmly on the floor. Adjust the operator chair properly, adjust your chair first.

Lifestyle changes required¹⁰⁻¹² are as follows: The vast body of literature supports the idea that workers should vary their work positions as often as possible to shift the workload from one group of muscles to another and adapt to the dynamic postures. Aerobic exercises as the name suggest

concentrates more on improving oxygen transport by increasing blood flow to the tissues and thereby increasing their efficiency. Since the velocity of the blood flows increases, they also wash out blood triglycerides. The exercise program should contain warm-up, exercise period, and cool down. Finsen *et al.*¹³ suggested that an increased variation in work postures may reduce the risk of overloaded spine and lower and upper limbs. A study by Heuchet *al.*¹⁴ on dental students in Norway reported that a high body mass index was significantly associated with an increased prevalence of low back pain. Åkesson *et al.*¹⁵ observed that the work posture of dentists plays an important role as a risk factor for the development of work-related disorders. Therefore, postural awareness and lifestyle changes are essential for dentists. Postural awareness among dentists has been observed to be average according to some studies.¹⁶⁻¹⁸ The American Dental Association (ADA) has given valuable tips to improve the posture and reduce muscle fatigue in a clinical environment.¹⁹ The general agreement is to schedule patients so that you alternate between long, difficult cases and short, easier cases and use a surgical magnification system when necessary. It should allow for a comfortable posture while maintaining a close view of the task at hand. Adapt the lifestyle changes required for a healthy dental practice.

CONCLUSION

A majority of the surveyed dental students had average-to-poor postural awareness and a significant association was detected between postural awareness and Musculo skeletal Disorders. This study showed that those students with better postural awareness had reduced incidence of MSDs thus emphasizing the need for training programs aimed at increasing the awareness and knowledge of ergonomics.

REFERENCES

1. Valachi B, Valachi K. Mechanisms leading to musculoskeletal disorders in dentistry. J Am Dent Assoc
www.neuroquantology.com



- 2003;134:1344-50.
2. Kierklo A, Kobus A, Jaworska M, Botuliński B. Work-related musculoskeletal disorders among dentists – A questionnaire survey. *Ann Agric Environ Med* 2011;18:79-84.
 3. Abdul Samat R, Shafei MN, Yaacob NA, Yusoff A. Prevalence and associated factors of back pain among dental personnel in North-Eastern State of Malaysia. *Int J Collab Res Intern Med Public Health* 2011;3:576-86.
 4. Wai-Ping Lau J. Occupational shoulder pain in the dental profession-orthodontics, oral and maxillofacial surgeons, and general dental practitioners: Hong Kong Dent J 2006;3:29-36.
 9. Garbin AJ, Garbin CA, Diniz DG, Yarid SD. Dental students' knowledge of ergonomics postural requirements and their application during clinical care. *Eur J Dent Educ* 2011;15:31-5.
 10. Kisner C, Colby LA. Therapeutic exercise foundations and techniques. , PART II - Application of Therapeutic Exercise Techniques to Regions of the Body 6th ed. Ch. 9. New Delhi: Jaypee Brothers; 1996. p. 319-325.
 11. Milerad E, Ericson MO, Nisell R, Kilbom A. An electromyographic study of dental work *Ergonomics* 1991;34:953-62.
 12. Ergonomics and Dental Practice: Preventing work-related musculoskeletal problems Nov 01,2014.Available from:<http://www.ada.org/.ada/ergonomics-and-dental-practicing-preventing-work>.
 13. Finsen L, Christensen H, Bakke M. Musculoskeletal disorders among dentists and variation in dental work. *Appl Ergon* 1998;29:119-25.
 14. Heuch I, Hagen K, Heuch I, Nygaard Ø, Zwart JA. The impact of body mass index
 5. Gambhir RS, Singh G, Sharma S, Brar R, Kakar H. Occupational health hazards in current dental profession- A review. *The Open Occupational Health and Safety Journal* 2011;3:57-64.
 6. Vyshka GM. Genetics of pain and spinal cord stimulation: A review of the literature. *Webmed Central Neurology* 2011;2:WMC001635, 1-13.
 7. Sarkar PA, Shigli AL. Ergonomics in general dental practice. *Peoples J Sci Res* 2012;5:56-60.
 8. Nordander C, Ohlsson K, Akesson I, Arvidsson I, Balogh I, Hansson GA, *et al*. Risk of musculoskeletal disorders among females and males in repetitive/constrained work. *Ergonomics* 2009;52:1226-39.
 - on the prevalence of low back pain: The HUNT study. *Spine (Phila Pa 1976)* 2010;35:764-8.
 15. Åkesson I, Hansson GA, Balogh I, Moritz U, Skerfving S. Quantifying work load in neck, shoulders and wrists in female dentists. *Int Arch Occup Environ Health* 1997;69:461-74.
 16. Hutcheson CL. The dental professional's quiet health culprit. *Int J Orthod Milwaukee* 2004;15:31-2.
 17. Samotoi A, Moffat SM, Thomson WM. Musculoskeletal symptoms in New Zealand dental therapists: Prevalence and associated disability. *N Z Dent J* 2008;104:49-53; quiz 65.
 18. Salik Y, Ozcan A. Work-related musculoskeletal disorders: A survey of physical therapists in Izmir-Turkey. *BMC Musculoskeletal Disord* 2004;5:27.
 19. Ergonomics for dental students. Available from: <http://www.richmondinstitute.com/.Ergonomics-for-Dental-Students-2011.p.2-3>.