A STUDY OF AQUATIC INSECTS WITH THEIR ECOLOGY AND LIFE CYCLES

Amy Paul

Research Scholar, Department of Zoology, Gauhati University

Sankha Datta

Research Scholar, Department of Zoology, Gauhati University

ABSTRACT

The ecosystems' biodiversity heavily relies on insects' presence. "Aquatic insects" refers to insects that dwell mostly in freshwater habitats such as swamps, ponds, lakes, springs, streams, and rivers. Insects are found in many freshwater environments, with 8600 species in 12 orders and 150 families. They perform a vital ecological role in maintaining the health of freshwater habitats. Most freshwater ecosystems, including rivers and lakes, can support a variety of aquatic insect species. Hence there are many distinct types of aquatic insects. These species may be found in both lentic and lotic settings. Since they are abundant, reproduce quickly, have such huge biomass, and quickly colonize new freshwater habitats, aquatic insects are often used as models for studying freshwater ecosystem structure and function. Various variables impact aquatic biodiversities, such as industrial pollution and human activity.

Keywords: *Aquatic, Biodiversity, Conservation, Ecological indicator, Insects, life cycle*

DOI Number: 10.14704/nq.2022.20.9.NQ44598 Neuro Quantology 2022; 20(9):5169-5179

I.INTRODUCTION

Regarding diversity and abundance, insects are the most successful animal group in the animal world, making them the biggest and most diversified group of invertebrates. They outnumber all other animal species combined, and their diversity rivals that of plants and microorganisms. “Aquatic insects are a specialized subset of insects that have a wide range of adaptability, making them a more interesting and diversified species.” Various aquatic insect species can be found in micro and macro habitats because of their varied lifestyles. These include those that cling to the bottom substrates, those that crawl along protected surfaces, those that climb aquatic plant stems and other shoreline substrates, and those that burrow into the soft bottom (which are not associated with a substrate for attachment).

An aquatic habitat area of 3,166,414 Km2 with a large range in rainfall and altitude, topography, and latitude makes India one of the super-diverse nations. Freshwater insects comprise the most varied assemblage of creatures. Aquatic insect species originating from the fauna of North America, Australia, and Europe are estimated to number over 45,000 globally, with roughly 5,000 of these species inhabiting India's interior wetlands [1]. Ephemeroptera, dragonflies (Odonata), and caddisflies (Ephemeroptera) are some of the well-known aquatic insects found in inland wetlands (Trichoptera). It is not uncommon to see brightly colored aquatic insects such as dragonflies and damselflies (Odonata) in wetlands. This nutrient recycling process is facilitated by several aquatic insect feeding groups such as shredder, scraper, filter feeder, and predators [2]. Wood and leaf litter from the surrounding terrain are the primary food sources for aquatic insects. Fungal and bacterial activity further degrades the nutrients digested by aquatic insects into an absorbable form. Plants at the water's edge take up this nutrient-rich sludge. Aquatic insects play an important role in the environment and are an important food source for fish and amphibians.

Aquatic insects, classified into 12 families and numbering over 100,000 species, have been shown to support a freshwater environment and

to be an important part of the aquatic food chain. Orders like Diptera, Coleoptera, and Trichoptera, which account for 43%, 18%, and 15% of the total, are the most diversified. One or more phases of these insects' life cycles are spent inside the freshwater environment, where they provide critical ecological functions such as digesting organic materials, providing food for predators, and moving energy across trophic levels. Biological interactions among aquatic insects can significantly affect community structure in a freshwater ecosystem because of their role as links in the nutrient cycling process and as food sources for various functional feeding groups such as shredders, scrapers, gatherers, collectors-filter feeders, and predators. Aquatic organisms are very useful for biomonitoring strategies in aquatic systems because they are excellent indicators of the effects of human activity on the aquatic environment. Biodiversity and the function of aquatic insects in fisheries management and practice are further supported by the economic value of biodiversity and the role of aquatic insects. Indian rivers and lakes support a wide variety of aquatic invertebrates and other plant and animal species because of the diversity of the country's freshwater ecology. However, compared to other affluent nations, India has paid far less attention to research on freshwater creatures, particularly insects. Physico-chemical habitat characteristics were shown to significantly impact aquatic insect populations in Tamil Nadu, Kerala, Karnataka, and Andhra Pradesh, according to an extensive assessment of aquatic insect research. To analyze pesticide and heavy metal biomagnification, these studies extend the usefulness of aquatic insects in monitoring changes in water quality and aquatic life in freshwater settings. Toxicological research on aquatic insects includes a wide range of species. Insects like this one may help us determine whether or not a body of water is polluted.

Human activities may have a detrimental impact on terrestrial and aquatic ecosystems due to rapid rates of change in natural settings [3]. For example, riparian vegetation clearance, river damming or climate change all have the potential to alter water flow and frequency of flooding in lotic ecosystems [4]. The elimination of riparian vegetation and the linearization of watercourses reduces the space available for water infiltration and increases the incidence of flood and flash flood events [5]. The frequency and intensity of heavy precipitation events are projected to rise in tropical areas as precipitation becomes more concentrated over shorter time periods [6]. Changes in precipitation are projected to lead to an increase in disturbance events in lotic systems, which will alter flow patterns. For a better understanding of how biodiversity responds to environmental change, manipulative experiments are essential, since they can isolate the impacts of various disturbance sources.

To fully comprehend ecosystem dynamics and biodiversity patterns, it is critical to conduct research on the long-term effects of hydrological perturbations on aquatic habitats [7]. Hydrological disturbances (e.g., water withdrawal or flash floods) have historically been studied using taxonomic measurements like species richness and beta diversity indices [8]. According to a recent study, environmental perturbations may influence species in a variety of ways, depending on their functional features [9]. To understand biodiversity, one must look at how creatures interact with their environment, as well as how these interactions affect their identity and their ability to adapt to their surroundings [10].

In the event of a hydrologic disruption, traits such as permanence and recolonization are critical for aquatic populations. The resistance/resilience axis may accommodate qualities of persistence and recolonization in relation to the presence of the organism [11]. Resilience is distinct from resistance in that characteristic sets are distinct. It is possible for animals to maintain a stable population number in the face of perturbations if they have morphological adaptations for substrate fixation and body form [12]. The duration of a species' life cycle has a direct correlation to its resilience features, hence organisms with shorter life cycles are better equipped to colonise freshly disturbed habitats. There is a trade-off between strong resistance to disturbance and lower resilience in organisms, while low resistance to disturbance tends to improve resilience [13].

Communities may be shaped by functional features that are not directly connected to resistance or resilience (e.g., food acquisition). Hydrological disruptions may have an impact on

the importance of these other features on community organisation [14]. On the other hand, more disruption, either in severity or frequency, reveals resistance–resilience trade-offs. The value of features associated to competition in structuring communities in a particular environment is reduced when the environment is more disturbed. Aquatic communities tend to be organised based on features of resistance and resilience if environmental disturbance is considerable. We might anticipate organisms with varied features associated to resistance and resilience to respond differently to hydrological perturbations in these kinds of situations [15]." Resistance or resilience qualities are more favourable depending on the intensity and frequency of disruption.

It is possible to conduct manipulative research involving the effects of water flow on aquatic species in Cerrado biome streams. The Cerrado is one of the world's most endangered tropical biomes and one of the world's top five most biodiverse regions. The Cerrado's headwater streams are well known for their vulnerability to human-caused changes, especially those caused by shifting land uses. Large water resources in South America such as Prata River or Amazon River depend on this biome for its dynamic stability. Cerrado rivers provide 70% of Brazil's hydroelectric electricity, which is generated in the Cerrado territory. Storm surges, popularly known as "trombas d'água" (meaning "waterspouts"), cause streams in the Cerrado to undergo unusually brief hydrological pulses. As a result of climate change, extreme weather events such as wildfires, droughts, and flooding are predicted to become more common in the area [16]. The significance of extremely diversified headwater streams necessitates the use of monitoring networks that can indicate whether there is adequate data to identify any trends in reducing biodiversity.

An experiment was carried out in the Cerrado of Brazil to see whether altering stream flow affects the prevalence and colonisation of aquatic insects with varied functional features. Since aquatic insects have been studied for their functional qualities and abundance, we've decided to work with them because of their reliance on environmental circumstances.

As a result, aquatic insect testing is required for licensing any pesticide. The OECD has provided several recommendations for pesticide research in Daphnia species as part of the Chironomus immobilization test. Aside from their significance in biodiversity, insects may also be used as a pollution indicator. Stonefly populations declined due to a reduction in water oxygen, and Chironomus larvae due to an increase in temperature. Pesticides also hurt a variety of insects. The increase in insect population has a direct effect on pH, electric conductivity, and dissolved oxygen. Insects in the water were able to detect human contaminations. Insect bacterial growth suggests that the wetland habitat has high phosphate and nitrate levels. There was no pollution in the aquatic habitat, as shown by the abundance of the Gerrisspinolae population. There is no way to determine the extent of the ecosystem's contamination with chemicals alone; hence using a bioindicator is preferable. Seasonal fluctuations in the population of aquatic insects have a direct effect. The abundance of water in aquatic habitats makes them a popular target for human consumption. Forest fragmentation, the growth of agriculture, the building of dams, and mining have a significant impact on the vegetation area and faunal variety linked with it. It has been more common in recent years for uncontrolled and unregulated human activities to devastate natural regions by turning them into commercial plantations, resulting in a significant loss of plant biomass and variety. With their wide range of microhabitat distribution, high abundance, high fecundity, short generation period, and rapid colonization, aquatic insects are the greatest instruments for studying the freshwater ecosystem's structure and function. Before developing conservation and management plans for freshwater environments, further study is required to determine the health of aquatic insects.

II.ECOLOGY AND BIOLOGY OF AQUATIC INSECTS

The study of aquatic insects' ecology and biology is critical to understanding their involvement in water quality, disease ecology, climate change, biodiversity, community structure, and ecosystem functioning. Aquatic insect ecology and biology study have advanced significantly over the last century, allowing us to understand better their diversity, life cycles, and ability to

serve as ecological surrogates and energy sources [17]. It is very uncommon for aquatic insects to be found along the boundary between terrestrial environments (e.g., wetland areas) and freshwater ecosystems (e.g., wetlands). However, few aquatic insects are found in completely marine environments. These aquatic insects play a crucial role in freshwater ecosystems regarding nutrient cycling and total secondary production, depending on how humans have impacted the land around the waterways where they live. A wide range of aquatic insects contributes to the ecosystem's trophic structure, from detritivores to predators and food supplies for vertebrate and invertebrate predators. "This extends beyond the water's surface to the riparian area since many aquatic insects have aquatic (larval and adult) and terrestrial (adult) life phases, A Special Issue on "Ecology and Biology of Aquatic Insects" will focus on the most recent research on aquatic insect evolution, habitat partitioning, and community response to land use/land change, as well as their application as surrogates for the ecosystem, attributes such as pollution, and their use in the court of law, among other things". It has been a big challenge for aquatic insects to successfully adapt to their new environment because of the difficulty of breathing [18].

With the help of cutaneous and gill respiration and air extraction from plants, insects have overcome this problem. Molecular approaches and field ecology research may be utilized to understand better how these physiological systems have permitted unique ways for aquatic insects to respire and how they have adapted to flowing waterways and established control measures for pest species. Transcriptomic analysis may be utilized to manage a pest aquatic beetle species in this journal edition. More than 88,500 aquatic insects have been identified, spanning around 13 different orders [19]. Since taxonomy and fundamental biological and ecological investigations have long been recognized as intertwined, key groups of water insects, such as Plecoptera, Trichoptera, and even Neuroptera, remain mostly unknown. Ecological understanding concerning habitat usage or partitioning is lacking for many of these species, such as the Neuropteran family Sisyridae (Spongillaflies) in the aquatic Neuropteran family. Our knowledge of aquatic insect life cycles and their role in production and bioenergetics may be hindered by a lack of species-level information [17]. "To improve our ecological understanding of various life stages and life history features, this issue will contain research that addresses habitat usage and partitioning by two of these orders".

When it comes to aquatic insect communities, there is a wide range of variety and abundance that may be found, depending on environmental variables. These creatures can form separate communities based on their unique hydraulic, substrate, and oxygen needs. Several variables may influence the makeup of an insect population, including natural and artificial features of the environment. This is because insects play vital roles in the functioning of the ecosystem. Insect community composition in disease ecology is influenced by various factors, including stream flow and human effects. There might be a shift in the relative abundance of aquatic insect groups due to changes in aquatic habitats. Pollution effects on freshwater ecosystems may be assessed using aquatic insects as surrogates for water quality. Much aquatic research relies on the EPT (Ephemeroptera, Trichoptera, and Plecoptera) richness index as an example of this kind of surrogate analysis. The diversity of these three pollutant-intolerant orders is an important indicator of environmental health, water quality, and organisms' ability to respond to pollution [20]. It will be discussed in this issue how ecosystem traits are substituted in the rising human-mediated microplastic contamination. Fly-fishing, for example, is a leisure activity that draws on aquatic insect ecology and biology. These facets of human culture are all linked. Death scene investigations, specifically, are a new use that has recently emerged for the employment of aquatic insects by humans. "The use of aquatic Chironomidae in death investigations and how this family of aquatic insects may give vital information in determining a post-mortem submersion period will be evaluated in this issue (PMSI) [21]." Aquatic Insects: Ecology and Biology is a Special Issue that provides fresh empirical evidence and helps to untangle the aquatic insects' intricate functions in the aquatic community and ecosystem function.

III.EVOLUTION OF AQUATIC INSECTS

Although the origins of aquatic insects have been debated, it is still unclear if insects have evolved specifically for aquatic conditions or whether their adaptation to aquatic environments is secondary. Many scientists believe that millipede, centipede, and insect progenitors resided in leaf litter regions near the edges of ponds. We can trace our lineage to insects that evolved in wet environments [22]. Their Paleozoic fossil record goes back to the Devonian period. Insects with aquatic young are found solely in the Odonata and Ephemeroptera suborders of the genus Dragonfly. Insect evolution and phylogeny have been hindered by a lack of fossils of freshwater creatures. There are 12 orders of aquatic insects (Figure 1). Mayflies (Ephemeroptera), lacewings (Neuroptera), alder flies (Megaloptera), caddisflies (Trichoptera), moths (Lepidoptera), and wasps (Hymenoptera) are all aquatic, but their larvae are terrestrial adults. The aquatic beetles (Coleoptera) and bugs (Hemiptera) are all aquatic in their larval, nymphal, and adult phases.

IV.ROLE OF AQUATIC INSECTS IN FRESHWATER ECOSYSTEM

Because of their abundance, high birth rate, short generation time, enormous biomass, and quick colonisation of freshwater environments, aquatic insects may serve as model species for studying the structure and function of the freshwater ecosystem. Since aquatic insects spend the majority of their lives in or near water, any shift in the population's size or composition at a particular point in time or place might point to a shift in water quality. They may also be used as a gauge for aquatic ecosystem trophic structure, water quality, and eutrophication. Ecologically and economically, insects that live in the aquatic environment are a vital element of the ecosystem. In the study of ecology, population growth, genetics, and many other aspects of biology, they have been the major instrument. Dead leaves and other plant components that fall into bodies of water from the land are broken down by some of the aquatic insect life. In aquatic habitats, this substance serves as the foundation of the food chain. The algae that grows on rocks, logs, leaves, and stems of plants with roots may be scraped off using a metal scraper. Algae are more productive when they are thinned out by water insects and invertebrates, which provide oxygen and food to other species below. Species of aquatic insects are capable of removing microscopic particles from water. If you want to maintain the water clear enough for algae and other plants to thrive in, this is a great tool. As they dig for food, other aquatic insects stir the fragile bottom sediments. Adding oxygen from the water to the bottom gives it a better habitat for creatures.

Fish and ducks rely on chironomid larvae as a food source [23]. There is a wide variety of Chironomid species and their susceptibility to eutrophic conditions have been utilised to develop trophic categorization of lakes. Disease-carrying arthropods, such as dipteran flies, are ubiquitous across the world's ecosystems. Among the most common diseases in tropical regions is malaria. Anopheles mosquitoes are responsible for an estimated 500,000 cases of malaria each year, according to the CDC. A single mosquito, Aedes aegypti, transmits yellow fever. Aedes aegypti and A. albopictus are the mosquitoes that carry dengue and broken bone fever. Hemipterans are important predators for many wild and domesticated fish, but they may also be a problem in fish culture nursery ponds, where they prey on the young. Mosquito larvae may be controlled biologically by using certain insect families [25]. It's common in many cultures to consume and appreciate these insects. It's a sort of large water insect that Thais adore in stirfries and salads, for example.

V.AQUATIC INSECT LIFE CYCLES

Aquatic or semi-aquatic insects make for 7 percent of North America's 91,000+ species of insects. These creatures may be found in any kind of river. All aquatic insects have in common that they all depend on the water at some point. The bulk of waterway-blocking aquatic macroinvertebrates is the larval stage of an adult land-dwelling bug. Many aquatic insects begin their lives submerged in the ocean or other bodies of water. Immature aquatic bodies have evolved to better adapt to living in the water. Therefore they typically do not look like their adult land-based counterparts. The aquatic bug must undergo a series of physical transformations before leaving its watery

habitat and beginning its new existence on land. Metamorphosis is a term used to describe the transformation of a person's physical appearance. From one aquatic creature to the next, the degree of the transformation varies. Incomplete metamorphosis and full metamorphosis are two different kinds of transformation.



Almost 90% of the insects that undergo metamorphosis complete their transformation. Four steps are required for a complete transformation. As an egg, it develops into a larval (which frequently looks nothing like an adult), a pupa, and a cocoon, from which it emerges as an adult land bug. Insect larvae are generally very different from their adult counterparts. The bodies of many larvae resemble those of worms.

Some people are missing all three of their limbs. However, they will all ultimately mature into different-looking adults with wings that fly away, breed, and lay eggs to repeat the cycle. Aquatic insects that go through a full metamorphosis include the caddis fly, dobson fly, crane fly, mosquito, midge, and a wide variety of beetle species.

The metamorphosis that is not complete goes through just three phases. An insect's life cycle starts with an egg, which hatches into a nymph. The adult form is a tiny replica of the nymph, which frequently resembles the adult in appearance. On their backs, you may see wing pads or the beginnings of wings. It takes them a series of molts to grow to full size and produce wings from their wing pads. They are sexually immature at this point. They emerge from the canals as fully fledged creatures in their mature form, complete with wingtips. Mayflies, stoneflies, dragonflies, and damselflies are aquatic insects that do not complete their metamorphosis.

Various stages require a wide range of physical and behavioral modifications of aquatic insects. When it comes to where these phases occur, whether on land or in the water, insects may vary greatly. As a result, a thorough understanding of aquatic life's life cycles and phases is critical.

VI.AQUATIC INSECTS AND THEIR HABITATS

Running water (rivers and creeks) or stagnant water (ponds and lakes) are both suitable habitats for aquatic insects (ponds and lakes). Erosion-prone ecosystems like this one may be found on the shores of many lakes. Many depositional ecosystems may be found in floodplain pools and backwaters. Various geographical and temporal scales may illustrate the water insect habitats. “Particles vary in size from a few millimeters to the whole of the drainage basin, which covers an area of kilometers squared.” Habitat changes may be seen from a few days to tens of thousands of years. The permanence of its physical structures varies depending on how large the ecosystem is. Depending on the grain and microhabitat, this may take a few days to a few thousand years. This geographical and temporal fluctuation also affects insect communities in the wetlands. It is common for water insects to cling, swim, skate, or burrow their way into a habitat [26].

Flow, turbulence, and food availability all have a role in the distribution of aquatic insects in an environment. The frequency of movement in an environment is determined by a species' habit (method of locomotion, attachment, or hiding). The substratum, which is a crucial habitat physical component, is a very complicated one. The physical nature of the substrate is determined by the water current and the type of the available parental material. Complexity is added to the substrate by the organic debris, which may have a significant impact on the organism's reaction to the substrate. Fossil makeup varies throughout continents and biomes because of the environment. Low (rivers and creeks) or depositional water levels make sand a bad habitat (ponds and lakes). “Lake basins, river flood plain pools, and stream/river backwaters offer depositional circumstances, whereas stream/river currents and lake shoreline waves generate erosional habitats. Abundance and variety of species suited to erosional conditions [2].” Generally speaking, silty sand has high diversity, whereas muddy

substrata have low diversity and high biomass. There is a decrease in fauna due to sand and silt. It is recognized that the amount of area available for colonization impacts the number of species in stony substrates. With stable substrates and organic debris, the variety and quantity of organisms increase [27].

VII.SOCIETAL BENEFITS OF AQUATIC INSECTS

Humans and animals benefit from a wide range of aquatic bug species, but four, in particular, stand out. Biomonitoring, fishing, and noxious weed removal rely on these insects [28].

An introduction to the food web: When there are many aquatic insects, they ensure that enormous quantities of nutrients can be processed. Diversity (or taxon richness) of aquatic insects (e.g., nutrients, habitats) helps ensure that all available resources are well used in space and time.

Biomonitoring using aquatic insects: The systematic use of live organisms or their reactions to evaluate the health of aquatic habitat is known as "biomonitoring." In water quality assessments, fish, algae, protozoans, and other species are utilized, but macroinvertebrates, the majority of which are insects, are more usually used. In addition to their widespread distribution, the large number of species and their ability to respond to diverse forms of environmental stress make them ideal indicators of water quality and ecosystem health. They can also determine the extent of a perturbation's geographic reach, and their long life cycles make it possible to track changes over time [29].

There is possible to learn a lot about the health of a waterway by studying the aquatic insects that inhabit it. Biological, chemical, and physical variables impact aquatic insects, making them excellent water quality indicators. Short-term and long-term pollution episodes have a cumulative impact, and they cannot avoid. When it comes to things like dissolved oxygen content, they are quite finicky. Disturbances of many types are putting a growing strain on aquatic ecosystems [130].

In flowing rivers, mayflies are regarded as a "keystone" species because of their ability to serve as an early warning system for the existence of oligotrophic or mesotrophic conditions.

The saprophytic Diptera species suggests that water bodies are seriously contaminated since low oxygen levels and high nutrient concentrations are typical of poor water quality (eutrophic). Many pollution-tolerant chironomids sometimes indicate poor water quality (characterized by low dissolved oxygen and high nutrient concentrations). Great variety and low density are common characteristics of excellent water quality conditions. The presence of Chironomussp. in a water body shows that it is eutrophic. Using these bugs as an indicator for environmental contaminants is common since they can live in highly contaminated settings [31]. The whirling beetles of the Gyrinidae family (freshwater ponds, lakes, open-flowing streams, etc.) are discovered. It is common to see Haliplidae (water beetles) crawling through aquatic vegetation along the banks of wetlands and waterways such as lakes, rivers, or creeks.

Fishing for insects in the water Aquatic insect biodiversity is of particular interest to the general public because of the prevalence of these critters in the diets of many fish species, some of which are eaten by people. For a long time, anglers who use natural or synthetic baits have been interested in them. To fool fish into swallowing their hooks and being caught, anglers have sought to mimic the color and shape of numerous aquatic insects for ages. When it comes to mimicry, the most prevalent mimicry is that of the mayflies (caddisflies, stoneflies, and non-biting midges). Imitated forms of larvae, pupae, and adults are offered to fish as they develop, drift, emerge from the surface, or return to the water as egg-laying females or dying adults to mimic their behavior. Fish food may be found throughout the year and at various times of day if the stream has a wide variety of these insects, each having its unique emergence period.

Noxious weed control: Native species are being out-competitive, waterways and water intake systems are being jammed, and noxious invading plants are excluding feeding fish. Insect herbivores have successfully controlled these weeds, although herbicides are also often used. Three important herbivores in the United States have effectively reduced an invasive South American plant, the alligator weed

(Alternanthera philoxeroides): alligator weed flea beetles and alligator weed thrips. "Eichhornia crassipes (common water hyacinth), an invasive species from Brazil, was successfully controlled by two imported species of weevils and one species of imported moth. Weeds may be reduced or eliminated by the discovery of aquatic insect species."

Using aquatic insects in Forensic Entomology may be very helpful in drowning situations, such as the death of a person submerged for a long time. Anax Parthenope, Lestes Sponsa, Scarlet Skimmer, etc., are only a few aquatic insects that might be quite helpful in investigating a death. Several organizations can help examine submerged corpses and aquatic insects, particularly when working with natural water bodies. "True flies, or Diptera, are the most common insects in forensic examinations. Calliphoridae (blow flies), Sacrophagidae (meat flies), and Muscidae are the most common species in this group (house flies). Calliphoridae (blow flies) and Sacrophagidae (fleshflies) may arrive within minutes of death". Until the corpse reaches the bloat stage of decomposition, Muscidae (house flies) do not colonize [32].

VIII.APPROACHES TO AQUATIC INSECTS' CONSERVATION

The protection of freshwater systems on their whole is an important management goal. Headwater streams and high-elevation ponds are essential to protect endemic insects. River dynamics must be preserved, including seasonality and physicochemical parameters (Figure 4). "Consideration of river network connectivity, sensitive land use, topographical heterogeneity, and biotic interactions are important location-specific factors that need to be taken into account, as well as the promotion of macrophyte/riparian/bank vegetation quality for adult aquatic insects and protection of open water areas". Because it diminishes insect variety and promotes local hydrological dryness, channelization should be avoided at all costs. A good side effect of this is the creation of artificial, shallow, well-vegetated shorelines and the preservation of substrates that are rich in organic materials. Beavers, as well as other vertebrate engineers, should be rediscovered. Although non-native trees may occasionally fill in for the loss of native ones, alien trees must generally be cleared to enhance insect habitats significantly. For many terrestrial and aquatic insects, preserving or restoring intact hydrology and careful management of salty systems has become critical [33].

A landscape network consisting of several interconnected ponds relies on connectivity like other systems. There should be plenty of variety in pond size and shape, as well as high levels of functional connection between ponds and the deposition pools of nearby streams and rivers, to create high-quality pond shapes. A more diverse range of plants and insects may be achieved by preserving the natural dynamics of freshwater ecosystems. Due to the brief hydroperiods of certain aquatic insects, the landscape heterogeneity must include both permanent and ephemeral lakes and deposition pools of streams in order to maintain a diverse ecosystem. For certain aquatic insects, ephemeral ponds may be colonized from permanent ponds and pools. Around ponds, buffer strips have been established to reduce the impact of cultivation.

In addition, well-designed artificial ponds, high-quality irrigation ditches for marshland insects, and stormwater ponds for aquatic insects may offer essential supplemental habitat. Natural vegetation variability improves rural and urban ponds, with the additional advantage of raising insect conservation awareness in city ponds. Improved artificial ponds can considerably enhance the landscape's functional connection (i.e., improves the landscape). Many artificial environments, such as garden ponds, golf course roughs, and military training grounds provide excellent opportunities for aquatic insect conservation. Aquatic insect conservation is especially important in several unusual situations. "Biological control of invasive water plants, preservation of river-lake ecotones, rehabilitation of mining pools, retention of well-managed Sphagnum bogs, removal of alien predators such as fish, erection of physical diversion structures to deflect certain threatened flying adults, and in some special habitats reducing tourist impact through the use of designated paths and duckboards."

IX.CONCLUSION

Natural resource protection and biodiversity preservation have risen to the top of the priority

list in the quest for a more ecologically sound future. Due to a lack of data, many taxa have previously been precluded from being used as indicators. As more and more research is done on aquatic insect habitats and distributional patterns, their use is becoming more appropriate. "Because of their varying reactions to stimuli in their aquatic home and the ability to determine the quality of that environment, aquatic insects are often employed to monitor the health of aquatic habitats." Water entomologists are working hard to enhance and create novel biomonitoring methods that use aquatic insects.

REFERENCES

[1]. Amaravathi, D., Raja Sekhar, P.S. and Rama Rao, K. (2018). Entomofaunal diversity in Kondakarla freshwater lake ecosystem at Visakhapatnam, Andhra Pradesh, India. Journal of Entomology and Zoology Studies, 6(6): 880-886.

[2]. Subramanian, K.A. and Sivaramakrishnan, K.G. (2007). Aquatic Insects of India-A Field Guide. Ashoka Trust for Ecology and Environment (ATREE), Bangalore, India. pp. 50-62.

[3]. Secretariat of the Convention on Biological Diversity (2020). *Global Biodiversity Outlook 5*. Secretariat of the Convention on Biological Diversity.

[4]. Allan, J. D. (2004). Landscapes and riverscapes: The influence of land use on stream ecosystems. *Annual Review of Ecology Evolution and Systematics*, **35**, 257–284. https://doi.org/10.1146/annurev.ecolsys.35.12

[5]. Resh, V. H., Brown, A. V., Covich, A. P., Gurtz, M. E., Li, H. W., Minshall, G. W., Reice, S. R., Sheldon, A. L., Wallace, J. B., & Wissmar, R. C. (1988). The role of disturbance in stream ecology. *Journal of the North American Benthological Society*, **7**, 433–455. https://doi.org/10.2307/1467300

[6]. Arias, P. A., Bellouin, N., & Coppola, E. (2021). Chapter 4: Future global climate: Scenario-based projections and near-term information. In V. Masson-Delmotte, P. Zhai, & A. Pirani (Eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1–195). Cambridge University Press.

[7]. Lake, P. S. (2000). Disturbance, patchiness, and diversity in streams. *Journal of the North American Benthological Society*, **19**, 573–592. https://doi.org/10.2307/1468118

[8]. Death, R. G. (1996). The effect of patch disturbance on stream invertebrate community structure: The influence of disturbance history. *Oecologia*, **108**, 567–576. https://doi.org/10.1007/BF00333735

[9]. Menezes, S., Baird, D. J., & Soares, A. M. V. M. (2010). Beyond taxonomy: A review of macroinvertebrate trait-based community descriptors as tools for freshwater biomonitoring. *Journal of Applied Ecology*, **47**, 711–719. https://doi.org/10.1111/j.1365-2664.2010.01819.x

[10]. Verberk, W. C. E. P., van Noordwijk, C. G. E., & Hildrew, A. G. (2013). Delivering on a promise: Integrating species traits to transform descriptive community ecology into a predictive science. *Freshwater Science*, **32**, 531–547. https://doi.org/10.1899/12-092.1

[11]. sarremejane, R., Cid, N., Stubbington, R., Datry, T., Alp, M., Cañedo-Argüelles, M., Cordero-Rivera, A., Csabai, Z., Gutiérrez-Cánovas, C., Heino, J., Forcellini, M., Millán, A., Paillex, A., Pařil, P., Polášek, M., Tierno, J. M., de Figueroa, P., Usseglio-Polatera, C. Z.-M., & Bonada, N. (2020). DISPERSE: A trait database to assess the dispersal potential of aquatic macroinvertebrates. *Science Data*, **7**, 386. https://doi.org/10.6084/m9.figshare.13148333

[12]. Mazzucco, R., Van Nguyen, T., Kim, D.-H., Chon, T.-S., & Dieckmann, U. (2015). Adaptation of aquatic insects to the current flow in streams. *Ecological Modelling*, **310**, 143–152. https://doi.org/10.1016/j.ecolmodel.2015.04.01

[13]. Ferzoco, I. M. C. (2019). *Testing trade-offs in dispersal and competition in a guild of semi-aquatic backswimmers*. University of Toronto.

[14]. Wilkes, M. A., Edwards, F., Jones, J. I., Murphy, J. F., England, J., Friberg, N., Hering, D., Poff, N. L. R., Usseglio-Polatera, P., Verberk, W. C. E. P., Webb, J.,

& Brown, L. E. (2020). Trait-based ecology at large scales: Assessing functional trait correlations, phylogenetic constraints and spatial variability using open data. *Global Change Biology*, **26**, 7255–7267. https://doi.org/10.1111/gcb.15344

[15]. Kappes, H., Tackenberg, O., & Haase, P. (2014). Differences in dispersal- and colonization-related traits between taxa from the freshwater and the terrestrial realm. *Aquatic Ecology*, **48**, 73–83. https://doi.org/10.1007/s10452-013-9467-7

[16]. Arias, P. A., Bellouin, N., & Coppola, E. (2021). Chapter 4: Future global climate: Scenario-based projections and near-term information. In V. Masson-Delmotte, P. Zhai, & A. Pirani (Eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1–195). Cambridge University Press

[17]. Cummins, K.W.; Merritt, R.W.; Berg, M.B. Ecology and distribution of aquatic insects. In An Introduction of Aquatic Insects of North America, 5th ed.; Merritt, R.W., Cummins, K.W., Berg, M.B., Eds.; Kendall Hunt Publishing Company: Dubuque, IA, USA, 2019; pp. 117–140.

[18]. Huryn, A.D.; Wallace, J.B. Habitat, life history, secondary production, and behavioral adaptations of aquatic insects. In An Introduction of Aquatic Insects of North America, 5th ed.; Merritt, R.W., Cummins, K.W., Berg, M.B., Eds.; Kendall Hunt Publishing Company: Dubuque, IA, USA, 2019; pp. 65–116.

[19]. Starr, S.M.; Benstead, J.P.; Sponseller, R.A. Spatial and temporal organization of macroinvertebrate assemblages in a lowland floodplain ecosystem. Landsc. Ecol. 2014, 29, 1017–1031

[20]. Herman, M.R.; Nejadhashemi, A.P. A review of macroinvertebrate and fish-based stream health indices. Ecohydrology. Hydrobiol. 2015, 15, 53–67

[21]. Wallace, J.R.; Merritt, R.W. Chapter 7: The role of aquatic organisms in forensic investigations. In The Utility of Arthropods in Legal Investigations, 3rd ed.; Byrd, J.H., Tomberlin, J.K., Eds.; CRC Press: Boca Raton, FL, USA, 2020; pp. 155–186.

[22]. Sharma, K., Lodhi, R.K. and Rao, R.J. (2020). Study the diversity of aquatic insects in the ramada reservoir of Gwalior district (M.P.)." International Journal of Research-granthaalayah, 8(2): 140-146.

[23]. Cibrowski, J. J. H., and Corkum, L. D. (2003). United Earth Fund.Appendix 9: Sediment- zoobenthos interactions. In Evaluating Ecosystem Results of PCB Control Measures within the Detroit River–Western Lake Erie Basin : 78-82

[24]. Saether, O. (1975). Chironomid communities as water quality indicators. Holarctic Ecology, 2 : 65-74

[25]. Ohba, S. and Nakasuji, F (2006). Dietary items of predacious aquatic bugs (Nepoidea: Heteroptera) in Japanese wetlands. Limnology, 7 : 41-43

[26]. Hershey, A.E., Lamberti, G.A., Chaloner, D.T. and Northington, R.M. (2010). Aquatic insect ecology. In Ecology and classification of North American freshwater invertebrates, Academic Press, pp. 659-694

[27]. Collins, N.M. (Ed.). (2012). The conservation of insects and their habitats. Academic Press.Fonnesu, A., Sabetta, L. and Basset, A. 2005. Factors affecting macroinvertebrate distribution in a Mediterranean intermittent stream. Journal of Freshwater Ecology, 20: 641-647

[28]. Nair, G.A., Morse, J.C. and Marshall, S.A. (2015). Aquatic insects and their societal benefits and risks. Journal of Entomology and Zoology Studies, 3(3): 171-177

[29]. Bonada, N., Prat, N., Resh, V.H. and Statzner, B. (2006). Developments in aquatic insect biomonitoring: a comparative analysis of recent approaches. Annual Review of Entomology, 51: 495-523.

[30]. Tachet, H., Richoux, P., Bourneau, M. and Usseliopolatera, P. (2003). Invertebratesd‟ eaudouce; systematique, biologie, ecologie. Freshwater invertebrates; systematic, biology, ecology. CNRS edition, Paris.

[31]. Papacek, M. (2001). Small aquatic and ridiculous bugs (Heteroptera: Nepomorpha) as predators and prey: The question of economic importance. European Journal of Entomology, 98(1): 1-12

[32]. Joseph, I., Mathew, D.G., Sathyan, P., and Vargheese, G. (2011). The use of insects in forensic investigations: An overview of

forensic entomology. Journal of Forensic Dental Sciences, 3(2): 80-89

[33]. Samways, M.J., Barton, P.S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T. and Cardoso, P. (2020). Solutions for humanity on how to conserve insects. Biological Conservation, 242: 10-27