



Effectiveness of cinnamon, exercise and counselling on hyper androgenic symptoms and level of anxiety among young girls with Polycystic Ovarian Syndrome

Niranjani.S^{1*}, Dr.G. Bhuvaneshwari², Dr.M. Hemamalini³, Dr.R. Vijayaraghavan⁴

¹Ph.D scholar, Department of research and development, Saveetha Institute of Medical and Technical Sciences, (SIMATS), Chennai, Tamil Nadu, India.

niranmay26@gmail.com

²Guide / HOD and Professor, Department of Community Health Nursing, Saveetha College of Nursing, SIMATS, Chennai, Tamil Nadu, India

bhuvana.prabha1981@gmail.com

³Co-guide/ Principal and Professor, Hindu Mission College of Nursing, Tambaram West, Chennai, Tamilnadu, India.

hemasrini1979@yahoo.com

⁴Department of Research and Development, Saveetha Institute of Medical and Technical Sciences, (SIMATS), Chennai, Tamil Nadu, India

jai_vijay@hotmail.com

ABSTRACT:

9534

Background:

Polycystic ovarian syndrome (PCOS) is an endocrine metabolic, multifaceted disorder commonly diagnosed among young girls and adult women. Girls with polycystic ovarian syndrome often exhibit hyper androgenic symptoms like hirsutism, acne, alopecia and obesity. Young girls who are overweight & obese are very concerned about their physical appearance, which in turn reflects in their psychological wellbeing too. So it is very relevant to assess the level of anxiety related to body image disturbance among young girls with polycystic ovarian syndrome. Girls with polycystic ovarian syndrome can modify their lifestyle - including dietary adjustments, exercise, and behavioural alteration, which can ease the metabolic dysfunction. On the other hand, lifestyle modification has now been recognized as the first-line management for PCOS.

Aim: To determine the effectiveness of cinnamon, exercise and counselling in reducing hyper androgenic symptoms, BMI, waist circumference and anxiety level among young girls with PCOS

Methods: Quasi-experimental pre and post test design was used in this study. A total of 824 young girls from different colleges in Chennai were screened based on Rotterdam criteria and among that 150 study participants were selected who met the inclusion criteria. self-administered questionnaire was used to collect the baseline data. Modified Ferriman Gallwey scale, Ludwig visual score, Global acne grading scale were used to assess the hyper androgenic symptoms like hirsutism, alopecia and acne respectively. Other markers like BMI, Waist circumference and level of anxiety using Hamilton anxiety scale were also assessed. Participants were partitioned into three groups, 50 in each group - Non interventional group (only education regarding PCOS), Group A with all three interventions (cinnamon tea, exercise program and anxiety



reduction counselling), group B with selected interventions (exercise program and anxiety reduction counselling). Interventions are planned for 6 months. Post-test I, II were conducted after 3rd and 6th month respectively. Non probability Purposive sampling technique was used. Prior permissions were obtained from all institutional heads. consent was acquired from every one of the study participants.

Results: The present study unveiled the following results. In Modified Ferriman Gallwey scale (Hirsutism), the non – interventional group from Pre-test to Post-test 2 had shown no improvement in the score, whereas Group A and Group B shown 1 score decrease. In Ludwig visual score (alopecia), the non – interventional group from Pre-test to Post-test 2 had shown no improvement in the score, whereas Group A shown 2 scores decrease and Group B shown 1 score decrease. In Global acne grading scale (acne), the non – interventional group from Pre-test to Post-test 2 shown 1 score increase, whereas Group A and Group B shown 2 scores decrease. This shows that there is reduction in hyper androgenic symptoms like hirsutism, alopecia and acne in experimental groups. In anxiety scale, the non – interventional group from Pre-test to Post-test 2 had no improvement, whereas Group A and Group B shown 2 score decrease and this shows that there is reduction in the level of anxiety among experimental groups. For all these four scales, Kruskal Wallis one-way ANOVA on ranks shown significance ($P < 0.001$). Tukey’s multiple comparison test was carried out and both the experimental groups shown significant decrease. This shows that the Group A and Group B are better than the non – interventional group. This study also revealed that in BMI, the non-interventional group from Pre-test to Post-test 2, shown 1.7% increase in BMI, while in group A and group B, 4.4% and 3.02 % reduction in BMI was observed. This clearly shows that the group A is superior to group B and non-interventional group. In waist circumference the non-interventional group from Pre-test to Post-test 2, 0.3% increase in waist circumference was observed, while in group A and group B, 1.6% and 0.6 % reduction was observed. Hence it revealed that group A and group B are better than the Non interventional group. The Group A is superior to group B.

Conclusion: This study spotlighted the beneficial role of cinnamon tea, exercise program and anxiety reduction counselling in reducing the hyper androgenic symptoms like hirsutism, acne, alopecia, BMI, waist circumference and also the level of anxiety among young girls with Polycystic ovarian syndrome. Hence, the study suggests this multi intervention package which cost effective and less time consuming for young girls with PCOS.

Keywords: effectiveness, cinnamon, exercise, counselling, hyper androgenic symptoms, psychological markers, Polycystic ovarian syndrome, young girls.

DOI Number: 10.14704/nq.2022.20.8.NQ22973

NeuroQuantology 2022;20(8):9534-9544

INTRODUCTION:

Polycystic ovary syndrome (PCOS) is a condition where the women produces increased amount of male hormones. This higher male hormone in the body causes menstrual irregularities, infertility, excessive hair growth on the face and body, acne, hair loss and also obesity. In addition to that long-term health problems like diabetes, thyroid dysfunction and heart disease can also occur. These challenges to self-body image gradually develop anxiety, depression and low self-esteem among women with PCOS. Hence the poor quality of life in PCOS may affect the women’s desire to follow healthy eating habits, ability to exercise and change lifestyle. Therefore, it is also important to identify psychological aspects that are supportive for self-esteem and behavioural

change. PCOS affects women during their reproductive age between 15 to 44 years. 2.2 and 26.7 percent of women in this age group have PCOS. The prevalence rate of PCOS in Pondicherry was 12.18% (23), and 9.8% (24) in Thiruvananthapuram. These findings alarm the rising prevalence of PCOS in south India. Although the aetiology of PCOS is still a mystery, we believe that genetic and environmental factors contribute highly to this disorder. Evidences shown that lifestyle changes along with planned counselling programs are now been recognized as the first-line management for reducing hyper androgenic symptoms and anxiety level among women with PCOS.

MATERIALS AND METHODS:



Quasiexperimentalpre and posttest design was used in this study. The Study was conducted in selected colleges at Chennai. A total of 824 young girls were screened based on Rotterdam criteria and recruited. Self-administered questionnaire was used to collect the baseline data. Modified Ferriman Gallwey scale, Ludwig visual score, Global acne grading scale were used to assess the hyper androgenic symptoms like hirsutism, alopecia and acne respectively. Other markers like level of anxiety using Hamilton anxiety scale, BMI and Waist circumference were also assessed. 150 young girls with PCOS were selected who met with the inclusion criteria. The participants were recruited from colleges in and around Chennai. Non probability Purposive sampling technique was used. The study was approved by Tagore institutional ethics committee. Written consent was acquired from every one of the study participants.

After recruiting the participants, they were partitioned into three groups (50 participants in every group).

Non interventional group(n=50):only education related to PCOS

Group A (n=50): cinnamon tea, exercise program and anxiety reduction counselling

Group B (n=50): exercise program and anxiety reduction counselling

Interventions:

I. Oral cinnamon tea:

1.5 grams of cinnamon powder mixed in 100ml of warm water was given during break time once in a day for 5 days in a week for 6 months. The cinnamon tea was prepared daily by the researcher and provided to the students during the break time.

II. Exercise program:

One-hour exercise program was given once a day in the evening for 5 days in a week for 6 months. Cardio exercise and strength training exercises were practiced in the alternate days.

III. Anxiety reduction Counselling:

Participants whose anxiety level was mild and moderate based on Hamilton anxiety scale were given counselling to reduce their anxiety level. Counselling was given by the researcher. The counselling sessions was planned as follows

Session I: Introduction or Orientation Phase

Session II: Psychoeducation regarding the PCOS. The researcher explained in detail about the signs and symptoms, co morbidities, self-image disturbances and anxiety related to PCOS.

Session III: Specific Concern: After 1 week Jacobson progressive Muscle Relaxation Technique (JPMR) was planned in III session

Session IV: After 1-week interval - Diaphragmatic breathing technique was given.

Session V: After a week, behavioural modification session was planned - Counselling on diet, exercise and sleep was given.

Session VI: After a week of time, feedback session was planned. The researcher obtained feedback from the participants.

Session VII: Termination phase.

Ethical consideration:

The present study was approved by Institutional Ethics Committee, Tagore Medical College and Hospital IEC NO: 32/MAR/2021 Dated: 18.03.202.

Data analysis:

The means were compared (parametric data) by two-way repeated measures analysis of variance (2-way RM ANOVA), for groups, tests and the group X test interaction comparisons. When significant difference was found, multiple comparison was done by Bonferroni ‘t’ test for between group and within group comparisons. The medians were compared (nonparametric data) by Friedman repeated measures ANOVA on ranks with Dunnett’s multiple comparison test, and Kruskal Wallis test with Dunnett’s multiple comparison test. A probability of 0.05 and less was considered statistically significant. SigmaPlot 14.5 version (Systat Software Inc., USA) was used for the statistical analysis and graph plotting.

Results:

Table 1 - Comparison of Non interventional, group A and group B of pretest, posttest 1 and posttest 2 on hyper androgenic symptoms (hirsutism, acne and alopecia) and psychological markers (level of anxiety) in young girls with poly cystic ovarian syndrome.

S.No.	Variable	Groups	Test	Median	Percentile	Statistical
-------	----------	--------	------	--------	------------	-------------



						analysis
1	Modified Ferriman Gallwey scale	Non interventional -	Pre test	3	2 – 3	H = 248.794 P < 0.001
		Group A		2	2 – 3	
		Group B		3	2 – 3	
		Non interventional -	Post-test 1	3	3 – 3	
		Group A		2	2 – 2	
		Group B		3	2 – 3	
		Non interventional -	Post-test 2	3	3 – 3	
		Group A		1	1 – 1	
		Group B		2	2 – 2	
2	Ludwig visual score	Non interventional -	Pre test	2	1 - 2	H = 222.258 P < 0.001
		Group A		2	1 - 2	
		Group B		2	2 - 2	
		Non interventional -	Post-test 1	2	1 - 2	
		Group A		1	1 - 1	
		Group B		1	1 - 2	
		Non interventional -	Post-test 2	2	2 - 2	
		Group A		0	0 - 1	
		Group B		1	1 - 1	
3	Global acne grading scale	Non interventional -	Pre test	2	2 - 3	H = 287.511 P < 0.001
		Group A		3	2 - 3	
		Group B		3	3 - 3	
		Non interventional -	Post-test 1	3	2 - 3	
		Group A		2	1 - 2	
		Group B		2	2 - 2	
		Non interventional -	Post-test 2	3	3 - 3	
		Group A		1	0 - 1	
		Group B		1	1 - 2	
4	Anxiety scale	Non interventional -	Pre test	2	2 - 2	H = 308.921 P < 0.001
		Group A		2	2 - 2	
		Group B		2	1 - 2	
		Non interventional -	Post-test 1	2	2 - 2	
		Group A		1	1 - 1	
		Group B		1	0 - 1	
		Non interventional -	Post-test 2	2	1 - 2	
		Group A		0	0 - 0	

9537



	Group B	0	0 - 0
Non –interventional group = Control; Group A = Experimental 1; Group B = Experimental 2 n = 50 each The ‘H’ and ‘P’ values are by Kruskal Wallis ANOVA on ranks with Tukey’s multiple comparison test			

Table 2 - Comparison of Non interventional, group A and group B of pretest, posttest 1 and posttest 2 on BMI in young girls with poly cystic ovarian syndrome.

S.No.	Group comparison	Test comparison	BMI	WAIST CIRCUMFERENCE
1	Non Interventional Group	Pre-test	28.79 ± 0.19	88.70 ± 0.18
	Group A		29.52 ± 0.13	89.41 ± 0.12
	Group B		29.83 ± 0.14	88.19 ± 0.37
	Non Interventional Group	Post-test 1	29.08 ± 0.19	88.96 ± 0.19
	Group A		28.87 ± 0.13	88.67 ± 0.12
	Group B		29.44 ± 0.15	87.66 ± 0.41
	Non Interventional Group	Post-test 2	29.27 ± 0.18	89.07 ± 0.18
	Group A		28.22 ± 0.12	87.94 ± 0.12
	Group B		28.98 ± 0.15	87.50 ± 0.37
2.	Significance within group A (Pre and Post-test 1)		t = 12.903 P < 0.001	t = 9.513 P < 0.001
	Significance within group A (Pre and Post-test 2)		t = 25.805 P < 0.001	t = 18.871 P < 0.001
	Significance within group A (Post-test 1 and Post-test 2)		t = 12.903 P < 0.001	t = 9.358 P < 0.001
	Significance within group B (Pre and Post-test 1)		t = 7.797 P < 0.001	t = 6.883 P < 0.001
	Significance within group B (Pre and Post-test 2)		t = 16.861 P < 0.001	t = 8.946 P < 0.001
	Significance within group B (Post-test 1 and Post-test 2)		t = 9.064 P < 0.001	t = 2.062 P = 0.120
Non –interventional group = Control; Group A = Experimental 1; Group B = Experimental 2				

9538



n = 50 each. The 'F' and 'P' values are by two-way RM ANOVA with Bonferroni 't' test.




Figure 1,2,3 and 4 shows the Comparison of Non – interventional group (Control), Group A (Experimental 1) and Group B (Experimental 2) groups of pre-test (pre), post-test 1 (Post1) and post-test 2 (Post2) on modified Ferriman Gallwey scale, Ludwig visual score, Global acne grading scale and Anxiety scale among young girls with polycystic ovarian syndrome. The 'H' and 'P' values are by Kruskal Wallis one-way ANOVA with Tukey's multiple comparison test. *Significantly different from the respective pre-test.

Figure 5: Comparison of control (Con), experimental 1 (Exp1) and experimental 2 (Exp2) groups of pre-test (pre), post-test 1 (Post1) and post-test 2 (Post2) on BMI and Waist circumference in women with polycystic ovary syndrome. The 'F' and 'P' values are by two-way RM ANOVA with Bonferroni 't' test. *Significantly different from the respective pre-test.

In Table 1, The median values of Modified Ferriman Gallwey scale on non – interventional group Pre-test, Group A Pre-test, Group B Pre-test, non – interventional group Post-test 1, Group A Post-test 1, Group B Post-test 1, non –

interventional group Post-test 2, Group A Post-test 2 and Group B Post-test 2 are 3, 2, 3, 3, 2, 3, 3, 1 and 2 respectively. Kruskal Wallis one-way ANOVA on ranks shown significance at (P < 0.001) level. In the non – interventional group from Pre-



test to Post-test 2 there was no improvement in the score, whereas Group A and Group B shown 1 score decrease. Tukey's multiple comparison test was carried out and both the experimental groups shown significant decrease. This shows that the Group A and Group B are better than the non – interventional group.

The median values of Ludwig visual score of non – interventional group Pre-test, Group A Pre-test, Group B Pre-test, non – interventional group Post-test 1, Group A Post-test 1, Group B Post-test 1, non – interventional group Post-test 2, Group A Post-test 2 and Group B Post-test 2 are 2,2,2,2,1,1,2,0 and 1 respectively. Kruskal Wallis one-way ANOVA on ranks shown significance ($P < 0.001$). In the non – interventional group from Pre-test to Post-test 2 no improvement was observed in the score, whereas Group A shown 2 scores decrease and Group B shown 1 score decrease. Tukey's multiple comparison test was carried out and both the experimental groups shown significant decrease. This shows that the Group A is superior to Group B and non – interventional group and Group B is better than non – interventional group.

The median values of Global acne grading scale of non – interventional group Pre-test, Group A Pre-test, Group B Pre-test, non – interventional group Post-test 1, Group A Post-test 1, Group B Post-test 1, non – interventional group Post-test 2, Group A Post-test 2 and Group B Post-test 2 are 2,3,3,3,2,2,3,1 and 1 respectively. Kruskal Wallis one-way ANOVA on ranks shown significance ($P < 0.001$). In the non – interventional group from Pre-test to Post-test 2 shown 1 score increase, whereas Group A and Group B shown 2 scores decrease. Tukey's multiple comparison test was carried out and both the experimental groups shown significant decrease. This shows that the Group A and Group B are better than the non – interventional group.

The median values of anxiety scale of non – interventional group Pre-test, Group A Pre-test, Group B Pre-test, non – interventional group Post-test 1, Group A Post-test 1, Group B Post-test 1, non – interventional group Post-test 2, Group A Post-test 2 and Group B Post-test 2 are 2,2,2,2,1,1,2,0 and 0 respectively. Kruskal Wallis one-way ANOVA on ranks shown significance ($P <$

0.001). In the non – interventional group from Pre-test to Post-test 2 no improvement was observed in the score, whereas Group A and Group B shown 2 scores decrease. Tukey's multiple comparison test was carried out and both the experimental groups shown significant decrease. This shows that the Group A and Group B are better than the non – interventional group.

The Non interventional group, group A and group B - BMI and Waist circumference are given in Table-2. The mean values of BMI of Non interventional Pretest, group A Pre-test, group B Pretest, Non interventional Posttest 1, group A Posttest 1, group B Posttest 1, Non interventional Posttest 2, group A Posttest 2 and group B Posttest 2 are 28.7,29.5,29.8,29.0,28.8,29.4,29.2,28.2 and 28.9 respectively. Comparison of within the Group A shown significance ($P < 0.001$). Comparison of within the Group B shown significance ($P < 0.001$). This study also revealed that in BMI, the non-interventional group from Pre-test to Post-test 2, shown 1.7% increase in BMI, while in group A and group B, 4.4% and 3.02 % reduction in BMI was observed. Hence, group A is superior to group B and non-interventional group.

The mean values of waist circumference of Non interventional Pretest, group A Pretest, group B Pretest, Non interventional Posttest 1, group A Posttest 1, group B Posttest 1, Non interventional Posttest 2, group A Posttest 2 and group B Posttest 2 are 88.7,89.4,88.1,88.9,88.6,87.6,89.0,87.9 and 87.5 respectively. Comparison of within the Group A shown significance ($P < 0.001$). Comparison of within the Group B of Pretest and posttest 1, pretest and posttest 2 shown significance ($P < 0.001$) whereas Posttest 1 and Posttest 2 doesn't shown significance ($P = 0.120$). In non-interventional group from Pre-test to Post-test 2, 0.3% increase in waist circumference was noted, while in group A and group B, 1.6% and 0.6 % reduction was observed. Hence it revealed that group A and group B are better than the Non interventional group. The Group A is superior to group B.

Discussion:

PCOS is complex which includes reproductive, metabolic, and mental health



issues. Women with PCOS have endocrine and ovarian impairment, as well as a high range of metabolic dysfunction, such as overweight and glucose intolerance. In women with PCOS, hyperinsulinemia as compensatory mechanism induces overproduction of ovarian androgens, leading to hyperandrogenism. Hyperandrogenism often exhibit symptoms like hirsutism, alopecia, acne and body weight issues. Cinnamon, a well-known spice for centuries has countless health benefits in it. The recent in-vitro and in-vivo research has discovered new healing properties of cinnamon. Various research evidences show that cinnamon supplementation improves menstrual cyclicity, reduces glucose, triglycerides, low-density lipoprotein (LDL), total cholesterol and also improved insulin sensitivity in women with Polycystic Ovarian Syndrome(PCOS).The other benefits of cinnamon, includes anti-inflammatory, antibacterial, antifungal, and antioxidant properties. Evidences shown that cinnamon can diminish insulin resistance by expanding activation of the IRS/PI-3 kinase insulin signalling pathway. The cinnamon also stimulates auto phosphorylation of the insulin receptor and inhibit protein tyrosine phosphatase I. Through these two components cinnamon make adipocytes to build the glucose take-up and glycogen synthesis. Cinnamon is also the well-known anti-oxidant which reduces the oxidative stress in the body and thus reducing the symptoms of polycystic ovarian syndrome.The present study shown that consuming cinnamon tea weekly 5 days for 6 months reduced symptoms like hirsutism, alopecia, acne problems, BMI and waist circumference in experimental groups at significant level of $P < 0.001$.

Due to the scientific advancements, the common lifestyle exercises are being replaced with modern inventions and technologies. Most of the young girls follow sedentary lifestyle like watching TV, playing video games, spending too much of time in social media, and other gadgets. Thus, the opportunity for physical exercises is limited. Lifestyle alterations such as proper diet, exercise, stress free life, a good rest and sleep are associated with less risk, least cost and do not cause any ill effects on the reproductive system of adolescent girls. Thus, it is beneficial considering

lifestyle modifications as a propitious approach for reducing the risk of PCOS. Weight loss is considered as an important treatment modality for PCOS. Weight loss improves physiological, biochemical and psychological parameters of PCOS. weight loss can restore ovulation and pregnancy rates, decreases insulin levels, diminishes hyperandrogenic symptoms, lowers testosterone levels and improves psychological considerations. This study shown that planned exercise program, which includes cardio and strength training exercises practiced in alternate days for 5 days in a week shown the favourable result in reducing PCOS symptoms.

Women with PCOS have both clinical and biochemical signs of hyperandrogenism. Evidences shown a relationship between anxiety and hyperandrogenic symptoms. It has been suggested that women with PCOS have a poor self-esteem, a more negative self-image, and have higher levels of anxiety, depression and psychological distress owing to their body image disturbance. Barry et al. shown that girls with PCOS were significantly had difficulty coping with stress, anxiety and depression than the control group (20). The planned counselling sessions and reduction in the hyperandrogenic symptoms together can help the women with PCOS in reducing their anxiety level.

This study revealed that consuming cinnamon tea, 1 hour of exercise program and anxiety reduction counselling for 6 months shows significant reduction in the Hyper androgenic symptoms and anxiety at the level of $P < 0.001$ which was in concordance with the study of Seyed Mohammad Mousavi et al (2020) illustrated that cinnamon administration significantly decreased Body Weight and Waist Circumference. Greater effects on Body Weight were observed in subjects with a baseline BMI of ≥ 30 kg/m². (25)

The present study was undertaken to observe the effectiveness of cinnamon, exercise and counselling in reducing hyperandrogenic symptoms and anxiety level of young girls with polycystic ovarian syndrome. There was a significant reduction ($P < 0.001$) in the PCOS symptoms like hirsutism, alopecia, acne, BMI, waist circumference and anxiety level in group A



with all three interventions(cinnamon tea, exercise and anxiety reduction counselling). There was also better reduction in group B with interventions like exercise program and anxiety reduction counselling. Whereas there is no reduction either in the hyperandrogenic symptoms or anxiety level among Non interventional group.

Conclusion:

As the aetiology of PCOS is not clear, poor lifestyle habits, stress, anxiety and lack of physical activity as considered to be the predisposing factors. Evidences shown that by modifying once lifestyle and consuming anti-oxidants will gradually reduce the symptoms of PCOS. The present study results support the beneficial effects of cinnamon, exercise and counselling in reducing the hyper androgenic symptoms like hirsutism, acne, alopecia, BMI, waist circumference and also the level of anxiety among young girls with Polycystic ovarian syndrome. These interventions are cost-effective, less time consuming and it is effective in decreasing the hyper androgenic symptoms, certain physiological symptoms and psychological parameters. Hence, the study recommends cinnamon, exercise and counselling for young girls with Polycystic ovarian syndrome. However, we want further investigations with larger participants to research the impact of cinnamon, exercise program and counselling in reducing the different factors like blood glucose, lipid profile and total testosterone levels among young girls with Polycystic ovarian syndrome.

Source of Funding: Self-funding

Conflicts of Interest:The authors declared no competing interests.

References:

1. Selma Feldman Witchel, Sharon E Oberfield, and Alexia S Peña (2019) Polycystic Ovary Syndrome: Pathophysiology, Presentation, and Treatment With Emphasis on Adolescent Girls. *Journal of Endocrine Society.*: 3(8): 1545–1573
2. Wendy M. Wolf, Rachel A. Wattick, Olivia N. Kinkade, Melissa D. Olfert (2018) . Geographical Prevalence of Polycystic Ovary Syndrome as Determined by Region and Race/Ethnicity. *International Journal of Environmental Research and Public Health*: 15(11): 2589
3. Ritu Deswal, Smiti Nanda, Veena S. Ghalaut, Prasanta S. Roy Amita S. Dang (2019) Cross-Sectional Study Of The Prevalence Of Polycystic Ovary Syndrome In Rural And Urban Populations. *International Journal Of Obstetrics And Gynaecology*. 10.1002
4. F J Broekmans, E A H Knauff, O Valkenburg, J S Laven, M J Eijkemans, B C J M Fauser (2006) PCOS according to the Rotterdam consensus criteria: Change in prevalence among WHO-II anovulation and association with metabolic factors. *British Journal of Obstetrics and Gynaecology*: 113(10)
5. Disha Shetty, Baskaran Chandrasekaran, Arul Watson Singh, Joseph Oliverra (2017) Exercise in polycystic ovarian syndrome: An evidence-based review. *Saudi Journal of sports medicine*: 17(3) Page : 123-128
6. Catherine Ulbricht, Erica Seamon, Regina C. Windsor (2011) An Evidence-Based Systematic Review of Cinnamon by the Natural Standard Research Collaboration. *Journal of Dietary Supplements*, 8(4):378–454
7. Kort DH, Lobo RA (2014) Preliminary evidence that cinnamon improves menstrual cyclicity in women with polycystic ovary syndrome: a randomized Non interventional group trial. *Journal of Obstetric Gynecological* 2014;211:487.
8. Mangala Gowri P, Mary Minolin T, Thenmozhi P (2017) Effectiveness of Cinnamon Tea in Reducing Weight Among Late Obese Adolescence: *Asian Journal of Pharmaceutical and Clinical Research*, Vol 10, Issue 4, 2017, 156-159
9. Nivetha.M and Susan .G.Suganya (2016) Survey of Poly Cystic Ovarian Disease (PCOD) Among The Girl Students of Bishop



- Heber College, Trichirapalli, Tamil Nadu, India. IOSR Journal of Nursing and Health Science. Volume 5, Issue 4 Ver. I PP 44-52
10. Brenda Bruner, Karen Chad, and Donna Chizen (2006) Effects of exercise and nutritional counseling in women with polycystic ovary syndrome. *Applied Physiology Nutrition and Metabolism* 31(4):384-91
 11. Neelika Tripathi, Renu Agrawal, Richa Singh. S. K. Misra (2021)_ Prevalence and associated risk factors of polycystic ovarian disease in professional college going girls of Agra City: a cross-sectional study. *International Journal of Community Medicine and Public Health* 8(4):1917
 12. Crystal C. Douglas, Barbara A. Gower, Betty E. Darnell, Fernando Ovalle, Robert A. Oster, Ricardo Azziz, (2006) Role of diet in the treatment of polycystic ovary syndrome. *Fertil Steril.* 2006 Mar; 85(3): 679–688.
 13. Julie D. Lamb, Erica B. Johnstone, Julie-Anne Rousseau, Christopher L. Jones, Lauri A. Pasch. Marcelle I. Cedars, Heather G. Huddleston (2011) Physical activity in women with polycystic ovary syndrome: prevalence, predictors, and positive health associations. *Research Reproductive endocrinology and infertility.* volume 204, issue 4, p352.e1-352.
 14. Amie Woodward, David Broom, Caroline Dalton, Mostafa Metwally. Markos Klonizakis (2020) Supervised exercise training and increased physical activity to reduce cardiovascular disease risk in women with polycystic ovary syndrome: study protocol for a randomized Non interventional groupled feasibility trial. *BMC Emergency Medicine.* volume 21, Article number: 101
 15. Anju Krishnan Nair, Bindu Nambisan, Sreekumary Radha, Jayasree Leelamma (2017)Effectiveness of lifestyle modification package among overweight and obese adolescent girls between 15-19 years with polycystic ovarian syndrome. *International Journal of Community Medicine and Public Health.* Vol 4.No 1((2)
 16. Somaya Ouda Abd Elmenim, Afaf Mohamed Mohamed Emam.(2016) Effect of Lifestyle Changes on Symptoms of Polycystic Ovarian Syndrome in Obese Girls. *IOSR Journal of Nursing and Health Science.* Volume 5, Issue 3 Ver. II ,PP 01-10
 17. Grei Shele, Jessica Genkil, Diana Speelman(2020) A Systematic Review of the Effects of Exercise on Hormones in Women with Polycystic Ovary Syndrome. *Journal of Functional Morphology and Kinesiology.* 5(2):35.
 18. Roger Hart , Martha Hickey, Stephen Franks (2004) Definitions, prevalence and symptoms of polycystic ovaries and polycystic ovary syndrome. *Best Practices in Research Clinical Obstetrics and Gynaecologyy.* 18; 671-683.
 19. Nidhi R, Padmalha V, Nagarathna R.,and Amritanshu R(2011).Prevalence of Polycystic Ovarian Syndrome in Indian adolescent. *Journal of PediatricsandAdolescent Gynecology*24(4)223-227
 20. Barry JA, Hardiman PJ, Saxby BK, Kuczmierczyk A. Testosterone and mood dysfunction in women with polycystic ovarian syndrome compared to subfertile controls. *J Psychosom Obstet Gynaecol.* 2011;32(2):104–11.
 21. Norman RJ, Davies MJ, Lord J and Moran LJ(2002) The role of lifestyle modification in polycystic ovary syndrome. *Trends in EndocrinologyMetabolism*13251–257
 22. Hajimonfarednejad, M. and Nimrouzi, M. and Heydari, M. and Zarshenas, M.M. and Raei, M.J. and Jahromi (2018),Insulin resistance improvement by cinnamon powder in polycystic ovary syndrome: A randomized double-blind placebo Non interventional groupled clinical trial, *Phytotherapy Research*,volume 32,number 2,pages 276-283.
 23. Vijayan CP, Sonia A. Prevalence of Polycystic Ovary Syndrome among students of a teaching collegiate hospital. *Health Sci.* 2013;2(1):4.
 24. Vijaya K, Bharatwaj RS. Prevalence and undetected burden of polycystic ovarian



syndrome (PCOS) among female medical undergraduate students in South India—a prospective study in Pondicherry. *GJRA—Glob J Res Analysis*. 2014;3:63–64

25. Seyed Mohammad Mousavi, Jamal Rahmani, Hamed Kord-Varkaneh, Ali Sheikhi, Bagher Larijani, Ahmad Esmailzadeh, Cinnamon supplementation positively affects obesity: A systematic review and dose-response meta-analysis of randomized controlled trials, *Clinical Nutrition*, Volume 39, Issue 1, 2020, Pages 123-133, ISSN 0261-5614,

9544

