



Dynamics of PIWI-interacting RNAs in Disease and Medicine

Arpita Parakh ^a

Email: parakha@rknec.edu

1879

Abstract

Small RNA plays important role in regulation of gene expression and decides the cell fate. They are found to regulate mRNA transcript which in turn are responsible for translating into proteins. Recent discovery of PIWI-interacting RNAs revealed many mysterious dimensions of biology as they are known to regulate “the ancient relics” called as transposons or “jumping genes”. Previously PIWI-piRNA pathway was known to be germline specific but recent studies shown their strong role in somatic cells and neurons in diverse organism. Hence, this review is the study of multifaceted role of piRNA in transposon regulation with the compilation of their recent discovery in different cell types in order to provide a strong understanding of their role in Disease diagnosis and medicinal therapeutics to research fraternity.

Keywords : PIWI-interacting RNA Transposons, neurons, Disease Diagnosis, Medicinal Therapeutics

DOI Number: 10.14704/nq.2022.20.9.NQ44218

Neuro Quantology 2022; 20(9):1879-1893

1.1 Background


P-Element induced wimpy testis (PIWI) proteins are known to interact and guide a class of noncoding small RNA called as PIWI-interacting RNAs (piRNAs). Recent discovery of millions of PIWI interacting RNAs exposed a mesmerizing and unexpected dimension of molecular biology. piRNAs are first discovered in *Drosophila Melanogaster* germ line cells and it was found that they play major role in transposon silencing, genome rearrangement, spermatogenesis and maintaining germ line integrity but many recent evidences proposed their presence in somatic cells, neurons and cancer cells which illustrated their multifaceted function like epigenetic programming, stem cell function and whole body regeneration [1-2]. The recent role of piRNA and its function is reported in neurons suggest their strong role in neurodegenerative disorders like *amyotrophic lateral sclerosis* (ALS), Alzheimer's disease (AD), Parkinson's disease (PD), Rett's Syndrome and aging [3]. Further studies on disease associated piRNA lead us to significant study of neurodegeneration and ^a RCOEM, Nagpur, India help in development of novel therapeutics to contribute improvement in

medicine, early prognosis and diagnosis as well. Recently Philip D. Zamore evidenced common biogenesis of piRNA across evolutionary broad range of animals which shows that if transposons are known as ancient relics as piRNAs also must have been associated with them and evolved differently in broad range of animal and different cell types.

1.2 Transposons “The ancient relics”

Transposons so called molecular parasites make up large percentage of genome volume in mammals and other lower organisms. These are known to be the ancient remnants accruing by time within the genome and responsible for influencing the genome by their property of mobilization. They composed 45% of the humans and one third of flies' genome and are known to be triggered by piwi-interacting RNAs (piRNAs), a class of very less known small non-coding RNAs responsible for maintaining genomic integrity, heterochromatin formation and epigenetic regulation of germ line organs in gonads of flies and other mammals.





1880

Figure 1.1: Transposable Elements and Types: A) showing Transposons use “Cut and Paste” Mechanism B)

Retrotransposons “Copy and Paste” mechanism.

Transposons were first discovered in maize by Barbara McClintok in 1940 [4-5]. Transposable elements are genetic material also called as “jumping genes” which are able to multiply within the genome by copying themselves to the other location and which leads to variation within the species as they are notorious enough to move from one genome location to another. Hence they are involved in genome rearrangements, genome size, and also lead to lethal or adaptive effects as well [6]. TE sequences are divided into those with or without a RNA transposition intermediate, insertion mechanism, features of TE sequences i.e. internal coding and non-coding sequence, terminal repeats, autonomous and non-autonomous TEs i.e. TE relying on other elements to move due to lack of certain proteins which help them to move from one place to other. Biologically, TEs are classified into two main classes i.e. retrotransposons called as Class I TEs and DNA transposons i.e. Class II TEs. The former transposon from RNA intermediate by transcription and again by reverse transcription i.e. “copy and Paste Mechanism”. It includes long terminal Repeat (LTR), Dictyostelium intermediate repeat sequence (DIRS – like elements), Penelope- like elements (PLEs), longinterspersed elements (LINEs) and Short Interspersed elements (SINEs). It ranges from 5-7 kb in drosophila which includes Copia, Gypsy, ZAM, Idefix and Tirant are infectious retrotransposon family found to infect other cells [7]. The latter can transpose via DNA intermediate. They are of four type i.e. Terminal

inverted repeats (TIR), Crypton, Helitron and Maverick. These can increase DNA volume by getting replicated with the chromosome or by gap repair mechanism. Transposon have their own machinery to replicate and can behave autonomous and in non- autonomous way. It is depicted in previous studies that transposons maniac is handled by class of small RNA called as piwi-interacting RNA which are known to silence transposons in germline cells as they are responsible for incorporating genetic variations and genome defense mechanism recruits piRNA to handle the transposons expression and maintain the integrity of the genome. In female flies the piRNA are found to silence an endogenous retrotransposon called as *gypsy* in somatic follicle cells during oogenesis [8] and in males they are found to silence a gene called as *stellate* a gene involved in male sterility [16]. Later on *flamenco* a piRNA producing locus was identified which is responsible for controlling other TE families [17]. Hence piRNAs previously called as rasiRNA (repeat associated small interfering RNAs) controls the expression of TE in germline cells to maintain the integrity as well as variation in the generations.

1.3 PIWI-interacting RNA (piRNA)

piRNAs are a class of non coding single stranded RNA with 24-36 nucleotides long which guide a mammalian class of Argonaute proteins called as PIWI proteins. Initially piRNAs were found in *Drosophila* testis and are named as rasiRNA in 2001 later a distinct class of piRNA is recognized as they were found to be associated with PIWI unlike other small RNAs. piRNAs are found to be



different according to their source of origin 1) piRNAs are known to silence transposons and are repeat specific piRNAs, they are formed from remnants of transposons and further silence active transposons. 2) piRNAs which are formed from 3'UTR regions and are mainly found in somatic cells like ovarian somatic cells of flies and spermatocytes of mice. 3) piRNAs derived from long noncoding RNA their function is still not much known [1].

1.4 Argonaute Protein (PIWI Protein)

Argonautes are known for small RNA guiding protein as they are known to guide them to target. PIWI is a subfamily of Argonaute also called as AUB and AGO3 in flies which is expressed in both germline cells and somatic cells of gonads. It has mainly PAZ and MID domain which makes PIWI protein to associate with piRNAs specifically. As, PAZ and MID domain of PIWI binds specifically with piRNA by recognising their 5'U end. Hence they play major role in piRNA specific transposon silencing.

1.5 Biogenesis: PIWI-interacting RNA and transposon silencing in germline

piRNA are class of small non-coding RNA ranging from 23-34nt in length present in all mammals including humans. These are found to bind with specific clade of Argonaut called as PIWI proteins. Many evidences suggest that piRNA biogenesis takes place in three phases in all organisms i.e. primary pathway, ping-pong mechanism and the phased primary pathway. The primary pathways produce pre-piRNA which is ready to bind with Piwi and Aub further ping-pong mechanism is initiated by primary pre-piRNA to produce secondary piRNAs and AGO3 slicer dependent trimming leads to production of secondary piRNA also termed as amplification of piRNA and increase the population of piwi bound piRNAs. This whole pathway is organized in a systematic manner to silence the transposon and require a range of proteins, transcription factor and determinants to procure piRNA biogenesis [18]. Phased primary pathway is also called as zucchini dependent pathway as, zucchini is responsible for trimming and producing piRNA transcript of specific size. There are recent evidences which present the advances in piwi- piRNA biogenesis

includes various studies like identification of *myb-A* transcription factor required for transcription of pre-pachytene piRNA in mice from piRNA cluster rich regions

1.5.1 Foundations and accreditation of piRNA precursor existence 1881

The piRNAs are characteristically distinguished from canonical RNAi pathways of miRNAs and siRNAs which include its dicer free processing and origin from single stranded precursors with the distinguished length of

23-34 nucleotides. These piRNAs are recognized by PIWI clade of ARGONAUTES containing structural features


with amino terminal domain, PAZ domain, MID domain and PIWI domain (animal specific). Other than this Argonautes are separated into three groups AGO, WAGO and PIWI clade which are associated with different small RNA classes. In flies PIWI is responsible for guiding piRNA to silence transposons. The existence and evolution of piRNA is a big mystery which needs a lot more studies. Various evidences revealed association of multiprotein complex is responsible for recognizing specific genomic landmark for transcribing piRNAs and it was found that piRNA precursors are transcribed from genomic clusters containing transposons fragments accumulated from years in the genome which in turn gives a clue of genetic memory of transposition invasion. This genetic memory lead us to a thought of adaptive immune response mechanism adapted by antisense primary piRNAs which are transcribed from clusters formed from the accumulation of transposons. The accumulation of transposon is found in various genomic locations which act as a trigger for piRNA silencing. It is predicted that moonshiner is recruited by piRNA cluster via heterochromatin protein-1 variant Rhino and moonshiner in turn recruits TBP related factor triggers piRNA transcription initiation [19]. Hence histone marks are responsible for triggering transcription in heterochromatin regions instead of any sequence motif. There are two types of piRNA cluster known till date known as unistrand and dual strand cluster according to their capacity of generating piRNAs. The main feature of unistrand cluster is formation of piRNA from only one genomic



strand. *Flamenco* and 20A are two unistrand cluster known in *Drosophila* till date. 20A cluster is found in somatic follicle cells and germ cells but flamenco is limited to follicle cells repress gypsy-like transposons. The specific features of flamenco cluster give the idea of production and foundation of piRNA as these clusters are most known and prevalent. Flamenco is located in the boundary of euchromatin and heterochromatin region colonized with fragments of gypsy family TEs in minus strand. The promoter is marked with H3K4me2 and cluster body is surrounded by H3K9me2/3 marks and transcribed with RNA polymerase II [19]. The transcript formed by RNA pol II are presented for alternative splicing

and transferred to Yb bodies via nuclear DotCOM (place for accumulation of *flam* transcript) [20]. Dual strand cluster are predominantly found in germ cells of *Drosophila* produces piRNA clusters. They are majorly located in pericentromeric and subtelomeric heterochromatin regions and boundaries of euchromatin regions. They are transcribed by RNA polymerase II. Recognizing transcription site on piRNA cluster and initializing transcription are hard-hitting process as it requires a multiprotein complex in germline cells.

1882



Figurer 1.2: piRNA transcription A) differences in transcription of unistrand and dual strand cluster former use alternative splicing and latter uses no splicing. B) the complex required for dual strand transcription of piRNAs which majorly happens in germline cells.

(adopted from Huang X et al., 2017).

It was found that Rhi and deadlock and cuff (A RDC complex) interact with each other and act as anchor between the transcription site and other

protein complex (TBP-TATA-box binding protein) which makes a platform for different proteins to get associated. Hence, Rhi-Del



platform which binds with cuff (cap binding protein) and moonshiner (a link between RDC and TBP-TATA-box binding protein) guides other protein to loci of piRNA producing cluster as, H3k9me3 decoration is found everywhere in genome and this methylation is found in areas which are highly silent heterochromatin region but they only act as recognition site in case of piRNA transcription [21]. This complex not only initiates transcription of piRNAs but also suppress the termination and stabilization by deciding the timing of RNA polymerase II binding which finally makes antisense piRNA transcript for transposon silencing. Further it was also studied that TREX is also responsible for transcription of dual strand cluster as knockout experiments lowered the rate of piRNA transcripts and was found to be co-localized with RDC in nuclear foci [22]. Finally, the transcripts formed by dual stranded clusters do not go for capping and splicing as they are bounded by cuff (which prevents them for capping) and DEADBOX helicase U2AF65 associated protein which takes them to nuclear periphery where vasa escorts them to naug for further processing.

1.5.2 Pre-piRNA processing in cytoplasm

In flies RNA piRNA transcript and requires rhino-Deadlock cutoff/moonshiner, U2AF65 associated protein (UAP56), a complex for transcription and stabilization of precursor piRNA [23-24] which are exported to cytoplasmic naug for processing and transposon silencing in germline cells of drosophila. The various export factors are associated to export piRNA from nucleus to naug followed by piRNA processing and recently it is found that MOV10L1 (mouse homologue) [25]/armi (in drosophila) is a factor which is associated with piRNA processing. MitoPLD/ zucchini is found to play a major role producing 5'end of precursor piRNA and produce specific 5'U bias for loading into PIWI protein. Zucchini a homolog of mice MitoPLD is found in flies which trims the piRNA which are generated from clusters rich in transposons [26-29]. Historically pre-piRNA formation is zucchini dependent also called as primary biogenesis of piRNA but it can be also be called as zuchhini dependent phased piRNAs as, it is also found that 3'end formation of piRNAs is also deployed by *zuc* and exhibit a signature of phased piRNAs.

Hence primary piRNA biogenesis can also be called as zuchhini dependent phased piRNA. Further piRNA maturation take place with *nibbler* which participates in exonucleolytic maturation of piRNA and *papi* [23,30] has been implicated in formation of 3'end formation which is the last step of piRNA processing as it decides the size of piRNA and then mature PIWI associated piRNAs are methylated by HEN 1 to produce 2' end of piRNA to make them single stranded stable structure and control transposons epigenetically [31-34].

The second phase which is also called as amplification of piRNA or ping-pong mechanism requires lot of determinants for its deployment. The ping pong signatures are also found in planarians [35], Koala genome [36], Hydra [37], Zebra fish and silk-worms [38]. The second phase of piRNA biogenesis is followed by transcription and stabilization of primary precursor in germline cells of *Drosophila Melanogaster* which require a protein complex called Rhino-Deadlock-Cutoff which stabilize and flag them for piRNA processing in Yb in case of follicle cells and naug in case of germline cells bodies [39]. Both the phases are interconnected as the primary piRNA are exported from the processing site of cytoplasm to nucleus once again for ping-pong amplification. The primary piRNA are trimmed very specifically with zucchini which makes the way for phased piRNA as trimming independent [40]. Zucchini interplays during primary pathway of piRNA biogenesis and Tudor interplays in secondary pathway of piRNA biogenesis i.e. during ping-pong amplification Tudor protein plays a major role. Piwi-piRNA complex interact with TUDOR family for amplification. Several TDRDs such as Spn-E, Tudor, Krimper, and tejas are found to be important for PIWI functioning [41-42]. The mature piRNA-Piwi complexes have different chance as the mode of localization of Argonaute is different. Piwi localize into nucleus and Aub localize in perinuclear cytoplasm hence there can be two forms in which piRNA can carry out silencing i) Transcriptional gene silencing, ii) post transcriptional gene silencing which is also involved in ping-pong mechanism. Researchers identified 1U bias in piRNAs strands bound with Aub derived from antisense piRNA clusters and 10A bias in sense piRNA strand derived from transposons and are bond with Ago3 and whole



process occurs in naugae a perinuclear germline structure reveals the un-spliced amplification of piRNA called as ping-pong mechanism which leads to silencing of transposons without any slicer enzyme. Ping-pong cycle leads to adaptive and sequence specific mechanism of PIWI-piRNA mediated silencing which do not require any slicer but needs TUDOR family to mediate its function and maintain its stability [43-44]. So, in germline cells piRNA biogenesis occurs in naugae and their licensing is different from somatic cells.


Ping-pong amplification occurs in germline cells of flies, the mechanism increases the piRNA population and responsible for dicer independent splicing. The production of phased

piRNA occurs by specific binding and overlaps of 1U bias of AUB associated piRNAs and 10A bias of Ago3 associated piRNAs. Many computational studies revealed the 10nt overlap between AUB-Ago3 associated sequences i.e. AUB is bound to 1U biased piRNAs and Ago3 is bound to 10A bias piRNA and their 10 nt. interaction is revealed by analyzing sequences computationally using various tools. This interaction of Aub- Ago3 bounded piRNA leads to formation of more piRNAs from the transposons transcripts. This amplification mediates from biogenesis of piRNA to its target silencing hence it plays very important role in transposon silencing via piRNAs. The primary piRNA produced from clusters are associated with AUB have the capacity to slice transposons transcript at the specific position with its slicer activity and the piRNAs which are produced have a 5' end and are in sense orientation of transposons and their loading and association with Ago3 is followed by maturation of piRNA includes trimming of their 3'end by unknown nuclease and forms a complex of Ago-piRNA which is again ready to detect and cleave more transcript of piRNA cluster to form antisense piRNA which are identical to target transcripts. In somatic cells maturation of Aub associated piRNAs also occurs via zucchini by trimming 3'end and production of phased piRNA occurs and these are loaded into PIWI to create adaptive silencing of transposons.

1.5.3 PIWI-piRNA biogenesis in soma

In somatic cells specifically in ovarian cells of drosophila melanogaster piRNA transcription takes place from the unistrand cluster and PIWI is expressed but AUB and AGO3 is not expressed. The piRNAs which are transcribed from unistrand clusters are same as that of formed in dual strand clusters of germline i.e. 23-34 nucleotide in size and 2' O methylated at 5' and 3' end. As structurally the piRNAs transcribed from unistrand clusters of somatic cells are same as that of germline dual strand cluster but, the process of transcription is entirely different. The well-known unistrand cluster is named as *flamenco* has its own promoter with H3K9me2 enables RNA polIII to transcribe clusters and with that the cluster body is decorated with H3K9me3 which shows the heterochromatin region. The promoter lacks TATA box but contains an initiator (Inr) motif and downstream promoter (DPE). The promoter region also ports a binding site of transcription factors called as Cubitus interruptus (Ci) in the upstream region -515 to -365 of transcription start site (Yamanaka S et al., 2014; Mohn F et al., 2014; Pane et al., 2011; Goriaux C et al., 2014). Co-immunoprecipitation experiments by

various authors discovered cytoplasmic localization of ARMI and YB which are mainly responsible for stabilizing piRNAs in somatic cells. Hence, the main protein involve in guiding the piRNA to sense transcript of transposons in somatic cells is PIWI protein and a putative helicase called as ARMI, a nuclease called as zucchini and a TUDOR protein called as YB is responsible for stabilization and biogenesis of piRNA. Unlike germline cell somatic cell piRNA do not undergo ping-pong mechanism for their amplification. Before piRNA assemble in Yb it was also found that the transcripts derived from different piRNA cluster loci are accumulated at one place on a single foci named as DOT Com. As the piRNA transcript derived from *flamenco* goes through alternative splicing and accumulated in a single nuclear structure of follicle cells called as Dot COM, the transcript are transferred from here to YB for further processing by different protein complexes. [48].



1.6 PIWI-piRNA in cancer

Cancer is most common and dreadful disease accounts for millions of deaths yearly, due to its late detection and reoccurrence [49] after treatment highlights the need for novel prognostic biomarker identification [50]. Recent discovery of piRNA in silencing transposons by interacting with various proteins and their domains in carrying out silencing shows its functions and mechanism in cancer. piRNA are responsible for silencing hence PIWI/piRNA complex interacts to form Transcriptional gene silencing and post transcriptional gene silencing and aberrant piRNA expression lead to many cancer types like lung cancer, breast cancer, colorectal cancer, kidney cancer, Hematological malignancy and Glioblastoma. At the time of transcription the complex interact with many proteins and transcriptional factors so, the overexpression of piRNAs lead to inhibition of apoptosis or aberrant expression like over expression of piR-021285 inhibits BC (Breast cancer) cells apoptosis [52], similarly piRNA-823 expression increases DNA methylation and inhibits tumor suppressor P16INK4 expression [57]. Many piRNAs also interact with mRNAs, long non coding RNAs and pseudogenes to mediate post transcription gene silencing and leads to suppression of cancer like piR-55490 binds to mTOR causing suppression of lung cancer cells. Similarly piR- 651 upregulation leads to tumor proliferation and apoptosis inhibition. piR-3481 and piR-52200 upregulation promotes cell proliferation by perturbing AMPK phosphorylation of ATM-AMPK-p53 pathway in lung cancer [59]. In Gastric cancer piR-651 up-regulates cell proliferation and piR-823 down

regulates proliferation of cancer cells and weakens methylation of tumor supporter genes [208, 54]. In colorectal cancer piR-1245, piR-54265, piR-025551, piR-823 promotes cell proliferation and growth by inhibiting apoptosis by regulating serine threonine phosphorylation and STAT3 phosphorylation. It is found that piR-32051, piR-39894 and piR-43607 are up-regulated and expressed during metastasis and high tumor state during kidney cancer [55-56]. piR-823 and piR-651 are also up-regulated in Hematological malignancy and promotes proliferation and cell cycle progression [57-59]. piR-52207 up- regulation is responsible for cell proliferation and tumor genesis by binding target mRNA (NUDT4, MTR) in ovarian cancer [60]. As, piRNA works in upstream of different regulatory networks shows their importance and for early cancer diagnosis and cancer detection. Aberrant PIWI expression is also seen in different cancer types like PIWIL1 up-regulated is responsible for DNA hypo methylation in lung cancer, regulate signaling pathway of gastric cancer, and used as a prognostic markers for predicting colorectal cancer [61-63], PIWIL2 is responsible in cervical cancer and renal cancer [64-65], PIWIL3 differential regulation leads to gastric cancer and multiple myeloma [66] and PIWIL4 are regulation expression leads to triple negative breast cancer [67]. Advanced detection of technologies can help to detect the piwi/piRNA expression in normal and cancer stages. PIWI/piRNA during biogenesis and function interact with different proteins but their overexpression or under expression may lead to many effects like phosphorylation in cellular signaling pathways, DNA methylation at a



specific loci and specific interaction with mRNA leads to aberrant expression of genes.

1.7 PIWI-piRNA in neuron

Evidences revealed the function of piRNA in germline and their key role in maintaining fertility [68, 39]. Many strong evidences showed the presence of Piwi/Hiwi/Mili proteins and piRNAs in somatic cells like neurons in brain and kidney in mice [69]. Lee et al., reported distribution of piRNA in brain for example Miwi expression was identified in Hippocampus region of mice by in Situ hybridization and confirmed by real time PCR. piRNA knockout studies by Lee et al. also shown reduced spine density in axons. Rajasethupathy et al. (2012) observed piRNA in neurons of Aplysia and interpreted their unique biogenesis pattern [70]. The amount of piRNA in brain is one tenth of the amount of piRNA in germline evidenced by [71-72, 59]. A neurotransmitter named as serotonin (5 Hydroxytryptamine) highly synthesized in neurons of Central Nervous System responsible for regulation of mood, appetite and sleep and its modulation alter the level of piwi/piRNA complex in Aplysia brain. CREB2 a cAMP responsive element binding protein 2 responsible for inhibitory constraint of memory also regulated by piwi/piRNA complex and hence responsible for memory loss and various memory disorders. As the complex interferes with the serotonin dependent methylation of conserved promoter sequence of CREB2 which is further responsible for the formation of synaptic plasticity and memory storage [73]. These evidences strongly depict the interference of Piwi/piRNA in neuronal development and memory storage on flies, mice and further in other higher organisms which in turn create a future enigma of involvement of piRNAs in neurodegenerative diseases like Alzheimer's, Huntington's, Parkinson's and Brain tumor to many other neuronal disorders. It was studied that brain tumors also had a high number of transposons and which gives a clue of presence of piRNA in tumors and hypothesis was confirmed by doing knockout experiment of piwi/Aub in drosophila suppresses the tumor growth.

1.8 DNA methylation and piRNA in neurodegenerative Disorders

piRNA dependent DNA methylation and Demethylation takes place in mammalian and germ line cells. Many recent studies depicted role of piRNA in DNA methylation of retrotransposons [74-75, 16]. Itou et al., identified role of piRNA/ PIWI protein in DNA methylation by silencing EGFP transgenes through induction of artificial piRNA and validating the results by bisulphite sequencing. However it is elucidated that piwi specific piRNA are mainly evolve from embryonic piRNA clusters and intergenic piRNA clusters. The latter are present in somatic cells including neuronal cells too. Recent research challenges these beliefs by designing an experimental system containing antisense EGFP transgenes which are methylated by producing artificial piRNA. In the same line DNA methylation is also a major cause of many neurodegenerative disorders like Alzheimer's, Dementia, Parkinson's, Rett Syndrome. Recent studies prove epigenetics as a major player in mediating neurodegenerative disorders. Many genes involved in maintaining neuronal plasticity and differentiation are epigenetically deregulated which gives a glimpse of involvement of small non coding RNAs including piRNAs in neurodegenerative disorders. Latest studies performed DNA methylation analysis of Alzheimer's, Dementia with Lewy bodies, Parkinson's and observed that neurodegenerative disorders shares same aberrant cpG methylation. The disruption of DNA methylation by piRNA (DQ599803) transcribed from piRNA cluster which is essential for neuronal plasticity and neurodegenerative disorders [73].

1.9 Conclusion

piRNAs are not only specific to germline but their presence in other somatic cells illustrates their involvement in many diseases not only limited to fertility or sterility but also to various memory related disorders and may be some neurodegenerative disorders. The androgens also play a strong role in regulating the proteins and determinants involved in biogenesis of piRNA like, Estrogen upregulates the transcription factor MYB which is involved in transcription of piRNA clusters and Testosterone

1886



upregulates various components of naugue/YB like TUDOR protein. Similarly piRNA are also depicted to be found in brain of mammalian tissue. Various researches show the role of piRNA in memory storage and also they take part in DNA methylation which is further responsible for causing neurodegenerative disorders. This debate demonstrated presence of piRNA in somatic cells other than germline cells. Although there is a remarkable progress in understanding the impairment of piRNA biogenesis and piwi proteins, but still many questions arise and remain elusive inline. Is there any reproductive disease in which interference of these piRNAs occurs, as knockout effects of any determinant of piRNA machinery maximally leads to sterility? How and why other RNAs escape from these piRNA machinery importantly mRNA? There are certain homologs of PIWI named as piwi like protein in humans, do they also interfere in piRNA biogenesis if not than what is there function? As Androgens and myb family components also play a key role in regulating the genes and proteins of piRNA machinery and take part in cell cycle and cell proliferation so, there are some links which warrants the role of piRNAs in cancer, and it is of great interest to examine mystery of piRNA and piwi proteins in neuronal disease and cancer would inhibit tumorigenesis. As many deep sequencing studies revealed presence of piRNA expression signatures in different carcinoma tumors but brain tumor studies are still elusive and unexplored. This review is mainly focused on identifying role of piRNAs in brain development and memory storage.

References

- [1] Gainetdinov I, Colpan C, Arif A, Cecchini K, Zamore PD. A Single Mechanism of Biogenesis, Initiated and Directed by PIWI Proteins, Explains piRNA Production in Most Animals. *Mol Cell*. 2018 Sep 6;71(5):775-790.e5. doi:10.1016/j.molcel.2018.08.007. PMID: 30193099; PMCID: PMC6130920
- [2] Ross RJ, Weiner MM, Lin H. PIWI proteins and PIWI-interacting RNAs in the soma. *Nature*. 2014;505(7483):353-359. doi:10.1038/nature12987
- [3] Wakisaka KT, Tanaka R, Hirashima T, Muraoka Y, Azuma Y, Yoshida H, Tokuda T, Asada S, Suda K, Ichiyanagi K, Ohno S, Itoh M, Yamaguchi M. Novel roles of *Drosophila* FUS and Aub responsible for piRNA biogenesis in neuronal disorders. *Brain Res*. 2019 Apr 1;1708:207-219. doi: 10.1016/j.brainres.2018.12.028. Epub 2018 Dec 19. PMID: 30578769.
- [4] Feschotte C. Transposable elements and the evolution of regulatory networks. *Nat Rev Genet*. 2008 May;9(5):397-405. doi: 10.1038/nrg2337. PMID: 18368054; PMCID: PMC2596197.
- [5] Ravindran S. Barbara McClintock and the discovery of jumping genes. *Proc Natl Acad Sci U S A*. 2012 Dec 11;109(50):20198-9. doi: 10.1073/pnas.1219372109. PMID: 23236127; PMCID: PMC3528533.
- [6] Mérel V, Boulesteix M, Fablet M, Vieira C. Transposable elements in *Drosophila*. *Mob DNA*. 2020 Jul 3;11:23. doi: 10.1186/s13100-020-00213-z. PMID: 32636946; PMCID: PMC7334843.
- [7] Marlor RL, Parkhurst SM, Corces VG. The *Drosophila melanogaster* gypsy transposable element encodes putative gene products homologous to retroviral proteins. *Mol Cell Biol*. 1986 Apr;6(4):1129-34. doi: 10.1128/mcb.6.4.1129. PMID: 3023871; PMCID: PMC367623.
- [8] Chalvet F, Teyssset L, Terzian C, Prud'homme N, Santamaria P, Bucheton A, Pélisson A. Proviral amplification of the Gypsy endogenous retrovirus of *Drosophila melanogaster* involves env-independent invasion of the female germline. *EMBO J*. 1999 May 4;18(9):2659-69. doi:10.1093/emboj/18.9.2659. PMID: 10228177; PMCID: PMC1171345.
- [9] Cimmino A, Calin GA, Fabbri M, Iorio MV, Ferracin M, Shimizu M et al. miR-15 and miR-16 induce apoptosis by targeting BCL2. *Proc Natl Acad Sci USA* 2005; 102: 13944-13949
- [10] Leucht, C, Stigloher C, Wizenmann A, Klafke R, Folchert A, and Bally-Cuif, L. (2008). MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary. *Nat. Neurosci*. 11, 641-648. doi:10.1038/nn.2115
- [11] Johnston DM, Sedkov Y, Petruk S, Riley, KM, Fujioka J., Jaynes J., Mazo A. (2011).



- Ecdysone- and NO- Mediated Gene Regulation by Competing EcR/Usp and E75A Nuclear Receptors during Drosophila Development. *Mol. Cell* 44(1): 51--61.
- [12] Gaudet P, Livstone M, Thomas P. (2011). Gene Ontology annotation inferences using phylogenetic trees. *GO Reference Genome Project. Brief. Bioinform.* 12(5): 449--462.
- [13] Pinnola A, Naumova N, Shah M, Tulin AV. (2007). Nucleosomal core histones mediate dynamic regulation of poly(ADP-ribose) polymerase 1 protein binding to chromatin and induction of its enzymatic activity. *J. Biol. Chem.* 282(44): 32511--32519.
- [14] Marlor RL, Parkhurst SM, Corces VG. The Drosophila melanogaster gypsy transposable element encodes putative gene products homologous to retroviral proteins. *Mol Cell Biol.* 1986 Apr;6(4):1129-34. doi: 10.1128/mcb.6.4.1129. PMID: 3023871; PMCID: PMC367623.
- [15] Chalvet F, Teyssset L, Terzian C, Prud'homme N, Santamaria P, Bucheton A, Pélisson A. Proviral amplification of the Gypsy endogenous retrovirus of Drosophila melanogaster involves env- independent invasion of the female germline. *EMBO J.* 1999 May 4;18(9):2659-69. doi:10.1093/emboj/18.9.2659. PMID: 10228177; PMCID: PMC1171345.
- [16] Aravin AA, Naumova NM, Tulin AV, Vagin VV, Rozovsky YM, Gvozdev VA. Double-stranded RNA- mediated silencing of genomic tandem repeats and transposable elements in the D. melanogaster germline. *Curr Biol.* 2001 Jul 10;11(13):1017-27. doi: 10.1016/s0960-9822(01)00299-8. PMID:11470406.
- [17] Zanni V, Eymery A, Coiffet M, Zytynicki M, Luyten I, Quesneville H, Vaury C, Jensen S. Distribution, evolution, and diversity of retrotransposons at the flamenco locus reflect the regulatory properties of piRNA clusters. *Proc Natl Acad Sci U S A.* 2013 Dec 3;110(49):19842-7. doi:10.1073/pnas.1313677110. Epub 2013 Nov 18. PMID: 24248389; PMCID: PMC3856796
- [18] Sato K, Siomi MC. Functional and structural insights into the piRNA factor Maelstrom. *FEBS Lett.* 2015 Jun 22;589(14):1688-93. doi: 10.1016/j.febslet. 2015.03.023. Epub 2015 Mar 30. PMID: 25836734.
- [19] Andersen PR, Tirian L, Vunjak M, Brennecke J. A heterochromatin-dependent transcription machinery drives piRNA expression. *Nature.* 2017 Sep 7;549(7670):54-59. doi: 10.1038/nature23482. Epub 2017 Aug 23. PMID: 28847004; PMCID: PMC5590728.
- [20] Théron E, Maupetit-Mehouas S, Pouchin P, Baudet L, Brasset E, Vaury C. The interplay between the Argonaute proteins Piwi and Aub within Drosophila germarium is critical for oogenesis, piRNA biogenesis and TE silencing. *Nucleic Acids Res.* 2018 Nov 2;46(19):10052-10065. doi: 10.1093/nar/gky695. PMID: 30113668; PMCID: PMC6212714.
- [21] Chen YA, Stuwe E, Luo Y, Ninova M, Le Thomas A, Rozhavskaya E, Li S, Vempati S, Laver JD, Patel DJ, Smibert CA, Lipshitz HD, Toth KF, Aravin AA. Cutoff Suppresses RNA Polymerase II Termination to Ensure Expression of piRNA Precursors. *Mol Cell.* 2016 Jul 7;63(1):97-109. doi: 10.1016/j.molcel.2016.05.010. Epub 2016 Jun 9. PMID: 27292797; PMCID: PMC4980073.
- [22] Hur JK, Chung YD. A novel model of THO/TREX loading onto target RNAs in metazoan gene expression. *BMB Rep.* 2016 Jul;49(7):355-6. doi:10.5483/bmbrep.2016.49.7.099. PMID: 27312604; PMCID: PMC5032001
- [23] Mohn F, Sienski G, Handler D, Brennecke J. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila. *Cell.* 2014 Jun 5;157(6):1364-79. doi: 10.1016/j.cell.2014.04.031. PMID: 24906153.
- [24] Yu B, Lin YA, Parhad SS, Jin Z, Ma J, Theurkauf WE, Zhang ZZ, Huang Y. Structural insights into Rhino- Deadlock complex for germline piRNA cluster specification. *EMBO Rep.* 2018 Jul;19(7):e45418. doi:10.15252/embr.201745418. Epub 2018 Jun 1. PMID: 29858487; PMCID: PMC6030695.



- [25] Vourekas A, Zheng K, Fu Q, Maragkakis M, Alexiou P, Ma J, Pillai RS, Mourelatos Z, Wang PJ. The RNA helicase MOV10L1 binds piRNA precursors to initiate piRNA processing. *Genes Dev.* 2015 Mar 15;29(6):617-29. doi: 10.1101/gad.254631.114. Epub 2015 Mar 11. PMID: 25762440; PMCID: PMC4378194.
- [26] Watanabe T, Chuma S, Yamamoto Y, Kuramochi-Miyagawa S, Totoki Y, Toyoda A, Hoki Y, Fujiyama A, Shibata T, Sado T, Noce T, Nakano T, Nakatsuji N, Lin H, Sasaki H. MITOPLD is a mitochondrial protein essential for nuage formation and piRNA biogenesis in the mouse germline. *Dev Cell.* 2011 Mar 15;20(3):364-75. doi: 10.1016/j.devcel.2011.01.005. PMID: 21397847; PMCID: PMC3062204.
- [27] Izumi N, Shoji K, Suzuki Y, Katsuma S, Tomari Y. Zucchini consensus motifs determine the mechanism of pre-piRNA production. *Nature.* 2020 Feb;578(7794):311-316. doi: 10.1038/s41586-020-1966-9. Epub 2020 Jan 29. PMID: 31996847.
- [28] Ipsaro JJ, Haase AD, Knott SR, Joshua-Tor L, Hannon GJ. The structural biochemistry of Zucchini implicates it as a nuclease in piRNA biogenesis. *Nature.* 2012 Nov 8;491(7423):279-83. doi:10.1038/nature11502. Epub 2012 Oct 14. PMID: 23064227; PMCID: PMC3493678.
- [29] Rogers AK, Situ K, Perkins EM, Toth KF. Zucchini-dependent piRNA processing is triggered by recruitment to the cytoplasmic processing machinery. *Genes Dev.* 2017 Sep 15;31(18):1858-1869. doi: 10.1101/gad.303214.117. Epub 2017 Oct 11. PMID: 29021243; PMCID: PMC5695087.
- [30] Hayashi R, Schnabl J, Handler D, Mohn F, Ameres SL, Brennecke J. Genetic and mechanistic diversity of piRNA 3'-end formation. *Nature.* 2016 Nov 24;539(7630):588-592. doi: 10.1038/nature20162. Epub 2016 Nov 16. PMID: 27851737; PMCID: PMC5164936.
- [31] Mickute M, Nainyte M, Vasiliauskaite L, Plotnikova A, Masevicius V, Klimašauskas S, Vilkaitis G. Animal Hen1 2'-O-methyltransferases as tools for 3'-terminal functionalization and labelling of single-stranded RNAs. *Nucleic Acids Res.* 2018 Sep 28;46(17):e104. doi: 10.1093/nar/gky514. PMID: 29901763; PMCID: PMC6158503.
- [32] Horwich MD, Li C, Matranga C, Vagin V, Farley G, Wang P, Zamore PD. The *Drosophila* RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Curr Biol.* 2007 Jul 17;17(14):1265-72. doi: 10.1016/j.cub.2007.06.030. Epub 2007 Jun 28. PMID: 17604629.
- [33] Wang H, Ma Z, Niu K, Xiao Y, Wu X, Pan C, Zhao Y, Wang K, Zhang Y, Liu N. Antagonistic roles of Nibbler and Hen1 in modulating piRNA 3' ends in *Drosophila*. *Development.* 2016 Feb 1;143(3):530-9. doi: 10.1242/dev.128116. Epub 2015 Dec 30. PMID: 26718004; PMCID: PMC4760310.
- [34] Saito K, Sakaguchi Y, Suzuki T, Suzuki T, Siomi H, Siomi MC. Pimet, the *Drosophila* homolog of HEN1, mediates 2'-O-methylation of Piwi- interacting RNAs at their 3' ends. *Genes Dev.* 2007 Jul 1;21(13):1603-8. doi: 10.1101/gad.1563607. PMID: 17606638; PMCID: PMC1899469.
- [35] Kim IV, Duncan EM, Ross EJ, Gorbovytska V, Nowotarski SH, Elliott SA, Sánchez Alvarado A, Kuhn CD. Planarians recruit piRNAs for mRNA turnover in adult stem cells. *Genes Dev.* 2019 Nov 1;33(21-22):1575-1590. doi: 10.1101/gad.322776.118. Epub 2019 Sep 19. PMID: 31537626; PMCID: PMC6824462.
- [36] Yu T, Koppetsch BS, Pagliarini S, Johnston S, Silverstein NJ, Luban J, Chappell K, Weng Z, Theurkauf WE. The piRNA Response to Retroviral Invasion of the Koala Genome. *Cell.* 2019 Oct 17;179(3):632-643.e12. doi: 10.1016/j.cell.2019.09.002. Epub 2019 Oct 10. PMID: 31607510; PMCID: PMC6800666.
- [37] Lim RS, Anand A, Nishimiya-Fujisawa C, Kobayashi S, Kai T. Analysis of Hydra PIWI proteins and piRNAs uncover early evolutionary origins of the piRNA pathway. *Dev Biol.* 2014 Feb 1;386(1):237-51. doi: 10.1016/j.ydbio.2013.12.007. Epub 2013 Dec 16. PMID: 24355748.
- [38] James AR, Unnikrishnan BS, Priya R, Joseph MM, Manojkumar TK, Raveendran Pillai K, Shiji R, Preethi GU, Kusumakumary P, Sreelekha TT. Computational and



- mechanistic studies on the effect of galactoxyloglucan: Imatinib nanoconjugate in imatinib resistant K562 cells. *Tumour Biol.* 2017 Mar;39(3):1010428317695946. doi: 10.1177/1010428317695946. PMID: 28345463.
- [39] Mohn F, Sienski G, Handler D, Brennecke J. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in *Drosophila*. *Cell.* 2014 Jun 5;157(6):1364-1379. doi:10.1016/j.cell.2014.04.031. PMID: 24906153.
- [40] Wang W, Han BW, Tipping C, Ge DT, Zhang Z, Weng Z, Zamore PD. Slicing and Binding by Ago3 or Aub Trigger Piwi-Bound piRNA Production by Distinct Mechanisms. *Mol Cell.* 2015 Sep 3;59(5):819-30. doi: 10.1016/j.molcel.2015.08.007. PMID: 26340424; PMCID: PMC4560842.
- [41] Nishida KM, Okada TN, Kawamura T, Mituyama T, Kawamura Y, Inagaki S, Huang H, Chen D, Kodama , Siomi H, Siomi MC. Functional involvement of Tudor and dPRMT5 in the piRNA processing pathway in *Drosophila* germlines. *EMBO J.* 2009 Dec 16;28(24):3820-31. doi: 10.1038/emboj.2009.365. PMID:19959991; PMCID: PMC2797066.
- [42] Kirino Y, Vourekas A, Sayed N, de Lima Alves F, Thomson T, Lasko P, Rappsilber J, Jongens TA, Mourelatos Z. Arginine methylation of Aubergine mediates Tudor binding and germ plasm localization. *RNA.* 2010 Jan;16(1):70-8. doi: 10.1261/rna.1869710. Epub 2009 Nov 19. PMID:19926723; PMCID: PMC2802038.
- [43] Aravin AA, Hannon GJ. Small RNA silencing pathways in germ and stem cells. *Cold Spring Harb Symp Quant Biol.* 2008;73:283-90. doi: 10.1101/sqb.2008.73.058. Epub 2009 Mar 6. PMID: 19270082.
- [44] Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. Discrete small RNA- generating loci as master regulators of transposon activity in *Drosophila*. *Cell.* 2007 Mar23;128(6):1089-103. doi: 10.1016/j.cell.2007.01.043. Epub 2007 Mar 8. PMID: 17346786.
- [45] Yamanaka S, Siomi MC, Siomi H. piRNA clusters and open chromatin structure. *Mob DNA.* 2014 Aug 1;5:22. doi: 10.1186/1759-8753-5-22. PMID: 25126116; PMCID: PMC4131230.
- [46] Pane A, Jiang P, Zhao DY, Singh M, Schüpbach T. The Cutoff protein regulates piRNA cluster expression and piRNA production in the *Drosophila* germline. *EMBO J.* 2011 Nov 16;30(22):4601-15. doi: 10.1038/emboj.2011.334. PMID: 21952049; PMCID: PMC3243597.
- [47] Goriaux C, Théron E, Brasset E, Vaury C. History of the discovery of a master locus producing piRNAs: the flamenco/COM locus in *Drosophila melanogaster*. *Front Genet.* 2014 Aug 4;5:257. doi:10.3389/fgene.2014.00257. PMID: 25136352; PMCID: PMC4120762.
- [48] Dennis C, Zanni V, Brasset E, Eymery A, Zhang L, Mteirek R, Jensen S, Rong YS, Vaury C. "Dot COM", a nuclear transit center for the primary piRNA pathway in *Drosophila*. *PLoS One.* 2013 Sep 9;8(9):e72752. doi: 10.1371/journal.pone.0072752. PMID: 24039799; PMCID: PMC3767702.
- [49] Scully OJ, Bay BH, Yip G, Yu Y. Breast cancer metastasis. *Cancer Genomics Proteomics.* 2012 Sep- Oct;9(5):311-20. PMID: 22990110.
- [50] Fan L, Strasser-Weippl K, Li JJ, St Louis J, Finkelstein DM, Yu KD, Chen WQ, Shao ZM, Goss PE. Breast cancer in China. *Lancet Oncol.* 2014 Jun;15(7):e279-89. doi: 10.1016/S1470-2045(13)70567-9. PMID: 24872111.
- [51] Fu A, Jacobs DI, Hoffman AE, Zheng T, Zhu Y. PIWI-interacting RNA 021285 is involved in breast tumorigenesis possibly by remodeling the cancer epigenome. *Carcinogenesis.* 2015 Oct;36(10):1094-102. doi: 10.1093/carcin/bgv105. Epub 2015 Jul 25. PMID: 26210741; PMCID: PMC5006152.
- [52] Li D, Luo L, Zhang W, Liu F, Luo F. A genetic algorithm-based weighted ensemble method for predicting transposon-derived piRNAs. *BMC Bioinformatics.* 2016 Aug 31;17(1):329. doi:10.1186/s12859-016-1206-3. PMID: 27578422; PMCID: PMC5006569.



- [53] Liu X, Zhou B, Xue L, Shih J, Tye K, Lin W, Qi C, Chu P, Un F, Wen W, Yen Y. Metastasis-suppressing potential of ribonucleotide reductase small subunit p53R2 in human cancer cells. *Clin Cancer Res*. 2006 Nov 1;12(21):6337-44. doi: 10.1158/1078-0432.CCR-06-0799. PMID: 17085643.
- [54] Martinez VD, Enfield KSS, Rowbotham DA, Lam WL. An atlas of gastric PIWI-interacting RNA transcriptomes and their utility for identifying signatures of gastric cancer recurrence. *Gastric Cancer*. 2016 Apr;19(2):660-665. doi: 10.1007/s10120-015-0487-y. Epub 2015 Mar 17. PMID: 25779424; PMCID: PMC4573768.
- [55] Weng W, Li H, Goel A. Piwi-interacting RNAs (piRNAs) and cancer: Emerging biological concepts and potential clinical implications. *Biochim Biophys Acta Rev Cancer*. 2019 Jan;1871(1):160-169. doi: 10.1016/j.bbcan.2018.12.005. Epub 2018 Dec 30. PMID: 30599187; PMCID: PMC6392428.
- [56] Chu H, Hui G, Yuan L, Shi D, Wang Y, Du M, Zhong D, Ma L, Tong N, Qin C, Yin C, Zhang Z, Wang M. Identification of novel piRNAs in bladder cancer. *Cancer Lett*. 2015 Jan 28;356(2 Pt B):561-7. doi: 10.1016/j.canlet.2014.10.004. Epub 2014 Oct 8. PMID: 25305452.
- [57] Yan H, Wu QL, Sun CY, Ai LS, Deng J, Zhang L, Chen L, Chu ZB, Tang B, Wang K, Wu XF, Xu J, Hu Y. piRNA-823 contributes to tumorigenesis by regulating de novo DNA methylation and angiogenesis in multiple myeloma. *Leukemia*. 2015 Jan;29(1):196-206. doi: 10.1038/leu.2014.135. Epub 2014 Apr 15. PMID: 24732595.
- [58] Ai L, Mu S, Sun C, Fan F, Yan H, Qin Y, Cui G, Wang Y, Guo T, Mei H, Wang H, Hu Y. Myeloid-derived suppressor cells endow stem-like qualities to multiple myeloma cells by inducing piRNA-823 expression and DNMT3B activation. *Mol Cancer*. 2019 Apr 13;18(1):88. doi: 10.1186/s12943-019-1011-5. PMID: 30979371; PMCID: PMC6461814.
- [59] Li B, Hong J, Hong M, Wang Y, Yu T, Zang S, Wu Q. piRNA-823 delivered by multiple myeloma-derived extracellular vesicles promoted tumorigenesis through re-educating endothelial cells in the tumor environment. *Oncogene*. 2019 Jun;38(26):5227-5238. doi: 10.1038/s41388-019-0788-4. Epub 2019 Mar 19. PMID: 30890754.
- [60] Singh G, Roy J, Rout P, Mallick B. Genome-wide profiling of the PIWI-interacting RNA-mRNA regulatory networks in epithelial ovarian cancers. *PLoS One*. 2018 Jan 10;13(1):e0190485. doi: 10.1371/journal.pone.0190485. PMID: 29320577; PMCID: PMC5761873.
- [61] Xie K, Zhang K, Kong J, Wang C, Gu Y, Liang C, Jiang T, Qin N, Liu J, Guo X, Huo R, Liu M, Ma H, Dai J, Hu Z. Cancer-testis gene PIWIL1 promotes cell proliferation, migration, and invasion in lung adenocarcinoma. *Cancer Med*. 2018 Jan;7(1):157-166. doi: 10.1002/cam4.1248. Epub 2017 Nov 23. PMID: 29168346; PMCID: PMC5774002.
- [62] Araújo T, Khayat A, Quintana L, Calcagno D, Mourão R, Modesto A, Paiva J, Lima A, Moreira F, Oliveira E, Souza M, Othman M, Liehr T, Abdelhay E, Gomes R, Santos S, Assumpção P. Piwi like RNA-mediated gene silencing 1 gene as a possible major player in gastric cancer. *World J Gastroenterol*. 2018 Dec 21;24(47):5338-5350. doi: 10.3748/wjg.v24.i47.5338. PMID: 30598579; PMCID: PMC6305533.
- [63] Sun R, Gao CL, Li DH, Li BJ, Ding YH. Expression Status of PIWIL1 as a Prognostic Marker of Colorectal Cancer. *Dis Markers*. 2017;2017:1204937. doi: 10.1155/2017/1204937. Epub 2017 May 29. PMID: 28634417; PMCID: PMC5467314
- [64] Iliev R, Stanik M, Fedorko M, Poprach A, Vychytilova-Faltejskova P, Slaba K, Svoboda M, Fabian P, Pacik D, Dolezel J, Slaby O. Decreased expression levels of PIWIL1, PIWIL2, and PIWIL4 are associated with worse survival in renal cell carcinoma patients. *Onco Targets Ther*. 2016 Jan 8;9:217-22. doi: 10.2147/OTT.S91295. PMID: 26811690; PMCID: PMC4712976.
- [65] Qu X, Liu J, Zhong X, Li X, Zhang Q. PIWIL2 promotes progression of non-small cell lung



- cancer by inducing CDK2 and Cyclin A expression. *J Transl Med.* 2015 Sep 15;13:301. doi: 10.1186/s12967-015-0666-y. PMID: 26373553; PMCID: PMC4571108.
- [66] Jiang L, Wang WJ, Li ZW, Wang XZ. Downregulation of Piwil3 suppresses cell proliferation, migration and invasion in gastric cancer. *Cancer Biomark.* 2017 Dec 6;20(4):499-509. doi: 10.3233/CBM-170324. PMID: 28869440.
- [67] Wang Z, Liu N, Shi S, Liu S, Lin H. The Role of PIWIL4, an Argonaute Family Protein, in Breast Cancer. *J Biol Chem.* 2016 May 13;291(20):10646-58. doi: 10.1074/jbc.M116.723239. Epub 2016 Mar 8. PMID:26957540; PMCID: PMC4865913.
- [68] Grivna ST, Beyret E, Wang Z, Lin H. A novel class of small RNAs in mouse spermatogenic cells. *Genes Dev.* 2006 Jul 1;20(13):1709-14. doi: 10.1101/gad.1434406. Epub 2006 Jun 9. PMID: 16766680; PMCID: PMC1522066.
- [69] Lee EJ, Banerjee S, Zhou H, Jammalamadaka A, Arcila M, Manjunath BS, Kosik KS. Identification of piRNAs in the central nervous system. *RNA.* 2011 Jun;17(6):1090-9. doi: 10.1261/rna.2565011. Epub 2011 Apr 22. PMID: 21515829; PMCID: PMC3096041.
- [70] Rajasethupathy P, Antonov I, Sheridan R, Frey S, Sander C, Tuschl T, Kandel ER. A role for neuronal piRNAs in the epigenetic control of memory-related synaptic plasticity. *Cell.* 2012 Apr 27;149(3):693-707. doi: 10.1016/j.cell.2012.02.057. PMID: 22541438; PMCID: PMC3442366.
- [71] Dharap A, Nakka VP, Vemuganti R. Altered expression of PIWI RNA in the rat brain after transient focal ischemia. *Stroke.* 2011 Apr;42(4):1105-9. doi: 10.1161/STROKEAHA.110.598391. Epub 2011 Feb 10. PMID: 21311060; PMCID: PMC3566794.
- [72] Peng JC, Lin H. Beyond transposons: the epigenetic and somatic functions of the Piwi-piRNA mechanism. *Curr Opin Cell Biol.* 2013 Apr;25(2):190-4. doi: 10.1016/j.ceb.2013.01.010. Epub 2013 Mar 4. PMID: 23465540; PMCID: PMC3651849.
- [73] Nandi S, Chandramohan D, Fioriti L, Melnick AM, Hébert JM, Mason CE, Rajasethupathy P, Kandel ER. Roles for small noncoding RNAs in silencing of retrotransposons in the mammalian brain. *Proc Natl Acad Sci U S A.* 2016 Nov 8;113(45):12697-12702. doi: 10.1073/pnas.1609287113. Epub 2016 Oct 24. PMID: 27791114; PMCID: PMC5111663.
- [74] Carmell MA, Girard A, van de Kant HJ, Bourc'his D, Bestor TH, de Rooij DG, Hannon GJ. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell.* 2007 Apr;12(4):503-14. doi: 10.1016/j.devcel.2007.03.001. Epub 2007 Mar 29. PMID: 17395546.
- [75] Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Totoki Y, Toyoda A, Ikawa M, Asada N, Kojima K, Yamaguchi Y, Ijiri TW, Hata K, Li E, Matsuda Y, Kimura T, Okabe M, Sakaki Y, Sasaki H, Nakano T. DNA methylation of retrotransposon genes is regulated by Piwi family members MILI and MIWI2 in murine fetal testes. *Genes Dev.* 2008 Apr 1;22(7):908-17. doi: 10.1101/gad.1640708. PMID:18381894; PMCID: PMC2279202.
- [76] Domingo-Prim J, Endara-Coll M, Bonath F, Jimeno S, Prados-Carvajal R, Friedländer MR, Huertas P, Visa N. EXOSC10 is required for RPA assembly and controlled DNA end resection at DNA double- strand breaks. *Nat Commun.* 2019 May 13;10(1):2135. doi: 10.1038/s41467-019-10153-9. PMID: 31086179; PMCID: PMC6513946.
- [77] Burgess RC, Misteli T, Oberdoerffer P. DNA damage, chromatin, and transcription: the trinity of aging. *Curr Opin Cell Biol.* 2012 Dec;24(6):724-30. doi: 10.1016/j.ceb.2012.07.005. Epub 2012 Aug 17. PMID: 22902297; PMCID: PMC3524355.
- [78] De Cecco L, Berardi M, Sommariva M, Cataldo A, Canevari S, Mezzanzanica D, Iorio MV, Tagliabue E, Balsari A. Increased sensitivity to chemotherapy induced by CpG-ODN treatment is mediated by microRNA modulation. *PLoS One.* 2013;8(3):e58849. doi: 10.1371/journal.pone.0058849. Epub 2013 Mar 6. PMID: 23484053; PMCID: PMC3590172.
- [79] Rodriguez F, Arkhipova IR. Multitasking of the piRNA Silencing Machinery: Targeting Transposable Elements and Foreign Genes



- in the Bdelloid Rotifer *Adineta vaga*. *Genetics*. 2016 May;203(1):255-68. doi: 10.1534/genetics.116.186734. Epub 2016 Mar 26. PMID: 27017627; PMCID: PMC4858778.
- [80] José Renato Pinto Judy Muller-Delp P. Bryant Chase . Will you still need me (Ca²⁺, TnT, and DHPR), will you still cleave me (calpain), when I'm 64? 23 December 2016 , Volume 16, Issue 2; <https://doi.org/10.1111/accel.12560>
- [81] Burtner CR, Kennedy BK. Progeria syndromes and ageing: what is the connection? *Nat Rev Mol Cell Biol*. 2010 Aug;11(8):567-78. doi: 10.1038/nrm 2944. PMID: 20651707.
- [82] Sturm Á, Perczel A, Ivics Z, Vellai T. The Piwi-piRNA pathway: road to immortality. *Aging Cell*. 2017 Oct;16(5):906-911. doi: 10.1111/accel.12630. Epub 2017 Jun 27. PMID: 28653810; PMCID: PMC5595689.
- [83] Cruickshanks N, Hamed HA, Booth L, Tavallai S, Syed J, Sajithlal GB, Grant S, Poklepovic A, Dent P. Histone deacetylase inhibitors restore toxic BH3 domain protein expression in anoikis-resistant mammary and brain cancer stem cells, thereby enhancing the response to anti-ERBB1/ERBB2 therapy. *Cancer Biol Ther*. 2013 Oct 1;14(10):982-96. doi: 10.4161/cbt.26234. Epub 2013 Aug 22. PMID: 24025251; PMCID: PMC3926895.
- [84] Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sada S, Klotzle B, Bibikova M, Fan JB, Gao Y, Deconde R, Chen M, Rajapakse I, Friend S, Ideker T, Zhang K. Genome-wide methylation profiles reveal quantitative views of human aging rates. *Mol Cell*. 2013 Jan 24;49(2):359-367. doi:10.1016/j.molcel.2012.10.016. Epub 2012 Nov 21. PMID: 23177740; PMCID: PMC3780611.
- [85] Weidner CI, Lin Q, Koch CM, Eisele L, Beier F, Ziegler P, Bauerschlag DO, Jöckel KH, Erbel R, Mühleisen TW, Zenke M, Brummendorf TH, Wagner W. Aging of blood can be tracked by DNA methylation changes at just three CpG sites. *Genome Biol*. 2014 Feb 3;15(2):R24. doi: 10.1186/gb-2014-15-2-r24. PMID: 24490752; PMCID: PMC4053864.
- [86] Senti KA, Jurczak D, Sachidanandam R, Brennecke J. piRNA-guided slicing of transposon transcripts enforces their transcriptional silencing via specifying the nuclear piRNA repertoire. *Genes Dev*. 2015 Aug 15;29(16):1747-62. doi: 10.1101/gad.267252.115. PMID: 26302790; PMCID: PMC4561483.

