



Effect of Fsh On Lipid Profile in Postmenopausal Women

Shaima A. Mhadi^{1*}, Nihad N. Hilal², Mohammed M. Abdul-Aziz³

Abstract

The World Health Organization (WHO) and Stages of Reproductive Aging Workshop (STRAW) define menopause as a permanent endpoint of the menstrual cycle for one year that occurs naturally or by induction of surgical procedure chemotherapy, or radiation. The etiology of menopause is classified into physiological and non-physiological. Pathophysiology of menopause includes the decline of ovary function in menopause, response to a loss of ovarian feedback mechanism, the decline of hypothalamus and pituitary function. Endocrine changes in menopause lead to alteration of gonadotropin secretion cycle patterns, changes in steroid and peptide hormones through monophasic patterns to increase gonadotropin, a decrease of estrogen. Aim: The aim of this study is to investigate the effect of FSH on lipid profile in postmenopausal women. Study design: A cross-section observational study. Method: The study is a cross sectional study done on 90 women, these women aged more than 50 years and had cessation of menstrual cycle for more than one year. The blood samples were taken from 90 postmenopausal women and interviews were carried out with them using a questionnaire. FSH measure, by ElectroChemiLuminescence (ECL) technology for immunoassay analysis done by Cobas e411 device. Lipid profile measure, by manual techniques done by use of spectrophotometer device. Result: We observed increase Follicle-stimulating hormone (FSH) during postmenopause has positive significant correlation with body mass index (BMI) (r 0.350, p 0.001), total cholesterol (TC) (r 0.397, p 0.001) and *Low-density lipoprotein* (LDL) (r 0.421, p 0.001) FSH also correlated positively but insignificant correlation with triglyceride (TG) (r 0.175, p 0.098) and *very low-density lipoproteins* (VLDL) (r 0.055, p 0.604). FSH has negative significant correlation with Estradiol (r -0.509, p 0.001) and Vit.D (r -0.220, p 0.037) as well as FSH correlated negative but insignificant with age (r -0.142, p 0.183) and High-density lipoprotein (HDL) (r -0.048, p 0.656). Conclusion: The current study showed significant positive correlation of FSH with TC and LDL but insignificant positive correlation with TG and insignificant negative correlation with HDL.

2769

Key Words: Postmenopause: FSH: Lituening Hormone: HDL.

DOI Number: 10.14704/nq.2022.20.6.NQ22272

NeuroQuantology 2022; 20(6):2769-2774

Introduction

Menopause is the end of a woman's menstrual cycle, which usually occurs between the ages of 45 and 55. Women who smoke and those who have chronic illnesses may have an early menopause⁽¹⁾. This is not an illness, but rather a normal biological process. Menopause and perimenopause—the transitional phase that begins 2 to 8 years before and lasts up to a year after a woman's last menstrual cycle—occur when the ovaries begin to shut down as women become older. As women

become older, their ovary function decreases. Total follicle depletion, sclerosing blood vessels, and reduced sex steroid production (estrogen) are all symptoms of this condition^(2,3). Reduced ovarian function would impaired ovary response to gonadotropin stimulation, altering the hypothalamus-hypophysis-ovary connection^(4,5). The levels of Follicle Stimulating Hormone (FSH) and Lituening Hormone (LH) rise throughout menopause.

Corresponding author: Shaima A. Mhadi

Address: ¹Medical Technologist (for Pathological Analyzes), Kirkuk General Hospital, Iraq; ²Professor of Chemical Pathology, Tikrit University College of Medicine, Iraq; ³Professor of Urology, Tikrit University College of Medicine, Iraq

*E-mail: Sheimaa.a@st.tu.edu.iq



Furthermore, decreased estrogen levels would cause endometrium to atrophy⁽⁶⁾. Menopause symptoms include vasomotor problems, somatic complaints including dyspareunia, hair loss, dry vagina, arthralgia, palpitation, and psychological issues like sleep difficulty and insomnia, mood swings, and cognitive issues⁽⁷⁾. The lipid profile in menopause is characterized by increased cholesterol, triglycerides, low-density lipoprotein (LDL), and apolipoprotein B levels, as well as reduced HDL and apolipoprotein A levels⁽⁵⁾. Menopause results in a reduction in high-density lipoprotein (HDL) concentration as well as structural alterations in HDL. Total cholesterol (TC) and low-density lipoprotein cholesterol (LDL-C) levels are generally greater in postmenopausal women than in premenopausal women⁽⁸⁾. Menopause-related lipid alterations, particularly an increase in LDL-C, have been identified as independent variables contributing to an increased risk of cardiovascular disease in the postmenopausal years⁽⁹⁾. Although massive endocrine hormone changes associated with menopause may have a role in cholesterol levels worsening, the specific underlying mechanism is unknown⁽¹⁰⁾. Obesity in postmenopausal women is caused by both inherited and environmental factors, with poor lifestyle choices contributing significantly to the rise in BMI and waist circumference⁽¹¹⁾.

The aim of this study to evaluate the levels of FSH, lipid profile and BMI in postmenopausal women and to show the influence FSH to lipid profile in postmenopausal women.

Method and Materials

Study Design

A prospective cross-sectional observational study was carried out in Kirkuk Puplic Hospital during the period from the last of December 2021 to the end of March 2022. The study included 90 postmenopausal women. These women were aged more than 50 years and had had more than one year after postmenstrual cycle cessation. The information about postmenopausal women in this study was retrieved from the women themselves. The information collected from postmenopausal women was carried out by using a questionnaire designed by the investigator. The questionnaire includes demographic characteristics such as age, weight, length, etc.

Inclusion criteria:

Women more than 50 years and cessation from menstrual cycle more than one year.

Exclusion criteria:

Women less than 50 years and women more than 50 years but have liver or kidney disease or taking one of this drugs such as:-

- hormonal therapy
- Antihyperlipidemic drugs.
- VitD therapy

Ethical Approval

Each patient was educated about the research's specifics, filled out a questionnaire, and signed a consent form to participate in the study, according to the Tikrit University College of Medicine's scientific committee. Kirkuk's directorate of health has also granted permission to visit Kirkuk general hospital and collect samples from patients.

Sample Collection and Preparation

About five milliliters of blood were collected from the antecubital vein of fasting postmenopausal women at (8.0-10.0 Am) and put in gel tubes without any anticoagulant at room temperature for 10-15 minutes and allowed to clot. The tube was then centrifuged (3000 rpm) for 15 minutes. The clear serum was pipetted into three clear dry eppendorfs and stored at -20°C until used for the various investigations. The levels of FSH, and Lipid Profile were measured.

Method

FSH measured by ElectroChemiLuminescence (ECL) technology for immunoassay analysis done by Cobas e411 device. Lipid profile measured by manual techniques done by use of spectrophotometer device.

Statistical Analysis

All patients signed an informed consent to take part in the study, and the study was approved by ethical committee of Tikrit University, College of Medicine. All data were presented as mean and standard deviation (SD). Statistical analysis was implemented with correlation Analysis. A P value of less than 0.05 was regarded significant. Analysis was performed by IBM SPSS Statistics for Windows version 23.0.

Table 4.1. Descriptive Statistics



	N	Mean	Std. Deviation
Age (year)	90	58.056	4.9477
M.D.	90	8.733	2.00
BMI	90	32.140	6.4419
CHO. (mg/dl)	90	200.469	49.8266
TG. (mg/dl)	90	181.948	45.4592
VLDL (mg/dl)	90	35.2293	8.00
HDL (mg/ dl)	90	38.993	9.4027
LDL (mg/ dl)	90	124.602	30.2774
FSH (IU/L)	90	62.4774	15.56938
Vit.D(ng/ML)	90	17.8804	4.00780

Results

The study is a cross sectional study which is conducted on 90 women. This study evaluated the levels of parameters such as FSH, lipid profile and BMI. These parameters levels are presented in table (4.1), the (mean±SD) were: age was (58.07±4.95) years, menstrual duration was (8.73±2.00) years, BMI was (32.14±6.44) kg/m², TC. Was (200.47±48.83) mg/dl, TG was (181.95±45.46)mg/dl, VLDL was (35.23±8.00)mg/dl, HDL was (38.99±9.41), LDL was (124.60±30.28) and mean FSH was (62.48±15.57) .We observed FSH during menopause correlated positively significantly with BMI(r 0.350, p 0.001), TC (r 0.397, p 0.001) and LDL (r 0.421, p 0.001) FSH also correlated positively but insignificantly correlation with TG (r 0.175, p 0.098) and VLDL (r 0.055, p 0.604). FSH correlated negatively but insignificantly with age (r -0.142, p 0.183) and HDL (r -0.048, p 0.656).

Correlation between FSH and Age

The study showed Negative correlation between FSH and Age in postmenopausal women (r: 0.055), the result was insignificant (P:>0.05), as shown in the Figure (4 - 1) .




Figure (4- 1). The correlation of FSH in (IU/L) with Age (years) in postmenopausal women.

Relationship between FSH and BMI

The study showed positive relationship between FSH and BMI in postmenopausal women (r: 0.350), the result was significant (P:<0.05), as shown in the Figure (4 - 2).




Figure (4- 2). The relationship between FSH in (IU/L) and BMI in (kg/m²) in postmenopausal women

Correlation between FSH and TC

The study showed positive correlation between FSH and total cholesterol in postmenopausal women (r: 0.397) ,the result was significant (P:<0.05), as shown in the Figure (3-4) .




Figure (3-4). The correlation of FSH in (IU/L) with Total cholesterol in (mg/dl) in postmenopausal women

Relationship between FSH and LDL

The study showed positive relationship between FSH and LDL in postmenopausal women (r: 0.421), the result was significant (P:<0.05), as shown in the Figure (4 - 5).






Figure (4 - 5). The relationship between FSH in (IU/L) and LDL in (mg/dl) in postmenopausal women

Relationship between FSH and TG

The study showed positive relationship between FSH and triglycerides in postmenopausal women (r: 0.175), the result was insignificant (P:>0.05), as shown in the Figure (4 - 6).




Figure (4 - 6). The relationship between FSH in (mIU/L) and triglycerides in (mg/dl) in postmenopausal women

Correlation between FSH and VLDL

The study showed weak positive correlation between FSH and VLDL in postmenopausal women (r: 0.055), the result was insignificant (P:>0.05), as shown in the Figure (3 - 7).




Figure (3 - 7). The correlation between FSH in (IU/L) and VLDL in (pg/ml) in postmenopausal women

Relationships between FSH and HDL

The study showed weak negative relationship between FSH and HDL in postmenopausal women (r: - 0.048), the result was insignificant (P:>0.05), as shown in the Figure (4 - 8).




Figure (4 - 8). The relationship between FSH in (IU/L) and HDL (mg/dl) in postmenopausal women

Discussion

FSH Correlations

Correlation between FSH and Age

The study showed weak Negative correlation between FSH and Age in postmenopausal women (r: 0.055), the result was insignificant because p value more than (0.05), as shown in the Figure (4 - 1). The ovaries become less sensitive to follicle-stimulating hormone after menopause, which explains the FSH negative correlation with age



(FSH). As a result, the ovaries stop generating estrogen and progesterone, leading estrogen levels to decrease and FSH levels to rise in the body^(12,13). Studies of younger and older postmenopausal women reveal that aging has effects on the hypothalamus and pituitary that are independent of steroid feedback loss. Between the ages of 50 and 75, LH and FSH levels drop by 30–40% after menopause. The complicated effects of aging on GnRH secretion explain these gonadotropin alterations, with a 22 percent decrease in GnRH pulse frequency compensated by a 14 percent increase in total GnRH secretion above that attributable to ovarian function loss alone. There are also age-related effects in the pituitary, with older postmenopausal women having 30% lower LH and FSH responses to GnRH than younger postmenopausal women⁽¹⁴⁾.

Estrogen negative feedback at the hypothalamic level remains unchanged in older compared with younger postmenopausal women; low-dose estrogen administration is associated with a significant decrease in circulating levels of LH, FSH and a parallel decrease in the overall amount of GnRH, with no effect on pulse frequency. Addition of progesterone decreased pulse frequency in postmenopausal women with a concomitant decrease in overall amount of GnRH, neither of which effects differed with aging⁽¹⁴⁾. The effect of estrogen negative feedback on the LH response to GnRH is not affected by aging although the FSH response to GnRH is attenuated with aging. As well as Serviente, C., et al,⁽¹⁵⁾ showed no significant correlation between FSH and Age.

The current study showed positive correlation between FSH and BMI in postmenopausal women (r: 0.350), the result was significant because p value less than (0.05), as shown in the Figure (4 - 2). FSH, particularly during menopause, can influence total body weight and fat mass. FSH and FSHR have not been studied for their potential significance in fat formation and redistribution in people as they age. We hypothesized that FSHR might be expressed in human adipocytes (fat cells) and that high circulating FSH levels would affect adipocyte activities throughout aging because of the documented association between high circulating FSH levels and increasing female body weight. As well as the Huang, W.Y., et al⁽¹⁶⁾, observed similar result positive correlation of FSH with BMI in postmenopausal women. On the other hand, the current study disagreed with study done by Serviente, C., et al⁽¹⁵⁾.

In this study FSH showed significant positive correlation with TC (r: 0.397 p: 0.001) and LDL (r: 0.421 p: 0.001) in postmenopausal women, as shown in the Figures (3 - 4), (4 - 5) respectively. This study showed positive correlation of FSH with TG (r: 0.175, p: 0.098) and VLDL (r: 0.055, p: 0.604) and negative correlation with HDL (R: -0.048, P: 0.656) but these correlations insignificant because p value more than (0.05). In a cohort of 400 postmenopausal Chinese women there was similar relationship between serum FSH, TC and LDL-C. done by Song et al⁽¹⁷⁾. Song et al. found a similar association between serum FSH, TC, and LDL-C in a group of 400 postmenopausal Chinese women⁽¹⁷⁾. Song et al. also demonstrated that evidence from the animal literature described the molecular processes by which FSH might affect lipid levels directly. Lower LDL receptor expression produced by FSH was linked to reduced LDLR expression, resulting in a rise in TC and LDL levels. Human liver tissue has FSH receptors, and when exposed to FSH, the expression of the low-density lipoprotein receptor (LDLR) is lowered.⁽¹⁷⁾

FSH has also been shown to stimulate lipid biosynthesis in adipose tissue, Cui H., et al⁽¹⁸⁾ and lipid droplet formation in human adipose tissue, Liu X., et al⁽¹⁹⁾ again demonstrating that FSH may have direct influence on lipid metabolism and specifically on TC and LDL-C. The study done by Guo et al.⁽²⁰⁾ By screening molecules related to cholesterol metabolism based on mRNA and protein expression variations, they found FSH regulated cholesterol metabolism. By measuring the rate of hepatic de novo cholesterol biosynthesis and the expression of its rate-limiting enzyme, 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase (HMGCR), they demonstrated the direct influence of FSH on cholesterol biosynthesis. The work by Song et al reports that FSH may react with its receptors in hepatocytes and decreases LDLR levels, which subsequently attenuates the endocytosis of LDL-C, resulting in an increase circulating LDL-C level⁽¹⁷⁾.

Similarly, Serviente et al. found that participants with greater serum FSH levels had higher levels of both TC and LDL-C in a study of 588 postmenopausal women⁽¹⁵⁾. In this investigation, FSH had a positive association with TC, LDL, TG, and VLDL and a negative correlation with HDL, but no significant link with TG, VLDL, or HDL. Menopause causes a rise in FSH, which can cause dyslipidemia by raising TC, TG, and LDL while lowering HDL. This Dyslipidemia is a well-known independent risk factor for CVD and cerebral



ischemic stroke in postmenopausal women, and it's also linked to other metabolic disorders including diabetes, obesity, and fatty liver. A studies done by, Zhang, W., et al ⁽²¹⁾ agreed with this study and found that high FSH lead to increase of TC, LDL and TG and decrease HDL in postmenopausal women.

Conclusions

The study concluded that:

1. The study showed insignificant negative correlation between FSH and age in postmenopausal women.
2. The study demonstrated significant positive relationship between FSH and BMI in postmenopausal women.
3. There is significant positive correlation between FSH and TC and LDL, but insignificant positive correlation with TG and insignificant negative correlation with HDL in postmenopausal women.

References

- Saljoughian M. "Menopause: changes and challenges." *US Pharm* 1 (2018); 13-16.
- Washeel KhG, Sarhat ER, Jabir TH. Assessment of melatonin and oxidant-antioxidant markers in infertile men in Thi-Qar Province. *Indian Journal of Forensic Medicine & Toxicology*. 2019;13(4) ;1495- 1499.
- Entedhar Rifaat Sarhat, Siham Ajmee Wadee, Ban Ismael Sedeeq, Thuraia Rifaat Sarhat. *Biochemical and Histological Evaluation of Indomethacin-induced Hepatotoxicity in Rats*. 2019; 12(109) (Part I):23-35.
- Entedhar Rifaat Sarhat, Siham A. Wadi, B.I. Sedeeq, Th.R. Sarhat and N.A. Jasim. Study of histopathological and biochemical effect of Punica granatum L. extract on streptozotocin -induced diabetes in rabbits. *Iraqi Journal of Veterinary Sciences*, 2019. 33(1): 189-194.
- Ariadi, A., Jamsari, J., Yanwirasti, Y., Siregar, M.F.G., & Yusrawati, Y. Correlation between Estrogen Levels with Lipid Profile in Menopause Women in West Sumatera. *Open access Macedonian journal of medical sciences*. 2019; 7(13): 2084.
- Siregar MFG. Perimenopausal and post-menopausal complaints in paramedics assessed by menopause rating scale in Indonesia. *IOSR-JDMS*. 2014; 13(12): 38-42.
- Entedhar Rifaat Sarhat, Siham A. Wadi, Ayhan R. Mahmood. Effect of Ethanolic Extraction of Moringa oleifera on Paraoxonase and Arylesterase enzyme activity in High Fat Diet-induced Obesity in Rats. *Research J. Pharm. and Tech*. 2018; 11(10): 4601- 4604.
- Mešalić, L., Tupković, E., Kendić, S., & Balić, D. Correlation between hormonal and lipid status in women in menopause. *Bosnian journal of basic medical sciences*. 2008; 8(2): 188.
- Entedhar R. Sarhata*, Moayad M. Al Anzy, Takea Shaker Ahmed. Study of oxidant-antioxidant status in cerebrospinal fluid of children with meningitis. *Eurasian Chemical Communications*, 2022, 4(9), 863-869.
- Sarhat ER, Abid IM, Kamel NA, Sarhat TR, Abass KS. Changes of serum Interleukin and Chemerin levels in patients with Polycystic Ovary syndrome. *J Adv Pharm Educ Res*. 2021; 11(4): 11-4. <https://doi.org/10.51847/XP8rpqX3jx>
- Azizi F, Ainy E. Coronary heart disease risk factors and menopause: A study in 1980 Tehranian women, the Tehran Lipid and Glucose Study. *Climacteric*. 2003; 6(4): 330–336.
- Casper RF. Clinical manifestations and diagnosis of menopause. In: Barbieri RL, Crowley WF, eds. *UpToDate*. Updated March 20, 2020. Accessed April 13, 2021.
- Mohammed, E.R. Sarhat, M.A. Hamied, T.R. Sarhat, Assessment of salivary Interleukin (IL)-6, IL-10, Oxidative Stress, Antioxidant Status, pH, and Flow Rate in Dental Caries Experience patients in Tikrit Province. *Sys. Rev. Pharm.*, 2021, 12, 55-59.
- Hall, J.E. Endocrinology of the menopause. *Endocrinology and Metabolism Clinics*. 2015; 44(3): 485-496.
- Serviente, C., Tuomainen, T.P., Virtanen, J., Witkowski, S., Niskanen, L., & Bertone-Johnson, E. . Follicle Stimulating Hormone is Associated with Lipids in Postmenopausal Women. *Menopause (New York, NY)*, 2019; 26(5): 540.
- Huang, W.Y., Chen, D.R., Kor, C.T., Chen, T.Y., Lin, P.T., Tseng, J.T.C., & Wu, H.M. Relationships between Follicle-Stimulating Hormone and Adiponectin in Postmenopausal Women. *Metabolites*. (2020); 10(10): 420.
- Song, Y., Wang, E.S., Xing, L.L., Shi, S., Qu, F., Zhang, D., & Huang, H.F. (2016). Follicle-stimulating hormone induces postmenopausal dyslipidemia through inhibiting hepatic cholesterol metabolism. *The Journal of Clinical Endocrinology*. 2016; 101(1): 254-263.
- Cui H, Zhao G, Liu R, Zheng M, Chen J, Wen J. FSH stimulates lipid biosynthesis in chicken adipose tissue by upregulating the expression of its receptor FSHR. *J Lipid Res*. 2012; 53(5): 909– 917.
- Liu X, Chan HC, Ding G, et al. FSH regulates fat accumulation and redistribution in aging through the Gαi/Ca²⁺ /CREB pathway. *Aging Cell*. 2015; 14(3): 409–420.
- Guo Y, Zhao M, Bo T, Ma S, Yuan Z, Chen W, He Z, Hou X, Liu J, Zhang Z, et al. Blocking FSH inhibits hepatic cholesterol biosynthesis and reduces serum cholesterol. *Cell Research* 2019; 29: 151– 166.
- Zhang, W., Chen, Y., Zhang, X., Chen, Y., Han, B., Li, Q., & Lu, Y. Follicle-stimulating Hormone and Lipid Profile in Women Older than 55. (2021).
- Sarhat ER, Wadi SA, Mahmood AR, Sarhat TR. Measurement of the Levels of Salivary Lipocalin-2 and C - reactive protein in Women with Polycystic Ovarian Syndrome. *Tikrit J Dent Sci*. 2019; 7(1): 31-5.

