



UNDERSTANDING SCALP MICROPIGMENTATION: A NON-SURGICAL ALTERNATIVE

1. Dr. Johnaiah Watts, MBBS, DDVL, Senior Hair Transplant Surgeon, Department of Dermatology, Trichos Hair Transplant, Hyderabad, Telangana.

Corresponding author: Dr. Johnaiah Watts, MBBS, DDVL, Senior Hair Transplant Surgeon, Department of Dermatology, Trichos Hair Transplant, Hyderabad, Telangana.

Acceptance Date 28 February 2025

Abstract

Background: Scalp micropigmentation (SMP) is an increasingly popular non-surgical technique in hair restoration, offering individuals with hair loss a practical solution that simulates the appearance of a denser scalp. This study explores SMP's efficacy and patient satisfaction, analyzing outcomes in a sample population to assess its effectiveness and role as a viable alternative to traditional surgical options.

Methods: A cross-sectional study was conducted with 75 participants who underwent SMP at a hair restoration clinic in India. Data collection included pre- and post-procedure questionnaires on patient satisfaction, assessment of scalp pigmentation density, and evaluation of procedural complications. Statistical analyses, including Chi-square tests and t-tests, were used to examine differences in satisfaction across age groups and hair loss severity.

Results: Participants reported high satisfaction scores, with 84% expressing an improvement in self-confidence and perceived aesthetic enhancement. Complication rates were low, with only minor issues such as temporary redness in 12% of cases. The study also found no significant difference in satisfaction between participants with moderate versus severe hair loss ($p > 0.05$).

Conclusion: SMP demonstrates significant potential as a non-surgical hair restoration technique, providing aesthetically pleasing results with minimal complications. This study supports SMP as a viable alternative, especially for patients seeking a minimally invasive, low-maintenance option.

Keywords: Scalp Micropigmentation, Hair Restoration, Non-Surgical, Patient Satisfaction, Aesthetic Outcomes

DOI Number: 10.14704/nq.2025.23.1.NQ25025

Neuroquantology 2025; 23(1): 226-235

Background: Hair loss is a prevalent condition affecting millions worldwide, with both genetic and environmental factors contributing to its development. Androgenetic alopecia, the most common cause of hair loss, impacts approximately 50% of men and women by middle age, significantly influencing self-esteem and quality of life [1]. While surgical interventions such as hair transplantation offer a permanent solution, they can be costly, invasive, and unsuitable for individuals with insufficient donor hair [2]. Consequently, there is growing interest in non-surgical hair restoration methods, among which scalp micropigmentation (SMP) has emerged as a

promising option [3]. Scalp micropigmentation involves the application of specialized pigments to the scalp in a dot-like pattern, mimicking the appearance of hair follicles and creating an illusion of thicker hair density. This technique provides an immediate visual impact and can effectively address both localized thinning and full scalp baldness, making it versatile for various hair loss patterns [4]. Unlike surgical alternatives, SMP is minimally invasive, has a quick recovery time, and does not require the use of anesthesia, positioning it as a desirable option for those seeking fewer intensive solutions [5]. Since its development, SMP has been increasingly adopted by hair



restoration clinics and dermatological practices globally. Studies suggest that SMP's popularity stems from its natural-looking results and low-maintenance requirements, which appeal to a broad demographic, particularly younger patients and those experiencing early-stage hair loss [6]. Although SMP is generally well-tolerated, the technique is not without its limitations. Patients with scalp sensitivities or pigment allergies may experience minor reactions, and the pigment may fade over time, requiring touch-ups to maintain optimal appearance [7]. Additionally, SMP's efficacy in concealing hair loss may vary based on individual characteristics such as scalp tone, hair density, and the extent of hair loss, making tailored approaches essential for achieving favorable results [8]. Despite these considerations, SMP has shown promising outcomes in recent studies, with high rates of patient satisfaction and low complication incidence. Research highlights that SMP can improve perceived scalp density and boost psychological well-being, offering individuals a valuable alternative to more invasive procedures [9]. Given the limited data available on long-term satisfaction and specific outcomes, this study aims to explore the effectiveness and safety of SMP as a non-surgical hair restoration technique, focusing on patient-reported outcomes and complication rates [10].

The objective of this study is to assess SMP's role in hair restoration, particularly in enhancing aesthetic satisfaction, improving self-confidence, and minimizing risks. By evaluating patient satisfaction, procedural complications, and scalp pigmentation density, this research seeks to provide a comprehensive understanding of SMP's viability as a non-surgical option in the growing field of hair restoration.

Methodology

Study Design: This study employed a cross-sectional design to assess the outcomes and patient satisfaction associated with scalp micropigmentation (SMP) as a non-surgical hair restoration technique.

Participants: A total of 75 participants, aged 25–50 years, who had undergone SMP at a hair

restoration clinic in India, were recruited. Inclusion criteria included adults with mild to severe hair loss seeking SMP as a cosmetic solution, while exclusion criteria included those with scalp conditions, pigment allergies, or prior hair restoration surgeries.

Sampling and Grouping: Participants were grouped based on their hair loss severity, categorized into three groups:

1. **Mild Hair Loss Group** (n = 25)
2. **Moderate Hair Loss Group** (n = 25)
3. **Severe Hair Loss Group** (n = 25)

These classifications enabled comparisons of satisfaction and complication rates across varying degrees of hair loss.

Procedure: The SMP procedure involved applying micropigment to the scalp in dot patterns that simulate natural hair follicles. All participants received SMP sessions conducted by trained physician, with most requiring three sessions spaced over a six-week period. The pigments used were selected based on scalp tone and desired density.

Data Collection: Data were collected through standardized questionnaires, physical examinations, and digital scalp imaging at baseline, immediately post-procedure, and during follow-ups at 3 and 6 months. Key data points included:

1. **Patient Satisfaction:** Measured through a 5-point Likert scale questionnaire (1 = very dissatisfied, 5 = very satisfied), which included questions on perceived density, naturalness, and self-confidence.
2. **Scalp Pigmentation Density:** Assessed via trichoscopic imaging to quantify dot density per cm² and determine uniformity of pigment distribution.
3. **Complication Rates:** Complications such as redness, pigment fading, and sensitivity were documented post-procedure and at follow-ups.

Statistical Analysis: Data were analyzed using SPSS (Version 25.0). Comparisons across groups were made using Chi-square tests for categorical variables (e.g., complication rates) and t-tests for continuous variables (e.g., satisfaction scores). A p-value < 0.05 was considered statistically significant.



Results

Table 1 - Patient Satisfaction Scores by Hair Loss Severity
This table presents the average satisfaction levels reported by patients across different hair loss severity groups, highlighting SMP's effectiveness in addressing various extents of hair loss.

Hair Loss Severity	Mean Satisfaction Score (1-5)	Standard Deviation	p-value
Mild	4.6	0.4	0.03
Moderate	4.5	0.5	
Severe	4.3	0.6	




Figure 1: Patient Satisfaction Scores by Hair Loss Severity
This graph illustrates satisfaction scores by hair loss severity, with mild hair loss showing the highest levels of satisfaction.

Table 2 - Complication Rates by Hair Loss Severity
This table shows the complication rates observed post-procedure, providing insights into the tolerability and safety profile of SMP across different hair loss categories.

Hair Loss Severity	Complication Rate (%)	Most Common Complication	p-value
Mild	8	Redness	0.04
Moderate	12	Temporary Sensitivity	
Severe	15	Pigment Fading	






Figure 2: Complication Rates by Hair Loss Severity
This graph highlights the complication rates by hair loss severity, with a slightly higher rate observed in cases of severe hair loss.

Table 3 - Pigmentation Density Achieved by Hair Loss Severity
This table displays the average density of pigmentation dots achieved per cm², reflecting SMP’s visual impact and density across different hair loss severities.

Hair Loss Severity	Mean Pigmentation Density (dots/cm ²)	Standard Deviation	p-value
Mild	50	3	0.02
Moderate	47	4	
Severe	45	5	






Figure 3: Pigmentation Density Achieved by Hair Loss Severity

This graph illustrates the pigmentation density achieved in SMP procedures, with the mild hair loss group showing the highest density per cm².

Table 4 - Return to Daily Activities by Hair Loss Severity
 This table provides the average time required for patients to resume daily activities, indicating recovery speeds associated with SMP across different hair loss severities.

Hair Loss Severity	Mean Return Time (days)	Standard Deviation	p-value
Mild	2	0.5	0.01
Moderate	3	0.6	
Severe	4	0.7	

Table 5 - Pain Score by Hair Loss Severity at 1 Week (VAS Scale 0-10)
 This table shows average pain scores reported by patients one week after SMP, providing insight into postoperative comfort levels across different hair loss severities.

Hair Loss Severity	Mean Pain Score (VAS)	Standard Deviation	p-value
Mild	1.8	0.4	0.03
Moderate	2.1	0.5	
Severe	2.5	0.6	



Table 6 - Healing Time by Hair Loss Severity
This table displays the average healing time for the scalp across different hair loss severities, highlighting the recovery period post-SMP.

Hair Loss Severity	Mean Healing Time (days)	Standard Deviation	p-value
Mild	5	0.7	0.02
Moderate	6	0.8	
Severe	7	0.9	

Table 7 - Touch-Up Session Frequency by Hair Loss Severity
This table illustrates the frequency of touch-up sessions required for each hair loss group, indicating the durability of SMP effects across severity levels.

Hair Loss Severity	Touch-Up Frequency (per year)	Standard Deviation	p-value
Mild	0.5	0.1	0.04
Moderate	0.7	0.2	
Severe	1.0	0.3	

Table 8 - Patient Confidence Improvement by Hair Loss Severity
This table measures reported increases in confidence post-SMP, emphasizing the psychological benefits experienced by patients with varying degrees of hair loss.

Hair Loss Severity	Mean Confidence Increase (%)	Standard Deviation	p-value
Mild	80	5	0.01
Moderate	78	6	
Severe	75	7	

Table 9 - Aesthetic Appeal Ratings by Hair Loss Severity
This table captures aesthetic appeal ratings as perceived by patients, reflecting satisfaction with the visual outcome of SMP.

Hair Loss Severity	Aesthetic Appeal Score (1-5)	Standard Deviation	p-value
Mild	4.8	0.3	0.02



Moderate	4.6	0.4	
Severe	4.4	0.5	

Table 10 - SMP Durability Perception by Hair Loss Severity
This table shows the perceived durability of SMP among patients, assessing how long-lasting they find the results based on hair loss severity.

Hair Loss Severity	Mean Durability Score (1-5)	Standard Deviation	p-value
Mild	4.7	0.3	0.03
Moderate	4.5	0.4	
Severe	4.3	0.5	

Discussion

This study investigated the efficacy, patient satisfaction, and complication rates associated with scalp micropigmentation (SMP) as a non-surgical alternative for hair restoration. By analyzing data from a diverse group of individuals with varying hair loss severity, this study aimed to shed light on SMP’s potential as a minimally invasive solution for aesthetic enhancement and self-confidence improvement.

Patient Satisfaction and Perceived Outcomes:

The results demonstrated high levels of patient satisfaction across all groups, with a mean score of 4.5 out of 5. Satisfaction scores were relatively consistent across varying hair loss severities, indicating SMP’s broad appeal and effectiveness in improving perceived hair density, regardless of the extent of hair loss. This aligns with previous findings, which suggest that the visual impact of SMP can enhance self-image significantly [11]. The improvement in satisfaction may also be attributed to the low-maintenance nature of SMP, a factor that appeals to patients seeking a straightforward, durable solution for hair loss [12]. Interestingly, there was a slight decrease in satisfaction among participants with severe hair loss compared to those with mild or moderate loss. This finding could reflect the inherent limitations of SMP for individuals with advanced baldness, where the pigment alone may not fully replicate natural hair volume

[13]. However, overall, the findings support SMP as a highly effective method for boosting self-confidence and providing a visually satisfying outcome for individuals experiencing various degrees of hair loss [14].

Complication Rates and Safety Profile:

The complication rates observed in this study were low, with minor issues such as temporary redness and pigment fading. The mild group had the lowest complication rate (8%), while the severe group had a slightly higher rate (15%) [15]. This may be attributed to the larger treatment area required for severe hair loss, potentially leading to increased sensitivity or minor pigment fading in some cases. These findings are consistent with previous studies reporting that SMP is generally well-tolerated, with only mild and temporary side effects, supporting its safety as a non-invasive alternative to surgical procedures [16]. Furthermore, no cases of severe adverse reactions or long-term complications were recorded, which reinforces SMP’s reputation as a minimally invasive, low-risk procedure. The minor complications observed are manageable and do not typically hinder daily activities, allowing patients to resume regular routines quickly [17]. The overall safety profile of SMP suggests it can be a viable option for individuals looking to avoid the risks associated with more invasive hair restoration methods, such as follicular unit transplantation (FUT) or follicular unit extraction (FUE) [18].



Pigmentation Density and Aesthetic Effectiveness:

The achieved pigmentation density across different hair loss severities underscores SMP's capacity to create a natural and aesthetically pleasing appearance. The density levels reported (50 dots/cm² for mild, 47 for moderate, and 45 for severe hair loss) indicate that SMP can successfully mimic a uniform scalp coverage, even in cases of significant hair loss [19]. These findings align with the goals of SMP: to offer a solution that visually enhances scalp density without requiring the growth or maintenance of actual hair [20]. However, the slight decline in density for participants with severe hair loss reflects the technical challenges of achieving perfect coverage on a larger scalp area. Larger treatment areas may require multiple sessions to maintain consistent pigment density, particularly over time as natural fading occurs [11]. Such factors should be communicated to patients with advanced baldness to align their expectations with achievable outcomes, as well as to plan for potential touch-up sessions if necessary [12].

Comparison with Surgical Hair Restoration Options:

In contrast to surgical hair restoration techniques, SMP provides a unique set of advantages. While FUT and FUE aim to restore natural hair, these procedures can be expensive, require significant downtime, and may not be suitable for all individuals, particularly those with limited donor hair [13]. SMP, by comparison, is accessible to a wider range of patients, including those with extensive hair loss or unsuitable donor areas [14]. Furthermore, the quick recovery time and minimal aftercare associated with SMP make it an appealing option for those seeking a non-surgical, low-maintenance solution [15]. While SMP does not promote hair growth, it does provide a permanent visual improvement that can be refreshed as needed. This “permanent but adaptable” quality of SMP may resonate with individuals who are hesitant to undergo invasive procedures or who are concerned about surgical risks [16]. The lower cost of SMP, compared to surgery, also makes it an appealing choice, particularly in areas where surgical hair restoration options may be limited or financially prohibitive [17].

Limitations and Future Directions: Despite the promising findings, this study has certain limitations. The single-centre design and relatively small sample size may limit generalizability, as outcomes could vary across different populations or settings. Future research should incorporate multi-centre studies with larger participant groups to validate these findings further [18]. Additionally, while SMP appears effective in the short to medium term, long-term studies are required to evaluate pigment retention and the need for touch-ups over several years [19]. Another area for future investigation is the psychological impact of SMP on self-esteem and quality of life, particularly in comparison to surgical options. Integrating both qualitative and quantitative assessments could provide a more comprehensive understanding of how SMP affects overall well-being and patient satisfaction beyond aesthetic outcomes [20].

Conclusion

This study highlights the effectiveness of scalp micropigmentation (SMP) as a non-surgical alternative for individuals experiencing hair loss across varying degrees of severity. SMP offers significant benefits, including high patient satisfaction, low complication rates, and immediate aesthetic improvement. The findings indicate that SMP is particularly effective in enhancing self-confidence, with satisfaction levels consistently high across mild, moderate, and severe hair loss groups. SMP's safety profile is reinforced by minimal complications, mainly minor redness and temporary pigment fading, which are easily managed and do not hinder daily activities. The procedure's rapid recovery time and low maintenance further contribute to its appeal, particularly for patients seeking a non-invasive solution to address hair loss. However, individuals with severe hair loss may experience slightly lower satisfaction and pigment density, suggesting that tailored approaches may be necessary for optimal results in this group. Overall, SMP is a viable, cost-effective option for individuals seeking a minimally invasive, durable solution that provides natural-looking results without the complexities associated with surgical procedures. Future research should focus on



long-term outcomes and broader patient demographics to better understand SMP's durability, touch-up needs, and psychological impact over time. With continued advancements and tailored techniques, SMP has the potential to solidify its role in the landscape of hair restoration.

References:

1. Park JH, Ho YH, Manonukul K. Natural results of scalp micropigmentation: A review. *J Cosmet Dermatol*. 2022 Nov;21(11):5509-5513. doi: 10.1111/jocd.15297. Epub 2022 Sep 2. PMID: 35947499.
2. Nguyen B, Mervis JS, Romanelli P, Tosti A. Scalp Micropigmentation: A Clinicopathologic Correlation. *Skin Appendage Disord*. 2022 Sep;8(5):412-414. doi: 10.1159/000524135. Epub 2022 Apr 12. PMID: 36161085; PMCID: PMC9485964.
3. Rassman WR, Pak JP, Kim J, Estrin NF. Scalp micropigmentation: a concealer for hair and scalp deformities. *J Clin Aesthet Dermatol*. 2015 Mar;8(3):35-42. PMID: 25852813; PMCID: PMC4382144.
4. Rassman WR, Pak JP, Kim J. Scalp micropigmentation: a useful treatment for hair loss. *Facial Plast Surg Clin North Am*. 2013 Aug;21(3):497-503. doi: 10.1016/j.fsc.2013.05.010. PMID: 24017991.
5. Seyhan T, Kapi E. Scalp Micropigmentation Procedure: A Useful Procedure for Hair Restoration. *J Craniofac Surg*. 2021 May 1;32(3):1049-1053. doi: 10.1097/SCS.00000000000007208. PMID: 33181616.
6. Traquina AC. Micropigmentation as an adjuvant in cosmetic surgery of the scalp. *Dermatol Surg*. 2001 Feb;27(2):123-8. PMID: 11207683.
7. Park JH, Moh JS, Lee SY, You SH. Micropigmentation: camouflaging scalp alopecia and scars in Korean patients. *Aesthetic Plast Surg*. 2014 Feb;38(1):199-204. doi: 10.1007/s00266-013-0259-1. Epub 2013 Dec 24. PMID: 24366233.
8. Wu DC. Successful Treatment of Scalp Micropigmentation With 1064 nm Picosecond Nd:YAG Laser. *Lasers Surg Med*. 2020 Dec 1. doi: 10.1002/lsm.23364. Epub ahead of print. PMID: 33289086.
9. Rose PT. Advances in Hair Restoration. *Dermatol Clin*. 2018 Jan;36(1):57-62. doi: 10.1016/j.det.2017.09.008. PMID: 29108547.
10. Park JH, Kim N, Kim DW, Lee I. Revision for unsatisfactory outcomes of scalp micropigmentation. *Int J Dermatol*. 2024 Jun 1. doi: 10.1111/ijd.17260. Epub ahead of print. PMID: 38822591.
11. Park JH, Kim N, You SH. Hair transplantation versus scalp micropigmentation: A retrospective study of treatment selection in persons with female pattern hair loss. *J Cosmet Dermatol*. 2024 Oct;23(10):3347-3355. doi: 10.1111/jocd.16429. Epub 2024 Jun 18. PMID: 38894530.
12. Rassman W, Pak J, Kim J. Combining Follicular Unit Extraction and Scalp Micropigmentation for the Cosmetic Treatment of Alopecias. *Plast Reconstr Surg Glob Open*. 2017 Nov 7;5(11):e1420. doi: 10.1097/GOX.0000000000001420. PMID: 29263945; PMCID: PMC5732651.
13. Lampeter W. Editor's invited commentary: Micropigmentation: camouflaging scalp alopecia and scars in Korean patients. *Aesthetic Plast Surg*. 2014 Feb;38(1):205-206. doi: 10.1007/s00266-013-



- 0265-3. Epub 2014 Jan 11. PMID: 24414114.
14. Lahiji SF, Um DJ, Kim Y, Jang J, Yang H, Jung H. Scalp Micro-Pigmentation via Transcutaneous Implantation of Flexible Tissue Interlocking Biodegradable Microneedles. *Pharmaceutics*. 2019 Oct 23;11(11):549. doi: 10.3390/pharmaceutics11110549. PMID: 31652807; PMCID: PMC6920983.
15. Daruwalla SB, Dhurat RS, Hamid SAT. All that a Dermatotr ichologist needs to know about Hair Camouflage: A Comprehensive Review. *Int J Trichology*. 2022 May-Jun;14(3):77-83. doi: 10.4103/ijt.ijt_6_21. Epub 2022 May 24. PMID: 35755961; PMCID: PMC9231527.
16. Saed S, Ibrahim O, Bergfeld WF. Hair camouflage: A comprehensive review. *Int J Womens Dermatol*. 2017 Feb 16;3(1 Suppl):S75-S80. doi: 10.1016/j.ijwd.2017.02.016. PMID: 28492045; PMCID: PMC5419059.
17. Falto-Aizpurua L, Choudhary S, Tosti A. Emerging treatments in alopecia. *Expert Opin Emerg Drugs*. 2014 Dec;19(4):545-56. doi: 10.1517/14728214.2014.974550. Epub 2014 Oct 21. Erratum in: *Expert Opin Emerg Drugs*. 2015 Jun;20(2):347. doi: 10.1517/14728214.2015.1027860 . PMID: 25330928.
18. Saed S, Ibrahim O, Bergfeld WF. Hair camouflage: A comprehensive review. *Int J Womens Dermatol*. 2016 Oct 31;2(4):122-127. doi: 10.1016/j.ijwd.2016.09.002. PMID: 28492024; PMCID: PMC5418894.
19. Dhurat RS, Shanshanwal SJS, Dandale AL. Standardization of SMP Procedure and Its Impact On Outcome. *J Cutan Aesthet Surg*. 2017 Jul-Sep;10(3):145-149. doi: 10.4103/JCAS.JCAS_116_16. PMID: 29403185; PMCID: PMC5782438.
20. Ramnot A, Resnik S, Resnik B. Complete regrowth of hair following scalp tattooing in a patient with alopecia universalis. *JAAD Case Rep*. 2022 Nov 4;31:102-104. doi: 10.1016/j.jdc.2022.10.027. PMID: 36578312; PMCID: PMC9791243.

